

POSTER PRESENTATION

Open Access

Response to HAART according to sex and origin (immigrant vs autochthonous) in a cohort of patients who initiate antiretroviral treatment

JA Perez-Molina^{1*}, M Mora-Rillo², I Suarez-Lozano³, JL Casado⁴, R Teira⁵, P Rivas⁶, E Pedrol⁷, A Hernando⁸, P Domingo⁹, E Barquilla¹⁰, H Esteban¹⁰, J Gonzalez-Garcia¹⁰

From Tenth International Congress on Drug Therapy in HIV Infection
Glasgow, UK. 7-11 November 2010

Purpose

Although poorly studied, gender differences can affect the efficacy of HAART. Immigrant women (IW) may also be at risk of treatment failure due to greater marginalization, cultural differences, or reduced access to health care. This subanalysis examined differences in baseline characteristics and response to HAART according to sex and geographic origin.

Methods

Subanalysis of GES-5808 (retrospective comparative study autochthonous/immigrant patients initiating HAART Jan05-Dec06). Late diagnosis was defined as a CD4+ count ≥ 200 , and/or AIDS at initiation of HAART. The primary endpoint was time to treatment failure (TTF), which was defined as virological failure (VF),

death, opportunistic infection (OI), interruption of HAART, or loss to follow-up. Survival was analyzed using a univariate (Kaplan-Meier) and multivariate (Cox regression) approach.

Results

Patient Characteristics at Initiation of HAART (Table 1)

Conclusions

Response to HAART was similar in both sexes. Men started HAART later and women had higher loss to follow-up and more treatment switches. This was even more common among IW. Earlier diagnosis is necessary for men; measures to improve adherence should be promoted among women, especially IW.

Table 1

	WOMEN (318)	MEN (772)	P
Immigrants, %	45.6	31.1	<0.001
Age, years (IQR)	35 (29-41)	39 (33-44)	<0.001
Median viral load (IQR)/CD4 (IQR)	4.7 (4.2-5.2)/ 217 (113-300)	5.0 (4.5-5.4)/ 190 (69-280)	0.001/0.002
Coinfection with HBV or HCV %	25.2	29.3	0.32
Stage C/Late diagnosis, %	21.2/49.0	29.0/59.0	0.006/0.003
Median time from diagnosis of HIV infection to initiation of HAART, mo (IQR)	15 (2-43)	16 (2-49)	0.55
Median TTF, wk	147	171	<0.001
VF/OI, %	5.3	6.3	0.52

Educational level and occupational status were significantly poorer in women. The adjusted risk of treatment failure in women was not significantly different from that of men (HR, 1.101; 95% CI, 0.79-1.53). The increase in CD4 lymphocytes was equivalent (185 vs 205). TTF was shorter among (IW) than autochthonous women (AW): 124 weeks (95% CI, 64-183) vs 152 (95% CI, 127-174). Most immigrant women were African and Latin American, and their dropout rate (25.5 vs 11.6) was double that of AW.

¹Hospital Ramón y Cajal, Tropical Medicine Unit. Infectious Diseases Dpt, Madrid, Spain

Full list of author information is available at the end of the article

Author details

¹Hospital Ramón y Cajal, Tropical Medicine Unit, Infectious Diseases Dpt., Madrid, Spain. ²Hospital La Paz, Madrid, Spain. ³Cohorte VACH, Huelva, Spain. ⁴Hospital Ramon y Cajal, Madrid, Spain. ⁵Cohorte VACH, Torrelavega, Spain. ⁶Hospital Carlos III, Madrid, Spain. ⁷Cohorte VACH, Tarragona, Spain. ⁸Hospital 12 de Octubre, Madrid, Spain. ⁹Cohorte VACH, Barcelona, Spain. ¹⁰Fundación SEIMC-GESIDA, Madrid, Spain.

Published: 8 November 2010

doi:10.1186/1758-2652-13-S4-P21

Cite this article as: Perez-Molina *et al.*: Response to HAART according to sex and origin (immigrant vs autochthonous) in a cohort of patients who initiate antiretroviral treatment. *Journal of the International AIDS Society* 2010 **13**(Suppl 4):P21.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

