

ORAL PRESENTATION

Open Access

Limits to the scalability of cortical network models

Sacha J van Albada^{1*}, Moritz Helias¹, Markus Diesmann^{1,2,3}

From 24th Annual Computational Neuroscience Meeting: CNS*2015
Prague, Czech Republic. 18-23 July 2015

The size of the mammalian brain is inconveniently right in the middle between a few interacting particles and a mole of matter on a logarithmic scale. In physics, we learn that often in the limit where system size goes to infinity, simple mathematical expressions can be obtained uncovering the mechanisms governing the dynamics at the large but finite system size in nature. In neuroscience, however, we found that such an ansatz may fail because correlations drop so slowly that the mechanism governing the behavior in the infinite size limit [1] is not the mechanism relevant at the scale of the brain circuit in question [2]. The direct simulation of networks at their natural size has historically been difficult due to the sheer number of neurons and synapses. Therefore, neuroscientists also routinely explore the other side of the logarithmic scale and investigate downscaled circuits. In summary, it seems that brain networks are often too small for the infinity limit and too large for simulations.

In this contribution, we assess the scalability of networks in the asynchronous irregular state with a focus on downscaling. By extending the theory of correlations in such networks [2-5] and verifying analytical predictions by direct simulations using NEST [6], we formally demonstrate that generally already second-order measures cannot be preserved. The underlying mathematical reason is a one-to-one mapping between correlation structure and effective connectivity, which depends both on the physical connectivity and on the working point of the neurons [7]. Correlations are relevant because they influence synaptic plasticity [8] and large-scale measurements of neuronal activity [9], and are related to information processing and behavior [10,11].

Our results show that the reducibility of asynchronous networks is fundamentally limited, indicating the importance of considering networks with realistic numbers of neurons and synapses. Fortunately, corresponding simulation technology is becoming available to neuroscience [12]. Both the investigation of the infinity limit and the exploration of downscaled networks remain powerful methods of computational neuroscience. However, researchers should make explicit the rationale they apply in up- or downscaling.

Acknowledgements

We acknowledge funding by the Helmholtz Association: portfolio theme SMHB and Young Investigator's Group VH-NG-1028, and EU Grants 269921 (BrainScaleS) and 604102 (Human Brain Project).

Authors' details

¹Institute of Neuroscience and Medicine (INM-6) and Institute for Advanced Simulation (IAS-6), Jülich Research Centre and JARA, Jülich, Germany.

²Department of Psychiatry, Psychotherapy and Psychosomatics, Medical Faculty, RWTH Aachen University, Aachen, Germany. ³Department of Physics, Faculty 1, RWTH Aachen University, Aachen, Germany.

Published: 18 December 2015

References

1. Renart A, de la Rocha J, Bartho P, Hollender L, Parga N, Reyes A, Harris KD: **The asynchronous state in cortical circuits.** *Science* 2010, **327**(5965):587-590.
2. Helias M, Tetzlaff T, Diesmann M: **The correlation structure of local cortical networks intrinsically results from recurrent dynamics.** *PLoS Comput Biol* 2014, **10**(1):e1003428.
3. Ginzburg I, Sompolinsky H: **Theory of correlations in stochastic neural networks.** *Phys Rev E* 1994, **50**(4):3171-3191.
4. Gritsky D, Tetzlaff T, Diesmann M, Helias M: **A unified view on weakly correlated recurrent networks.** *Front Comput Neurosci* 2013, **7**:131.
5. Helias M, Tetzlaff T, Diesmann M: **Echoes in correlated neural systems.** *New J Phys* 2013, **15**:023002.
6. Gewaltig MO, Diesmann M: **NEST (NEural Simulation Tool).** *Scholarpedia* 2007, **2**:1430.
7. van Albada SJ, Helias M, Diesmann M: **Scalability of asynchronous networks is limited by one-to-one mapping between effective connectivity and correlations.** *arXiv preprint* 2014, **1411**:4770.
8. Morrison A, Aertsen A, Diesmann M: **Spike-timing dependent plasticity in balanced random networks.** *Neural Comput* 2007, **19**(6):1437-1467.

* Correspondence: s.van.albada@fz-juelich.de

¹Institute of Neuroscience and Medicine (INM-6) and Institute for Advanced Simulation (IAS-6), Jülich Research Centre and JARA, Jülich, Germany
Full list of author information is available at the end of the article

9. Lindén H, Tetzlaff T, Potjans TC, Pettersen KH, Grün S, *et al*: **Modeling the spatial reach of the LFP.** *Neuron* 2011, **72**(5):859-872.
10. Zohary E, Shadlen MN, Newsome WT: **Correlated neuronal discharge rate and its implications for psychophysical performance.** *Nature* 1994, **370**(6485):140-143.
11. Riehle A, Grün S, Diesmann M, Aertsen A: **Spike synchronization and rate modulation differentially involved in motor cortical function.** *Science* 1997, **278**(5345):1950-1953.
12. Kunkel S, Schmidt M, Eppler JM, Plesser HE, Masumoto G, *et al*: **Spiking network simulation code for petascale computers.** *Front Neuroinform* 2014, **8**:78.

doi:10.1186/1471-2202-16-S1-O1

Cite this article as: van Albada *et al.*: Limits to the scalability of cortical network models. *BMC Neuroscience* 2015 **16**(Suppl 1):O1.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

