



## ORIGINAL ARTICLE

# Normative scores for the Timed Up & Go in a Spanish sample of community-dweller adults with preserved functionality



Arturo X. Pereiro<sup>a,\*</sup>, María Campos-Magdaleno<sup>a</sup>, Esperanza Navarro-Pardo<sup>b</sup>,  
Onésimo Juncos-Rabadán<sup>a</sup>, David Facal<sup>a</sup>

<sup>a</sup> Faculty of Psychology, University of Santiago de Compostela, Santiago de Compostela, Spain

<sup>b</sup> Faculty of Psychology, University of Valencia, Valencia, Spain

Received 4 June 2020; accepted 26 February 2021

## KEYWORDS

Aging;  
Ambulation;  
Functionality;  
Normative scores;  
Timed-Up & Go test

**Abstract** Our aim was to calculate the 'Timed Up & Go' (TUG) normative scores in a Spanish sample composed of functional older adults. The TUG test provides a measure of global ambulation skills and its total score has been successfully related with functionality and other important health variables in older adults. Reliable norms are needed for adults 50 years and older that allow the early identification and intervention in motor disturbances. The study was carried out with adults from Galicia and Valencia living in the community. A total of 314 Spanish community-living participants, aged from 50 to 90 years and functionality preserved were assessed through the implementation of a cross-sectional design. Health, comorbidity, physical activity, cognitive status, functionality measures and TUG test scores were obtained. TUG scores were successfully predicted by age and gender, and significantly correlated with cognitive status and comorbidity. TUG norms were calculated by age-group for women and men. TUG normative scores were below 13 s and slightly lower in men. Normative scores for women and men were lower than those proposed in studies carried-out in our context. Our norms showed risk reference scores close to those obtained by meta-analytical procedures.

© 2021 The Authors. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [arturoxose.pereiro@usc.es](mailto:arturoxose.pereiro@usc.es) (A.X. Pereiro).

**PALABRAS CLAVE**

Envejecimiento;  
Deambulaci3n;  
Funcionalidad;  
Puntuaciones  
normativas;  
Test Timed Up & Go

## Puntuaciones normativas para el test Timed Up & Go en una muestra comunitaria espa~ola de adultos con funcionalidad preservada

**Resumen** Conocer las puntuaciones normativas del test Timed Up & Go (TUG) sobre una muestra de adultos mayores con funcionalidad preservada. El test TUG proporciona una medida global de deambulaci3n, y su puntuaci3n total se ha relaciona con medidas de funcionalidad y otras relacionadas con la salud en las personas mayores. Conviene disponer de valores normativos para adultos a partir de los 50 a~os que permitan realizar una identificaci3n e intervenci3n tempranas en las dificultades motrices. El estudio se llev3 a cabo con personas adultas de Galicia y Valencia que viven en la comunidad. Un total de 314 participantes mayores, de entre 50 y 90 a~os, que vivían en la comunidad fueron evaluadas implementando un dise~o transversal. Se obtuvieron medidas de salud, comorbilidad, actividad fí sica, cognici3n, funcionalidad, adem3s de las puntuaciones en el test TUG. Edad y g3nero predijeron de modo significativo la puntuaci3n total en el TUG que, adem3s, estableci3 correlaciones significativas con estatus cognitivo y comorbilidad. Se calcularon los valores normativos para el TUG por grupo de edad para varones y mujeres. Las puntuaciones normativas estuvieron por debajo de los 13 s, y fueron ligeramente inferiores en los varones. Las puntuaciones normativas estuvieron, en ambos g3neros, por debajo de los valores propuestos en estudios previos realizados con poblaci3n espa~ola. Nuestros valores normativos ofrecen puntuaciones de referencia para la detecci3n de riesgo semejantes a las obtenidas por procedimientos meta-analíticos.

© 2021 Los Autores. Publicado por Elsevier Espa~a, S.L.U. Este es un artí culo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Ambulation is a core motor skill for maintaining basic levels of functionality<sup>1</sup> and quality of life<sup>2</sup> in older adults. Impairs in global mobility, as other physical condition measures like grip strength or balance have been shown negatively associated to the ability to perform activities of daily living, and accurately predict morbidity and mortality outcomes in old age.<sup>3,4</sup> It is estimated that between 20 and 50% of adults over 65 suffer motor impairments that affects ambulation.<sup>5,6</sup>

Podsiadlo and Richardson<sup>7</sup> suggested a modification of the ‘‘Get-up-and-go’ test<sup>8</sup> where the score is the time spent while the subject is standing from an armchair, walking 3 meters, returning to the chair walking 3 additional meters and sitting down. This test, named ‘Timed Up & Go’ (TUG) involve motor skills to transfer (standing-up and sitting-down), ambulation, and turning, and it is considered an easy and simple tool to assess mobility in older adults.<sup>9</sup> Reliable, valid, cost-effective, and safe measures to evaluate overall functional mobility, as those provide by the TUG, are needed for the early identification of these difficulties that could be potentially reversed through training interventions.<sup>10,11</sup>

Suitable values for reliability and validity levels were reported for the TUG.<sup>11</sup> Even though TUG score did not seem to successful predict the risk of falling in older adults,<sup>12</sup> they were significantly associated to balance, general health, cognitive status or physical activity measures<sup>13–16</sup> and have been shown to be sensitive when used for frailty identification in older adults.<sup>17,18</sup> Available evidence pointed out that TUG scores significantly change by age, and mixed evidence were reported regarding differences by gender.<sup>11,14,19,20</sup>

TUG norms by age and gender for Spanish population were reported in two studies.<sup>14,19</sup> Participants in both studies were 70 years and older and were selected by stratified

sampling procedures among the community-dwelling population of two Spanish cities. Pondal and Del Ser<sup>14</sup> excluded 41.5% from a total sample of 627 participants arguing impossibility to do the test or medical conditions that interfere in motor performance. In Abizanda et al. study,<sup>19</sup> 201 (25.8%) participants were excluded from a total sample of 781 considered but specific reasons for exclusion were not reported in this subsample. In both studies, norms were reported in percentiles for adults older than 70 although different age groups were considered. Thus, Abizanda et al.<sup>19</sup> reported norms for 70–79 and >80 age groups meanwhile Pondal and Del Ser<sup>14</sup> reported normative values for 71–75, 76–79, 80–85, and 86–99 age groups. The high exclusion rates observed in both studies<sup>14,19</sup> suggest that, according to the high average age of the samples, there had a high prevalence of difficulties in global mobility.

In general, TUG scores in Abizanda et al.<sup>19</sup> are considerably higher than in Pondal and Del Ser.<sup>14</sup> Thus, Abizanda et al.<sup>19</sup> normative scores for participants in their eighties and older (men: 25.9 s; women: 22.9 s) were considerably higher than those reported by Pondal and Del Ser<sup>14</sup> for the 81–85 (men: 14.0 s; women: 17.6 s) and 86–99 (men: 18.2 s; women: 19.6 s) age groups in a comparable percentile (e.g., percentile equality to 1.28 SD above the mean). Similarly, Abizanda et al.<sup>19</sup> scores corresponding to the percentile equality to 1.28 SD above the mean for the age groups 70–79 (men: 14.1 s; women: 16.5 s) and >80 (men: 25.9 s; women: 22.9 s) were considerably higher than the cut-off points suggested by Bohannon<sup>11</sup> after meta-analysis of 21 studies for participants in the 70–79 (10.2 s) and 80–99 (12.7 s) age groups.

Abizanda et al.<sup>19</sup> and Pondal and Del Ser<sup>14</sup> studies reported TUG norms for adults older than 70 and it impedes an early detection of global mobility impairments in

Spanish younger adults. Further, the inconsistencies between the two norms with respect to age groups make difficult to follow them as standards for clinical and research purposes. Taking into account these criticisms the aims of this paper were: (a) to provide norms for the TUG test in a sample of Spanish adults aged from 50 years, recruited in the community with global mobility preserved without assistive devices, and free of significant cognitive decline, and (b) to know how the scores associate to the percentile rank distribution (3rd, 5th, 10th, 16th, 20th, 50th, 70th, 90th, 95th, and 97th) considering gender and several age groups (i.e., 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, 80–90).

## Methods

### Participants

A total sample of 421 community-dwelling participants was incidentally recruited in the Galicia and Valencia, two autonomous regions of Spain. Galicia (total surface area, 29,575 km<sup>2</sup>; population, 2,701,743) and Valencia (total surface area, 23,255 km<sup>2</sup>; population, 5,000,868) are located, respectively, in the northwest and the southeast of Spain. Criteria for the recruitment of the participants were: older adults aged 50 or more years, living in the community, non-severe functional impairment in lower limbs, independent ambulation (i.e., participants with assistive devices for walking were excluded), no previous diagnosis of neurological or psychiatric diseases including dementia, cognitive status above the cut-off considering norms for age and education level, and free from visual disturbance.

Community-dwelling participants were recruited from a large on-going study on cognitive aging being undertaken at the University of Santiago de Compostela and University of Valencia. Candidates were relatives or neighbors of university students and were invited to participate in the study when active life and the autonomy for the instrumental activities of daily life were maintained. The participants were evaluated in their own homes and received no incentives for their collaboration in the study.

Through the implementation of a cross-sectional design, seven age groups (50–54, 55–59, 60–64, 65–69, 70–74, 75–79, and 80–90) were established and gender of participants was considered in recruitment procedures to yield a balanced sample.

A 5.9% of the total sample (nine women and five men) was excluded because moderate functional disturbances and 18.5% (eighty-three participants) as a consequence of low cognitive status (see *Procedure* section for more specifications). In addition, outlier TUG scores were identified and excluded using the stem-and-leaf plots for each age-group (age-group 50–54: 1 participant; age-group 70–74: 3 participants; age-group 75–79: 2 participants; age-group 80–90: 4 participants). Age range was 50–90 ( $M = 68.65$ ;  $SD = 8.23$ ) and 57.5% of the sample participants were female. The norms for a final sample consisted of 314 participants were calculated. Descriptive statistics for age, functionality, cognitive status, health, and physical activity are shown in [Table 1](#).

### Procedure

Assessment were carried-out by ten interviewers trained in neuropsychological evaluation of older adults. A socio-demographic and health questionnaire, the Lawton Instrumental Activities of Daily Living (IADL-Lawton) Scale<sup>21</sup> (Spanish version of Vergara et al.<sup>22</sup>), the TUG test,<sup>7</sup> the Montreal Cognitive Assessment<sup>23</sup> (MoCA), the Minnesota Leisure Time Physical Activity Questionnaire<sup>24</sup> (VREM; Spanish version Ruiz et al.<sup>25</sup>), the Charlson Comorbidity Index<sup>26</sup> (CCI), and the General Health Questionnaire<sup>27</sup> (GHQ-12; Spanish version Rocha et al.<sup>28</sup>) were administered, following this order, to the participants in a 90-minute session.

Informed consent as required in the Declaration of Helsinki on research ethics was obtained from all participants. The research plan was approved by both ethics committees (the Clinical Research Ethics Committee of the Xunta de Galicia – Procedure number 2018/620), and by the Commission of Ethics in Experimental Research of the University of Valencia - Procedure number H1521026499251).

The TUG test was administered using an armless chair (between 45 and 47 cm height) and immediately aborted if the participant was at risk of falling while the standing-up transfer was performed.

Women with scores below 6 points and men with scores below 5 points in the IADL-Lawton were excluded in order to obtain norms from participants with preserved functionality. Similarly, participants with significant cognitive decline were excluded when the MoCA score was  $-1.88SD$  below the mean (3rd percentile) established for their age and education reference group.<sup>29</sup>

### Statistical analysis

SPSS package, version 20.0 (SPSS, Chicago, IL USA), was employed for the statistical analyses.

The enter method of Multiple linear regression analysis was carried out to examine the predictive relationship of age (in years) and gender on the TUG total score. Tolerance, Variance inflation factor (VIF) and Condition index statistics were calculated to test collinearity, and the coefficient of determination ( $R^2$ ) was obtained to estimate the effect size.

In order to obtain the exact percentile for an exact score, the procedure used by Van der Elst et al.<sup>30</sup> was applied. The predicted value was calculated separately for men and women using the following regression equation where  $\hat{y}_i$  corresponds to the TUG test total score ( $\hat{y}_i = b_0 + b_1 \cdot \text{age}$ ). Then, the residual value was calculated ( $c_i = y_i - \hat{y}_i$ ) and later standardized by z score transformation (i.e. dividing the residual value by the standard deviation for the residual obtained in the fitted regression model;  $z = e/i$ ). Finally, the accumulated probability associated with this z value was obtained using the standardized normal distribution to establish the normal values corresponding for the percentiles considered (i.e. 3rd, 5th, 10th, 16th, 20th, 50th, 70th, 90th, 95th, and 97th) in age-groups (i.e. 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, and 80–90).

**Table 1** Sample description values (Mean, Standard Deviation) for age, functionality (IADL-Lawton), cognitive status (MoCA), health (GHQ-12), comorbidity (CCI), and physical activity (VREM).

Age group	N	Age (M, SD)	IADL-Lawton (M, SD)	MoCA (M, SD)	GHQ-12 (M, SD)	CCI (M, SD)	VREM
50–54	15	51.73 (1.48)	7.67 (.71)	25.00 (2.97)	18.46 (5.27)	1.33 (.72)	Low-moderate: 26.66% High-very high: 73.33%
55–59	10	57.20 (1.47)	7.20 (.91)	23.50 (2.71)	21.20 (4.56)	1.30 (.48)	Low-moderate: 30% High-very high: 70%
60–64	80	61.53 (1.36)	7.70 (.62)	26.06 (2.88)	13.21 (6.02)	.70 (.98)	Low-moderate: 11.25% High-very high: 88.75%
65–69	73	67.05 (1.47)	7.81 (.49)	26.23 (2.85)	12.02 (5.45)	.64 (.85)	Low-moderate: 4.10% High-very high: 95.89%
70–74	56	72.12 (1.30)	7.64 (.64)	24.48 (3.03)	12.19 (5.99)	.63 (.84)	Low-moderate: 5.35% High-very high: 94.64%
75–79	46	76.93 (1.52)	7.57 (.68)	24.41 (2.89)	13.97 (6.55)	1.39 (1.68)	Low-moderate: 6.52% High-very high: 93.47%
80–90	34	82.73 (2.68)	7.44 (.70)	23.50 (3.00)	11.23 (4.50)	.82 (1.46)	Low-moderate: 5.88% High-very high: 94.11%

Note: IADL-Lawton: Lawton Instrumental Activities of Daily Living Scale; MoCA: Montreal Cognitive Assessment; GHQ-12: General Health Questionnaire; CCI: Charlson Comorbidity Index; VREM: The Minnesota Leisure Time Physical Activity Questionnaire; M = mean; SD = Standard Deviation.

## Results

The TUG total scores were not normally distributed ( $W_{Shapiro-Wilk's}$  (313) = .977,  $p < .001$ ). The distribution was slightly positively skewed (Skewness = .520; Standard error = .138) and adopted a soft platykurtic shape (Kurtosis = .378; Standard error = .275).

TUG total score was significantly correlated with MoCA ( $r = -.29$ ;  $p < .001$ ) and CCI ( $r = .11$ ;  $p = .039$ ) scores. Significant relationships were not observed between TUG total score and AIDL-Lawton, GHQ-12 and VREM scores.

TUG scores were successfully predicted ( $R^2 = .13$ ) by age ( $\beta = .098$ ;  $SE \beta = .016$ ;  $Standardized \beta = .336$ ;  $t = 6.34$ ,  $p < .001$ ) and gender ( $\beta = .648$ ;  $SE \beta = .257$ ;  $Standardized \beta = .134$ ;  $t = 2.52$ ,  $p = .012$ ). Colinearity statistics ( $Tolerance = .99$ ,  $VIF = 1.002$ , and  $Condition\ index = 20.66$ ) showed a very slight overlap between the predictive variables age and gender.

Norms for women (see Table 2) and men (see Table 3) were calculated to know how the scores associate to the percentile rank distribution (i.e. 3rd, 5th, 10th, 16th, 20th, 50th, 70th, 90th, 95th, and 97th) in each age group (50–54, 55–59, 60–64, 65–69, 70–74, 75–79, and 80–90). The correspondence between 97th, 95th, and 90th percentiles and,

respectively, 1.88, 1.64, and 1.28 SDs above the mean must be considered for less or more restrictive use of these norms for clinical purposes.

The TUG normative scores were, for women and men, always below 13 s and, generally, women were slightly slower than men.

## Discussion

TUG scores showed significant relationships with cognitive status and comorbidity. Although previous studies showed significant associations with functionality, health and physical activity measures,<sup>13–16</sup> we consider that these relationships were not found in our study because the ceiling effect as a consequence of the preserved functionality of the sample. The TUG score was successfully predicted by age<sup>11,14,20</sup> and in line with some previous studies,<sup>14,19,20</sup> gender was revealed as a predictive variable of TUG test scores, showing that women get higher scores in TUG test and therefore generally are affected by a greater slowing in motor performance.

Our normative scores for women and men were considerably below those reported by Abizanda et al.<sup>19</sup> and also slightly lower than those proposed by Pondal and Del Ser.<sup>14</sup>

**Table 2** TUG normative scores (in s) for women by age group.

Z score	Percentile	Age groups						
		50–54	55–69	60–64	65–69	70–74	75–79	80–90
1.88	97	9.35	9.80	10.24	10.69	11.13	11.58	12.02
1.64	95	9.11	9.56	10.00	10.45	10.89	11.34	11.78
1.28	90	8.76	9.20	9.65	10.09	10.54	10.98	11.43
0.52	70	8.00	8.44	8.89	9.33	9.78	10.22	10.67
0	50	7.48	7.92	8.37	8.81	9.26	9.70	10.15
–0.84	20	6.64	7.09	7.53	7.98	8.42	8.87	9.31
–1	16	6.48	6.93	7.37	7.82	8.26	8.71	9.15
–1.28	10	6.20	6.65	7.09	7.54	7.98	8.43	8.87
–1.64	5	5.84	6.29	6.73	7.18	7.62	8.07	8.51
–1.88	3	5.60	6.05	6.49	6.94	7.38	7.83	8.27

**Table 3** TUG normative scores (in s) for men by age group.

Z score	Percentile	Age groups						
		50–54	55–59	60–64	65–69	70–74	75–79	80–90
1.88	97	8.32	8.87	9.42	9.97	10.52	11.07	11.62
1.64	95	8.08	8.63	9.18	9.73	10.28	10.83	11.38
1.28	90	7.72	8.27	8.82	9.37	9.92	10.47	11.02
0.52	70	6.96	7.51	8.06	8.61	9.16	9.71	10.26
0	50	6.45	7.00	7.55	8.10	8.65	9.20	9.75
–0.84	20	5.61	6.16	6.71	7.26	7.81	8.36	8.91
–1	16	5.45	6.00	6.55	7.10	7.65	8.20	8.75
–1.28	10	5.17	5.72	6.27	6.82	7.37	7.92	8.47
–1.64	5	4.81	5.36	5.91	6.46	7.01	7.56	8.11
–1.88	3	4.57	5.12	5.67	6.22	6.77	7.32	7.87

Thus, for a comparable age group and 90th percentile (i.e., equality to 1.28 SD above the mean), our results pointed out as reference scores 11.43 for women and 11.02 for men, much lower than those proposed by Abizanda et al.<sup>19</sup> for the 80–90 age-group (men: 25.9 s; women: 22.9 s), and also below those reported by Pondal and del Ser<sup>14</sup> for the 81–85 age group (men: 14.0 s; women: 17.6 s). A healthier sample with higher levels of functionality in the participants of our study may explain this disparity in the results.

Our norms were closer to the reference values for older adult population reported by Bohannon<sup>11</sup> in his meta-analysis. Thus, considering the 97th percentile (equally to 1.88 SD above the mean), the reference scores found in our study (65–69 age group: 10.69 for women and 9.97 for men; 75–79 age group: 11.58 for women and 11.07 for men; 80–90 age group: 12.02 for women and 11.62 for men), much more close to the cut-off points considering the upper limit of the interval coefficient established in the Bohannon<sup>11</sup> study (60–69 age group: 9.0; 70–79 age group: 10.2; 80–99 age group: 12.7). In conclusion, TUG scores showed significant relationships with cognitive status and comorbidity, even on preserved functionality people. The TUG score was higher as age increases and in women compared to men. Our norms were closer to the reference values for older adult population obtained by meta-analytical procedures.

Abizanda et al.<sup>19</sup> and Pondal and Del Ser<sup>14</sup> studies reported TUG norms for adults older than 70 and it impedes an early detection of global mobility impairments in Spanish younger adults. We hope that the TUG's norms for older adults from the age of 50 and without significant functional impairments will be helpful for the early identification of frailty in older adults.<sup>17,18</sup>

Future studies should implement representative sampling procedures to establish norms for the TUG test and longitudinally test the predictive value of the TUG scores for the functional outcomes.

## Funding

This work was financially supported through FEDER funds by the National Research Agency (Spanish' Ministry of Science, Innovation and Universities) (Project Ref. PSI2017-89389-C2-1-R), the Galician Government (Consellería de Cultura, Educación e Ordenación Universitaria; axudas para a consolidación e estruturación de unidades de investigación competitivas do Sistema Universitario de Galicia; GI-1807-USC: Ref. ED431-2017/27), and by the Frailty Network IN607C 2016/08. The funders played no role in the designing, conducting, or reporting of this study.

## Conflicts of interest

Not applicable.

## References

- Ferrucci L, Cooper R, Shardell M, Simonsick EM, Schrack JA, Kuh D. Age-related change in mobility: perspectives from life course epidemiology and geroscience. *J Gerontol Ser A Biol Sci Med Sci*. 2016;71:1184–94, <http://dx.doi.org/10.1093/gerona/glw043>.
- Webber SC, Porter MM, Menec VH. Mobility in older adults: a comprehensive framework. *Gerontologist*. 2010;50:443–50, <http://dx.doi.org/10.1093/geront/gnq013>.
- Cooper R, Kuh D, Hardy R. Mortality Review Group, FALCon and HALCyon Study Teams. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *BMJ*. 2010;341:c4467, <http://dx.doi.org/10.1136/bmj.c4467>.
- Cooper R, Kuh D, Cooper C, Gale CR, Lawlor DA, Matthews F, et al., FALCon and HALCyon Study Teams. Objective measures of physical capability and subsequent health: a systematic review. *Age Ageing*. 2011;40:14–23, <http://dx.doi.org/10.1093/ageing/afq117>.
- Guralnik JM, Ferrucci L, Pieper CF. Lower extremity function and subsequent disability: consistency across studies, predictive models, and value of gait speed alone compared with the short physical performance battery. *J Gerontol Ser A Biol Sci Med Sci*. 2000;55:221–31, <http://dx.doi.org/10.1093/gerona/55.4.m221>.
- Shumway-Cook A, Ciol MA, Yorkston KM. Mobility limitations in the medicare population: prevalence and sociodemographic and clinical correlates. *J Am Geriatr Soc*. 2005;53:1217–21, <http://dx.doi.org/10.1111/j.1532-5415.2005.53372.x>.
- Podsiadlo D, Richardson S. The Timed Up & Go: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39:142–8, <http://dx.doi.org/10.1111/j.1532-5415.1991.tb01616.x>.
- Mathias S, Nayak USL, Isaacs B. Balance in the elderly patient: the “Get-up and Go” test. *Arch Phys Med Rehabil*. 1986;67:387–9.
- Chung J, Demiris G, Thompson HJ. Instruments to assess mobility limitation in community-dwelling older adults: a systematic review. *J Aging Phys Act*. 2015;23:298–313, <http://dx.doi.org/10.1123/japa.2013-0181>.
- Brach JS, Vanswearingen JM. Interventions to improve walking in older adults. *Curr Transl Geriatr Exp Gerontol Rep*. 2013;2, <http://dx.doi.org/10.1007/s13670-013-0059-0>.
- Bohannon RW. Reference values for the Timed Up and Go Test: a descriptive meta-analysis. *J Geriatr Phys Ther*. 2006;29:64–8, <http://dx.doi.org/10.1519/00139143-200608000-00004>.
- Barry E, Galvin R, Keogh C, Horgan F, Fahey T. Is the Timed Up and Go test a useful predictor of risk of falls in community dwelling older adults: a systematic review and meta-analysis. *BMC Geriatr*. 2014;14:14, <http://dx.doi.org/10.1186/1471-2318-14-14>.
- Schoene D, Wu SM, Mikolaizak AS, Menant JC, Smith ST, Delbaere K, et al. Discriminative ability and predictive validity of the Timed Up and Go Test in identifying older people who fall: systematic review and meta-analysis. *J Am Geriatr Soc*. 2013;61:202–8, <http://dx.doi.org/10.1111/jgs.12106>.
- Pondal M, del Ser T. Normative data and determinants for the Timed “Up and Go” test in a population based sample of elderly individuals without gait disturbances. *J Geriatr Phys Ther*. 2008;31:57–63.
- Rockwood K, Awalt E, Carver D, MacKnight C. Feasibility and measurement properties of functional reach and the Timed Up and Go test in the Canadian Study of Health and Aging. *J Gerontol*. 2000;55:70–3, <http://dx.doi.org/10.1093/gerona/55.2.m70>.
- Viccaro LJ, Perera S, Studenski SA. Is timed up and go better than gait speed in predicting health, function, and falls in older adults? *J Am Geriatr Soc*. 2011;59:887–92, <http://dx.doi.org/10.1111/j.1532-5415.2011.03336.x>.
- Clegg A, Rogers L, Young J. Diagnostic test accuracy of simple instrument for identifying frailty in community-dwelling older people: a systematic review. *Age Ageing*. 2015;44:148–52, <http://dx.doi.org/10.1093/ageing/afu157>.
- Savva GM, Donoghue OA, Horgan F, O’Regan C, Cronin H, Kenny RA. Using timed up-and-go to identify frail members of the older population. *J Gerontol A Biol Sci Med Sci*. 2013;68:441–6, <http://dx.doi.org/10.1093/gerona/gls190>.
- Abizanda P, López-Torres J, Romero L, Sánchez PM, García I, Esquinas JL. Valores normativos de instrumentos de valoración funcional en ancianos españoles: estudio FRADEA. *Aten Primaria*. 2012;44:162–71, <http://dx.doi.org/10.1016/j.aprim.2011.02.007>.
- Steffen TM, Hacker TA, Mollinger L. Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and gait speeds. *Phys Ther*. 2002;82:128–37, <http://dx.doi.org/10.1093/ptj/82.2.128>.
- Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969;9:179–86, <http://dx.doi.org/10.1093/geront/9.3.Part.1>.
- Vergara I, Bilbao A, Orive M, García-Gutierrez S, Navarro G, Quintana JM. Validation of the Spanish version of the Lawton IADL Scale for its application in elderly people. *Health Qual Life Out*. 2012;10:130, <http://dx.doi.org/10.1186/1477-7525-10-130>.
- Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc*. 2005;53:695–9, <http://dx.doi.org/10.1111/j.1532-5415.2005.53221.x>.
- Taylor HL, Jacobs DR Jr, Schucker B, Knudsen J, Leon AS, Debacker G. A questionnaire for the assessment of leisure time physical activities. *J Chronic Dis*. 1978;3:741–55, [http://dx.doi.org/10.1016/0021-9681\(78\)90058-9](http://dx.doi.org/10.1016/0021-9681(78)90058-9).
- Ruiz Comellas A, Pera G, Baena Díez JM, Mundet Tudurí X, Alzamora Sas T, Elosua R, et al. Validation of a Spanish Short Version of the Minnesota Leisure Time Physical Activity Questionnaire (VREM). *Rev Esp Salud Publica*. 2012;86:495–508, <http://dx.doi.org/10.4321/S1135-57272012000500004>.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic in longitudinal studies: development and validation. *J Chron Dis*. 1987;40:373–83, [http://dx.doi.org/10.1016/0021-9681\(87\)90171-8](http://dx.doi.org/10.1016/0021-9681(87)90171-8).
- Goldberg D, Williams P. *A user’s guide to the General Health Questionnaire*. Windsor, UK: NFER-Nelson; 1988.
- Rocha KB, Pérez K, Rodríguez-Sanz M, Borrell C, Obiols JE. Propiedades psicométricas y valores normativos del General Health Questionnaire (GHQ-12) en población general española. *Int J Clin Hlth Psych*. 2011;11:125–39.
- Pereiro AX, Ramos-Lema S, Lojo-Seoane C, Guardà-Olmos J, Facal D, Juncos-Rabadán O. Correction to: Normative data for the Montreal Cognitive Assessment (MOCA) in a Spanish sample of community-dweller adults. *Eur Geriatr Med*. 2018;9:903–4, <http://dx.doi.org/10.1007/s41999-018-0116-5>.
- Van der Elst W, Dekker S, Hurks P, Jolles J. The letter digit substitution test: demographic influences and regression-based normative data for school-aged children. *Arch Clin Neuropsychol*. 2012;27:433–9, <http://dx.doi.org/10.1093/arclin/acs045>.