

# Reactivation of hepatitis B virus infection in rheumatic diseases: risk and management considerations

Christos Koutsianas, Konstantinos Thomas and Dimitrios Vassilopoulos 

**Abstract:** In patients with rheumatic diseases undergoing immunosuppressive treatment, hepatitis B virus reactivation (HBVr) has been long recognized as a major treatment-related adverse event with substantial morbidity and mortality. Because HBVr is easily preventable with appropriate screening and monitoring strategies, and, when indicated, prophylactic antiviral treatment, awareness of this complication is of the utmost importance, especially in the era of biologic treatments. As a condition, it continues to be topical, in view of the emergence of novel classes of immunosuppressive drugs (i.e. Janus kinase inhibitors) acquiring licenses for a variety of rheumatic diseases. The class-specific risk of these agents for HBVr has not yet been determined. Moreover, ambiguity still exists for the management of patients planned to be treated with traditional agents, such as cyclophosphamide and glucocorticoids, particularly in the setting of resolved HBV infection. Clinicians in the field of rheumatic diseases should be tailoring their practice according to the host's profile and treatment-specific risk for HBVr. In this review, the authors attempt to critically review the existing literature and provide practical advice on these issues.

**Keywords:** hepatitis B virus, reactivation, rheumatic disease, screening, biologics, DMARD, immunosuppression

Received: 30 October 2019; revised manuscript accepted: 11 February 2020.

## Introduction

Hepatitis B virus (HBV) infection still poses a significant worldwide morbidity and mortality burden, despite the availability of an efficacious vaccine and antiviral treatment. The global prevalence of chronic HBV infection has been recently estimated at 3.61%,<sup>1</sup> and, although it seems to be in decline, approximately 887,000 deaths were attributed to HBV complications in 2015.<sup>2</sup> There is substantial geographical variance in the rates of chronic HBV infection, which is highly endemic in the Western Pacific (6.2%) and African (6.1%) areas, but less common in Europe (1.6%) and America (0.7%).<sup>2</sup> However, the streams of immigration and influx of refugees to Western Europe and North America is likely to be changing the prevalence and endemicity in these latter regions, which underlines the need for physicians to be familiar with the disease.

Most relevant studies suggest that the prevalence of HBV in patients with rheumatic disease follows

the pattern of the general population. Chronic HBV infection rates in rheumatoid arthritis (RA) and ankylosing spondylitis (AS) have been estimated at 3.0% and 3.5%, respectively, whereas resolved HBV infection rate ranges between 13% and 50%.<sup>3–5</sup>

HBV is a partially double-stranded DNA virus that is transmitted *via* the parenteral route, and can cause acute or chronic infection. The risk of chronicity is dependent mainly on the age of the host at the time of infection, as the vast majority of infants (>90%) and only <5% of adults develop chronic HBV infection after exposure. Distinguishing between acute, chronic, past (resolved), and occult infection lies in HBV serological markers and measurement of serum HBV-DNA (Table 1 and Figure 1).<sup>6,7</sup>

HBV is unique in its ability to integrate its viral genome in its host's DNA by forming covalently

*Ther Adv Musculoskel Dis*

2020, Vol. 12: 1–19

DOI: 10.1177/  
1759720X20912646

© The Author(s), 2020.  
Article reuse guidelines:  
sagepub.com/journals-  
permissions

Correspondence to:  
**Dimitrios Vassilopoulos**  
Clinical Immunology-  
Rheumatology Unit, 2nd  
Department of Medicine  
and Laboratory, National  
and Kapodistrian  
University of Athens  
School of Medicine,  
Hippokraton General  
Hospital, 114 Vass.  
Sophias Ave., Athens, 115  
27, Greece  
[dvassilop@med.uoa.gr](mailto:dvassilop@med.uoa.gr)

**Christos Koutsianas**  
Clinical Immunology-  
Rheumatology Unit, 2nd  
Department of Medicine  
and Laboratory, National  
and Kapodistrian  
University of Athens  
School of Medicine,  
Hippokraton General  
Hospital, Athens, Greece

Department of  
Rheumatology, The Dudley  
Group NHS Foundation  
Trust, Russells Hall  
Hospital, Dudley, West  
Midlands, UK

**Konstantinos Thomas**  
Clinical Immunology-  
Rheumatology Unit, 2nd  
Department of Medicine  
and Laboratory, National  
and Kapodistrian  
University of Athens  
School of Medicine,  
Hippokraton General  
Hospital, Athens, Greece

**Table 1.** HBV serology [adapted from Koutsianas and colleagues<sup>5</sup>].

Nomenclature for HBV infection based on biochemistry, serology and viral load						
	Acute HBV infection	Chronic HBV infection			Past infection	
		Chronic hepatitis		Inactive carrier	Resolved infection	Occult infection
		HBeAg (+)	HBeAg (-)			
HBsAg	+	+	+	+	-	-
Anti-HBc (total)	+/-	+	+	+	+	Usually +/-
Anti-HBc IgM	+	-	-	-	-	-
Anti-HBs	-	-	-	-	+/-	+/-
HBeAg	+	+	-	-	-	-
Anti-HBe	-	-	+	+/-	+	+/-
Serum HBV-DNA	Variable <sup>a</sup>	Usually >20,000 IU/mL <sup>b</sup>	Usually >2000 IU/mL <sup>b</sup>	Undetectable	Undetectable	Detectable (low levels)
ALT	Markedly elevated	Elevated	Elevated	Normal	Normal	Normal

<sup>a</sup>HBV-DNA levels are usually elevated, but may vary depending on the host's immune response and the exact timing of the test performed.  
<sup>b</sup>The 2,000 and 20,000 IU/mL cut-offs are arbitrary values deriving from the detection limits of historical non-PCR methods of measurement. As HBV-related complications have been identified in patients with lower HBV-DNA levels, it is important to interpret HBV-DNA taking into consideration other factors [age, duration, ALT, disease stage].  
 ALT, alanine aminotransferase; anti-HBc, antibody against hepatitis B core antigen; anti-HBe, antibody against HBeAg; anti-HBs, antibody against HBsAg; HBeAg, hepatitis B envelope antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HBV-DNA, HBV-deoxyribonucleic acid; PCR, polymerase chain reaction.


closed circular DNA (cccDNA), which is responsible for its persistent presence in hepatocyte nuclei.<sup>8</sup> The virus itself does not have a direct cytopathic effect. On the contrary, hepatocellular injury is mediated by innate and adaptive immunity. A strong immune response is associated with viral clearance in acute HBV infection, but is also responsible for hepatocyte damage and fibrosis in the immune active phases of chronic HBV infection.

### Hepatitis B virus reactivation

HBV reactivation (HBVr) is a well-recognized complication of immunosuppressive treatment in cancer, rheumatic diseases, and organ transplantation.<sup>9</sup> It typically occurs in patients with chronic HBV infection [Hepatitis B surface Antigen (HBsAg) positive], but, less commonly, it can complicate immunosuppressive treatment in patients with resolved HBV infection [HBsAg negative/antibody against Hepatitis B core antigen (anti-HBc) positive]. HBVr is usually defined as: a rise in serum HBV-DNA compared with

baseline levels or detection of HBV DNA if undetectable at baseline, or reverse seroconversion (re-emergence) of HBsAg from negative to positive (see Table 2 for full definitions).<sup>10,11</sup> Factors that influence the risk of HBVr are related to the patient, the virus, and the type and duration of immunosuppression used (Table 3).<sup>12</sup>

With the expected further increase in the use of biologics and new oral targeted synthetic agents [i.e. Janus kinase (JAK) inhibitors] in treating rheumatic diseases, rheumatologists should be aware of the potential risk, the screening recommendations, and management options for prophylaxis and monitoring for HBVr.<sup>15</sup> The majority of the literature data around HBVr derives from the fields of hematology and oncology, where combinations of biologic and cytotoxic agents with higher immunosuppressive potential are routinely used, usually for a short period of time (months). These data need to be interpreted with caution when applied to rheumatic diseases, where the potency of immunosuppression is lower and the duration of therapy longer (years).



**Figure 1.** Serologic screening for HBV and interpretation of results.

anti-HBc, antibody against hepatitis B core antigen; anti-HBs, antibody against hepatitis B surface antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus.

**Table 2.** HBV reactivation definitions (literature data).<sup>6,10</sup>

Definition	Population	Baseline serum HBV DNA	Change in virological, serological or biochemical markers during therapy
Virological	HBsAg (+) Anti-HBc (+)	(+)	↑ of HBV DNA by $\geq 2 \log_{10}$ (100-fold)
		(-)	HBV-DNA: $\geq 1000 \text{ IU/mL}$ (3 logs)
		Unknown	HBV-DNA: $\geq 10,000 \text{ IU/mL}$ (4 logs)
	HBsAg (-) Anti-HBc (+)	(-)	Detectable HBV DNA (+)
Serological 'reverse seroconversion'	HBsAg (-) Anti-HBc (+)		HBsAg (+)
Biochemical 'hepatitis flare'	All		↑ of ALT by $\geq 3\times$ from baseline levels + ALT $\geq 100 \text{ IU/mL}$ + No other explanation possible

ALT, alanine aminotransferase; anti-HBc, antibody against hepatitis B core antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus.

In a recent meta-analysis for the prevalence of HBVr in rheumatic diseases being treated with conventional or biologic agents, the pooled HBVr estimate was 1.4%, thus much lower than the relevant estimation in hematology/oncology.<sup>16</sup>

In recent years, there has been an effort to stratify HBVr risk according to the patient's serological status and the type and duration of the immunosuppressive treatment used. The American Gastroenterological Association (AGA) classified

**Table 3.** Risk factors for HBV reactivation (literature data).<sup>12–14</sup>

Host factors	Male sex Older age Cirrhosis Type of disease needing immunosuppression (lymphoma) Co-infection status (e.g. HCV, HIV, HDV)
Virological factors	High baseline HBV-DNA Chronic HBV infection HBeAg positivity Non-A HBV genotype
Immunosuppression	Type Duration (see main text for details)

ALT, alanine aminotransferase; anti-HBc, antibody against hepatitis B core antigen; HBeAg, hepatitis B envelope antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HBV-DNA, HBV-deoxyribonucleic acid; HCV, hepatitis C virus; HDV, hepatitis D virus; HIV, human immunodeficiency virus.

HBVr risk as low (<1%), moderate (1–10%), and high (>10%) based on the above factors.<sup>17</sup> However, the emergence of new agents and the long duration of biologic treatments, especially in rheumatic diseases where there usually are no set stopping rules for treatment, pose further controversy on HBVr management and monitoring guidance.

In the current paper, we will critically review the existing literature on HBVr risk in patients with rheumatic diseases treated with different biologic and nonbiologic agents, and provide practical advice for its prevention and management.

### Type of immunosuppression and HBVr risk and management

#### Glucocorticoids

Glucocorticoid (GC) use in patients with HBV infection was known to relate to HBVr as early as the 1970s,<sup>18</sup> although its reported incidence varies significantly in the published literature (4–50%). The proposed pathophysiological mechanism seems to be twofold: GCs suppress T-cell cytotoxic function, thus diminishing the host's immune check on the virus, but also directly stimulate HBV-DNA replication by activating a GC-responsive transcriptional regulatory element in the HBV genome.<sup>17,19</sup>

There is robust literature evidence that even short courses of more than moderate-dose GC treatment in chronic HBV infection can lead to HBVr.<sup>17,20–22</sup> Chemotherapy for chronically

HBV-infected patients with lymphoma was less likely to cause HBVr in 9 months if it did not contain GCs (38% versus 73% if containing prednisolone).<sup>23</sup> An HBVr incidence rate of 6.1% has been reported in chronic HBV patients with asthma or chronic obstructive pulmonary disease (COPD) receiving GCs. The risk was higher among patients receiving systemic GCs, especially when they were used continuously (for at least 3 months) and in medium to high (>20 mg/day) doses.<sup>24</sup> Patients with chronic HBV infection and rheumatic diseases are also at risk for HBVr and hepatic flare when treated with GCs. For patients not receiving antiviral prophylaxis, and especially when GCs are used in combination with conventional and biologic agents, the hazard ratio (HR) has been reported at 5.1.<sup>25</sup> Peak doses of >40 mg prednisolone equivalents were associated with an adjusted HR for HBVr of 1.64.<sup>26</sup> HBVr risk is higher depending on the patient's HBV status (i.e. chronic hepatitis versus inactive carrier state), on the continuous oral GC use versus intravenous (IV) pulses<sup>27,28</sup> and on the GC dose used.<sup>29,30</sup>

With regards to IV GC pulse therapy in HBsAg positive patients, data are limited and confounded by the concomitant use of other immunosuppressives and the continued use of oral GCs after the IV pulses. A small retrospective study showed no increased HBVr risk for low dose GC pulses (12.5–100 mg/day) given for up to 1 week,<sup>27</sup> whereas, on the contrary, a retrospective Taiwanese study revealed a HBVr rate of 15% (11/72) in patients treated with IV GC pulses (625–750 mg/day) for 3 days.<sup>28</sup> These data should be interpreted with caution, since most of these patients

continued therapy with oral GCs (mean daily dose = 23 mg) and other immunosuppressives (biologics or non-biologics).<sup>28</sup> Recent AGA guidelines do not offer any specific recommendations regarding patients treated with IV GC pulses due to the absence of data, but note that any GC dose given for <1 week is considered low risk (<1%), and, thus, no antiviral prophylaxis is recommended.<sup>17</sup>

There is a relevant paucity of data looking into the effect of GC in HBVr risk specifically for patients with resolved HBV infection, but certainly this risk seems to be lower than in chronic HBV infection. In a retrospective Chinese study of a large HBsAg negative/anti-HBc positive population treated with at least one dose of systemic GCs for all indications, the incidence rate of HBsAg seroreversion was 1.8% at 1 year and 5.5% at 10 years. GC peak daily dose >20 mg prednisolone equivalents and treatment duration for >4 weeks were independent risk factors for a hepatitis flare but not for HBsAg seroreversion.<sup>31</sup> The authors could not identify studies from geographical areas with less prevalent HBV infection to confirm this relatively high HBVr rate.

GC use at the doses needed for adrenal insufficiency has been reported to be safe, and does not increase the risk for HBVr.<sup>32</sup>

Based on the aforementioned data, guidance from AGA and expert advice suggests that daily doses of GCs >20 mg prednisolone equivalents for treatment durations >4 weeks should be considered as having at least moderate risk for HBVr and warrant antiviral prophylaxis in HBsAg positive patients.<sup>17,33</sup> On the contrary, the risk is low in HBsAg negative/anti-HBc positive patients and monitoring, rather than prophylaxis treatment, is advised.

### *Non-biologic agents*

*Methotrexate and other conventional synthetic disease-modifying antirheumatic drugs.* Methotrexate (MTX) is an inhibitor of folate metabolism, and has served as the anchor drug for RA for several decades, while it is also frequently used in the management of other rheumatic diseases [spondyloarthropathies, systemic lupus erythematosus (SLE), vasculitis, myositis, scleroderma, etc.]. MTX is associated with direct acute (hepatitis) and more rarely with chronic (fibrosis) hepatotoxicity. With appropriate pre-treatment screening and regular monitoring, these events are rarely seen today.

In a retrospective analysis of a Thai population with rheumatic diseases being treated with MTX for an average of 9.9 years, no cases of HBVr or hepatitis flares were identified,<sup>34</sup> whereas in a similar data analysis from a national Taiwanese health database, no increased risk of liver cirrhosis was seen (compared with non-users of MTX).<sup>35</sup>

With regards to other non-biologic agents such as leflunomide (LEF), sulfasalazine (SSZ), hydroxychloroquine (HCQ), and azathioprine (AZA), cases of HBVr are extremely rare, and these drugs should be generally considered safe.<sup>17</sup> In a recent prospective study of chronically HBV-infected RA patients treated with MTX, LEF, SSZ, or HCQ without antiviral prophylaxis, only 4/211 (1.8%) developed HBVr; all 4 patients were also receiving GCs, and, thus, the exact contribution of these agents to HBVr is unclear.<sup>36</sup> Similarly, among patients with rheumatic disease and resolved HBV infection taking part in a large prospective multi-center study, use of MTX was not associated with increased risk for HBVr (RR = 0.4).<sup>37</sup>

Overall, MTX and the other non-biologic agents should be considered as low risk treatment for HBVr in patients with both chronic and resolved HBV infection. Nevertheless, for those agents with potential direct hepatotoxic effect (MTX, LEF), appropriate monitoring precautions should always be in place.

*Mycophenolate mofetil.* Mycophenolate mofetil (MMF) is an oral inhibitor of *de novo* purine synthesis in lymphocytes, and is prescribed for a variety of approved and off-label indications in rheumatology, such as SLE, ANCA-associated vasculitides (AAV), scleroderma, etc. No clear association between MMF administration and HBVr has been identified so far. Interestingly, earlier *in vitro* data supported an inhibitive effect of MMF on replication of several viruses, HBV included,<sup>38,39</sup> whereas other clinical studies in solid organ transplant recipients exhibited neutral impact of MMF on HBV replication.<sup>40,41</sup>

While data from the rheumatology field are limited, Li and colleagues compared a standard prednisone regimen (1 mg/kg daily) *versus* MMF (500–1000 mg twice daily) and lower prednisone dose (0.5 mg/kg daily) in HBsAg-positive patients with idiopathic nephrotic syndrome and ‘undetectable’ HBV-DNA (<1.000 copies/mL).<sup>42</sup> HBVr



(defined as HBV-DNA >1.000 copies/mL) during follow up was recorded in 64% (14/22) and 37% (7/19) of patients on standard prednisone monotherapy and MMF/lower prednisone combination, respectively ( $p=0.047$ ), highlighting the role of high GC doses in the replicative potential of HBV. In a retrospective study of SLE patients with chronic or resolved HBV infection from Taiwan, MMF was not found to be among risk factors for HBVr.<sup>43</sup>

Taking into account the available data, one can conclude that MMF carries a low risk for HBVr, similar to that of other conventional synthetic disease modifying antirheumatic drugs (csDMARDs).

**Cyclophosphamide.** Cyclophosphamide (CYC)-containing regimens have been recognized among those carrying the highest risk for HBVr in patients with hematological malignancies. In this particular context, however, CYC is administered in combination with other cytotoxic drugs implicated in HBVr (i.e. doxorubicin), and the resulting state of immunosuppression cannot be compared with that of rheumatic patients treated with CYC. For the majority of circumstances in rheumatology, CYC is prescribed alongside GCs for induction of remission in patients with severe manifestations of SLE and systemic vasculitides, such as AAV, polyarteritis nodosa (PAN), and CNS vasculitis. Whereas things are more-or-less clear-cut with patients with chronic HBV infection being candidates for prophylactic antiviral treatment, data on the management of rheumatic patients with resolved HBV infection planned to be treated with CYC-containing regimens remain scarce. In a large cohort of 2054 rheumatic patients,<sup>44</sup> (anti-HBc positive: 5%,  $n=183$  with vasculitis), eight cases of HBVr were recorded, but only one occurred in a patient with granulomatosis with polyangiitis (GPA) treated with CYC.

The kinetics and outcomes of HBVr in various immune-mediated inflammatory diseases from a wide spectrum of medical specialties were described in a French study combining physician-collected and published patients.<sup>45</sup> The authors included a total of 138 HBV-infected patients, most with chronic HBV infection ( $n=99$ , 72%); among them, 11 (8%) had been treated with CYC and experienced HBVr. CYC-induced HBVr occurred earlier than that associated with rituximab (RTX) or tumor necrosis factor inhibitors (TNFi) (8 versus 13 versus 26 weeks, respectively,  $p=0.009$ ).

Lin *and* colleagues reported HBVr rates in a cohort of 195 SLE patients who were treated with GCs and various immunosuppressives (including CYC) with chronic ( $n=38$ ) or resolved ( $n=157$ ) HBV infection.<sup>43</sup> Among 34 patients with chronic HBV infection not on antiviral prophylaxis, HBVr occurred in 15 (44%), whereas in contrast, only 3 (1.9%) patients with resolved infection experienced HBsAg seroreversion. However, in multivariate analysis, CYC was not associated with increased HBVr risk, and the main driver of HBVr in this cohort was an average prednisolone daily dose of >5 mg.

From these limited data, it appears that the risk for HBVr in CYC-containing schemes is high, and occurs relatively early in HBsAg positive patients, while the respective risk in patients with resolved HBV infection is rather low. Prophylactic antiviral prophylaxis should be used for the former group. We believe that current evidence does not support universal antiviral protection for patients with resolved HBV infection prior to induction of remission with CYC-containing regimens.

#### *Biologic disease-modifying antirheumatic drugs*

**TNFi.** TNF inhibition has revolutionized the treatment of several rheumatic diseases since the late 1990s. When it comes to HBV infection, however, TNF $\alpha$  is an important cytokine to assist in the eradication of the virus from the liver. Low levels of TNF $\alpha$  have been associated with reduced cytotoxic CD8+ responses against HBV and TNF $\alpha$  neutralization may favor HBVr.<sup>46</sup>

The risk of HBVr in chronic HBV infection patients receiving TNFi has been identified and is well described in several publications. Incidence rates of HBVr in chronically HBV infected patients who did not receive antiviral prophylaxis range between 7.1% and 75% in the setting of rheumatic diseases.<sup>25,47-54</sup> Most HBVr cases were associated with a hepatitis flare and in their majority they appeared within the first year of TNFi treatment. Nucleos(t)ide analogue treatment at the time of HBVr usually results in prompt control of the HBV replication and improvement of the hepatic function, although there have been cases of severe liver injury and hepatic failure despite antiviral treatment at the time of HBVr. It has been also shown that antiviral treatment is efficacious in preventing HBVr as supported by the negligible rates of HBVr in this setting.<sup>5</sup>

On the contrary, it seems that the HBVr risk in resolved HBV infection patients treated with TNF inhibitors is significantly lower.<sup>47–49,52,55,56</sup> In a recent meta-analysis of 468 published HBsAg negative/anti-HBc positive rheumatic patients treated with TNFi, Lee and colleagues reported an HBVr incidence rate of 1.7%.<sup>57</sup> Similarly, in a prospective multicenter Italian study,<sup>58</sup> no cases of HBVr were found among 146 resolved HBV infection patients treated with TNFi for a mean period of 56 months, whereas in a prospective Japanese study, the respective rate of HBVr was 4% (8/199).<sup>37</sup> It is unclear why these differences in HBVr between European and Asian studies exist, but it could be due to different states of the underlying HBV infection (occult *versus* resolved) or due to different HBV strains.

On the basis of these data, TNF inhibition in patients with chronic HBV infection (HBsAg+) carries a significant HBVr risk and it is recommended that all patients receive appropriate oral antiviral prophylaxis with the newer antiviral agents (entecavir, tenofovir). However, in patients with resolved HBV infection, the HBVr risk is rather low, and careful monitoring of aminotransferases (AST, ALT) with prompt HBV-DNA measurement in the case of their elevation should be sufficient in managing these patients.

*RTX and other B-cell depleting therapies.* Anti-CD20 agents (rituximab, ofatumumab, ocrelizumab) are monoclonal antibodies targeting CD20, a surface cell antigen of B lymphocytes. The immune complexes formed after their binding are recognized and destroyed by phagocytes, thus leading to B-cell depletion. RTX has been used for various rheumatic diseases including RA, AAV, cryoglobulinemic vasculitis, and off-label for several connective tissue diseases including SLE, scleroderma, myositis, and Sjögren's syndrome.

The observation that anti-CD20 monoclonal antibodies can cause HBVr dates back to more than 20 years ago. So far, 183 RTX-associated HBVr cases were identified in a medical literature search between 1997 and 2009.<sup>59</sup> This eventually led to the addition of a black box warning for HBVr risk to the RTX product label.<sup>60</sup> Once again, the majority of studies of HBVr in HBsAg-positive patients derive from the hematology and oncology literature.<sup>61,62</sup> In studies looking into the benefit of using antiviral prophylaxis in chronic HBV infection patients receiving RTX-containing

regimens, the HBVr rate was as high as 53%, but was essentially nullified by the use of lamivudine or entecavir.<sup>63–65</sup>

In the setting of rheumatic diseases, the available HBVr data for HBsAg-positive patients receiving immunosuppression without antiviral prophylaxis is scarce, but the HBVr risk should be considered as high. Overall, it should be considered that there is robust data to support that RTX use in chronic HBV infection is associated with very high HBVr rates (30–60%)<sup>17,33</sup> and antiviral prophylaxis is warranted. There is also evidence that, in the setting of rheumatic disease, the duration of immunosuppression should be taken into account when choosing the agent for antiviral prophylaxis, as there are case reports of virological escape with the use of drugs with low genetic barrier (e.g. lamivudine).<sup>66</sup> Third-generation antiviral agents with high genetic barrier (entecavir, tenofovir) should be used.

In comparison with other biologic agents, RTX has been associated with high HBVr rates also in patients with resolved HBV infection.<sup>63,67–71</sup> In their technical review, which supported the AGA guidelines, Perrillo and colleagues calculated a pooled baseline risk estimate of HBVr at 16.9% in this group of patients, which was based on hematology/oncology studies.<sup>17</sup> However, in prospective or retrospective studies with patients suffering with rheumatic diseases, the rate of HBVr under RTX therapy was much lower (6/266 patients or 2%).<sup>58,72–76</sup> Interestingly, the majority of reported patients with HBVr (4/6) came from a high-incidence HBV country (Taiwan).<sup>76</sup>

Although it remains an area of controversy, the authors believe that in patients with resolved HBV infection and undetectable baseline serum HBV-DNA, serial monitoring of aminotransferases, HBsAg, or serum HBV-DNA is a preferable alternative to long-term commitment to antiviral prophylaxis.

*Abatacept.* Abatacept (ABA) is a fusion protein composed of the Fc portion of the immunoglobulin IgG1 fused to the extracellular domain of CTLA-4; ABA acts by binding to the CD80/CD86 molecule, thus inhibiting the activation of T-cells by an antigen-presenting cell. The agent has been approved for RA and psoriatic arthritis, and there are ongoing trials for other rheumatic diseases (giant cell arteritis, polymyalgia rheumatica, myositis).

The HBVr risk for chronically infected HBV patients on ABA treatment has been explored only in small retrospective studies. In a study by Kim and colleagues,<sup>77</sup> four of eight HBsAg positive patients who did not receive prophylaxis developed HBVr after a mean period of 10.5 months. An Italian retrospective multicenter study collected data on 72 RA patients treated with ABA who had either chronic or resolved HBV infection.<sup>78</sup> All of the chronically infected patients received antiviral prophylaxis. No HBVr cases were identified in either group.

In the setting of resolved HBV infection, a few case reports of HBVr cases have been published.<sup>79,80</sup> Three cases of HBVr in a recently studied Japanese cohort were reported on resolved HBV infection patients treated with ABA (total  $n=29$ , 10.3%).<sup>81</sup> In another study, among 24 patients with resolved HBV infection on ABA treatment, three cases of HBVr were reported (12.5%),<sup>37</sup> but these were mild elevations of HBV-DNA without overt hepatitis and without the need to use antivirals. Reassuringly, no HBVr cases were observed among the nine resolved HBV infection patients treated with ABA in the prospective trial by Barone and colleagues.<sup>58</sup> Considering the high number of patients who have been treated with ABA since its approval, cases of HBVr in resolved HBV infection appear to be rare, especially in geographical areas with lower prevalence, such as Europe and the United States.

Taken together, these data would suggest that, in ABA-treated HBV patients with rheumatic disease, there is a need for antiviral prophylaxis in chronic HBV infection, but close monitoring in resolved HBV infection seems to be a safe alternative.

*Interleukin-6 inhibitors.* Inhibition of interleukin (IL)-6 provides a new mechanism of action for the treatment of rheumatic disease, and two agents have been approved for use in RA and systemic Juvenile Idiopathic Arthritis (sJIA) (tocilizumab-TCZ, sarilumab) and giant cell arteritis (GCA; tocilizumab) with several others being investigated. Most of the available HBVr relevant data is limited,<sup>82</sup> and focus on the use of TCZ, which is a humanized IgG1k monoclonal antibody against the soluble and membrane forms of the IL-6 receptor that inhibits the characteristic signalling pathway of IL-6, a pleiotropic cytokine.

There are only few case reports and small studies showing that HBVr can occur with TCZ therapy in patients with chronic HBV infection without antiviral prophylaxis.<sup>25,83-86</sup> In all published data, the use of antivirals prevented HBVr in this setting. In rheumatic patients with resolved infection, HBVr is rare,<sup>58,87</sup> and has been reported only in the form of transient HBV viremia.<sup>37,88</sup>

Similarly to ABA and TNFi, antiviral prophylaxis is warranted for HBsAg-positive patients, whereas aminotransferase monitoring (with HBV-DNA measurement in the case of AST or ALT elevation) is sufficient for HBsAg negative/anti-HBc positive patients.

*Interleukin-17 inhibitors.* IL-17 inhibitors have recently been approved for the treatment of psoriasis, psoriatic arthritis, and axial spondyloarthritis. As patients with HBV infection were excluded from the initial controlled clinical trials for these novel agents, data regarding their potential for HBVr are scarce.

There are few case reports and case series showing no safety signals for HBVr with secukinumab (a human IgG1k anti-IL17A monoclonal antibody)<sup>89-92</sup> or ixekizumab (a humanized monoclonal anti-IL17A antibody).<sup>93,94</sup> The risk for HBVr with secukinumab was studied in a multicenter prospective cohort study of 49 Taiwanese patients with either chronic or resolved HBV infection. Among those with chronic HBV infection who did not receive antiviral prophylaxis, 27% (6/22) developed virological HBVr without a hepatitis flare, while the respective rate of virological HBVr in those with resolved HBV infection was 4% (1/24).<sup>95</sup>

In the absence of solid data showing that IL-17A inhibition is associated with a substantial increase in HBVr risk, an approach similar to that used with other biologic agents is suggested.

*Interleukin-12/23 inhibitors.* Ustekinumab is a fully human IgG1 monoclonal antibody that inhibits the p40 subunit of both IL-12 and IL-23, rendering them unable to bind to their receptors. It has been approved for psoriasis, psoriatic arthritis, and inflammatory bowel disease (Crohn's disease, ulcerative colitis).

Several case reports and two retrospective studies have looked into HBVr risk in patients treated



with ustekinumab. In a single-center Taiwanese retrospective cohort,<sup>96</sup> 11 chronic HBV and 3 resolved HBV infection patients with psoriasis received ustekinumab. HBVr was observed in two of the seven chronic HBV infection patients who did not receive antiviral prophylaxis; in both cases the HBVr was mild and only virological, without biochemical hepatitis. In a similar Asian study by Ting and colleagues of 54 psoriasis patients treated with ustekinumab,<sup>97</sup> the rate of HBVr was 17% for chronically infected patients and 1.5% for those with resolved infection.

Based on these data, it appears that the risk for HBVr with ustekinumab is similar to that of TNF inhibitors and the advice on prophylaxis and monitoring is the same.

*Interleukin-23 inhibitors.* Novel anti-IL23 agents, such as guselkumab, risankizumab, and tildrakizumab have been approved for the treatment of psoriasis and are currently under investigation for psoriatic arthritis. No literature data on their use in the setting of HBV infection are available and, thus, no relevant recommendations exist.

*csDMARD and biologic disease-modifying antirheumatic drug combinations.* The question of whether csDMARD and biologic disease-modifying antirheumatic drug (bDMARD) combination treatment in rheumatic patients carries higher risk for HBVr compared with bDMARD monotherapy has not been specifically addressed in the published literature. In a retrospective study by Chen and colleagues looking into HBsAg positive patients treated without antiviral prophylaxis, there was no significant difference in HBVr risk when comparing csDMARD monotherapy to csDMARD and bDMARD combination treatment.<sup>25</sup> The majority of bDMARD agents used were TNF inhibitors (26/36, 72%).

### Oral targeted therapies

*JAK inhibitors.* JAK inhibitors belong to a novel class of targeted synthetic agents that were recently introduced into daily clinical practice for treatment of RA, psoriatic arthritis, and ulcerative colitis. JAK inhibitors have a unique mode of action, inhibiting production of various proinflammatory cytokines from T-lymphocytes and dendritic cells.<sup>98</sup> To date, three agents of this family have been approved for use in RA (tofacitinib, baricitinib, and upadacitinib), with another currently under investigation (filgotinib). As for

psoriatic arthritis, tofacitinib is the only JAK inhibitor currently approved for use after methotrexate or other csDMARD failure. Data for their HBVr potential are rare, given that these agents' pivotal clinical trials excluded patients with evidence of chronic or resolved HBV infection.

Two recent studies attempted to investigate the risk for HBVr with JAK inhibitor use. In a retrospective cohort of 116 tofacitinib-treated RA patients from Taiwan, 6 (5%) had chronic and 75 (65%) resolved HBV infection.<sup>99</sup> Two of the HBsAg positive patients were prophylactically treated with antivirals upon tofacitinib initiation and did not experience HBVr. Amongst the four HBsAg-positive patients not treated with antivirals, two had HBVr after 6 and 12 months of tofacitinib therapy, with one having only virological and the other virological and biochemical reactivation. Both were being cotreated with low-dose corticosteroids (5 and 10 mg of prednisolone daily, respectively). Patients who were HBsAg negative/anti-HBc positive did not experience HBVr.

Harigai and colleagues studied the effect of baricitinib in RA patients with past HBV infection in a *post hoc* analysis of baricitinib clinical trials.<sup>100</sup> They included 269 patients, 255 of whom were anti-HBc positive/antibody against hepatitis B surface antigen (anti-HBs) positive and 14 anti-HBc positive/anti-HBs negative. Among the former group, seven (3%) patients had detectable baseline HBV DNA (median viral load 256 IU/mL), while among the latter group, one (7%) had marginally detectable HBV viral load (36 IU/mL). From the total of eight patients with detectable baseline HBV DNA, half discontinued baricitinib, and three out of four were treated with antivirals. The remaining four patients continued baricitinib without antivirals. None of those patients experienced clinical or biochemical hepatitis.

From these preliminary data, a profile of JAK inhibitors similar to that of anti-TNFs is emerging with regards to HBVr. Thus, the aforementioned strategy should also be used in this patient population.

*Role of anti-HBs titers in HBVr rates.* In a significant proportion (73–84%) of patients with resolved HBV infection (HBsAg negative/anti-HBc positive), circulating anti-HBs antibodies can be detected in the serum.<sup>101,102</sup> A number of

retrospective studies in hematology and oncology patients have shown that the presence of anti-HBs antibodies reduces the risk for HBVr during chemotherapy (including RTX-containing regimens).<sup>68,103,104</sup> The results of these studies were corroborated by a relevant meta-analysis of 20 studies looking into the role of anti-HBs levels in HBVr risk of patients with hematological malignancy, which found that the presence of detectable anti-HBs reduced HBVr [pooled odds ratio (OR)=0.21], even in patients with lymphoma receiving RTX chemotherapy.<sup>105</sup>

Similar data have been reported in rheumatic patients with resolved HBV infection treated with different immunosuppressives and prospective studies support the predictive value of anti-HBs. Over a period of 4 years, 380 rheumatic patients on biologic therapy from a single Chinese center were identified as having resolved HBV infection, and were classified according to anti-HBs levels to 'negative' (<10 IU/mL), 'low' (10–100 IU/mL), and 'high' (>100 IU/mL) groups. There were no HBVr cases in the 'high' group, whereas cases of transient HBV viremia were seen in the both the 'low' (2.5/100 person-years) and 'negative' group (4.7/100 person-years).<sup>106</sup> Among 1042 Japanese patients with rheumatic diseases and resolved HBV infection who received both conventional and biological treatments, HBVr incidence was overall low (1.93/100 person-years), but increased to 4.32/100 person-years in anti-HBc positive/anti-HBs negative patients. The risk ratios for HBVr were 2.8 and 3.1 for anti-HBs titres below the median (71.4 IU/mL) and the cut-off for detection (<10 IU/mL), respectively.<sup>37</sup>

Finally, a meta-analysis of HBVr in studies of patients with resolved HBV infection from all medical fields showed that the total pooled rate of HBV reactivation was lower in patients with positive anti-HBs (5.2% versus 17.0%; RR: 0.29).<sup>107</sup>

Overall, these (mainly retrospective) data suggest that the presence of anti-HBs antibodies in patients with resolved HBV infection, especially when present in high titers (>100 IU/mL), significantly decreases the risk for HBVr after immunosuppressive therapies. Physicians should be vigilant in the care of patients with isolated HBc positivity when high risk treatments are used. Nevertheless, in the absence of prospective data from randomized trials showing that patients with high anti-HBs titers do not need appropriate

monitoring, or, in certain cases, antiviral prophylaxis for HBVr, we cannot currently recommend serial anti-HBs measurements replacing monitoring with serum HBV DNA.

## Recommendations for HBVr screening and management

### Screening

The valued benefit of screening for HBV infection prior commencing immunosuppressive treatment has been established in several medical society guidelines. Increasing physician awareness for the risk of HBVr has improved clinical outcomes in the past,<sup>108</sup> whereas, with appropriate screening, up to 80% of HBVr could be preventable.<sup>45</sup>

The cost-effectiveness of HBV screening has been studied in several studies from the hematology-oncology field, where a universal screening strategy was found to be cost-effective in studies with patients having solid tumors and hematologic malignancies before chemotherapy.<sup>109,110</sup> A tool for HBV risk calculation was used in a US cohort with hematologic or solid malignancies,<sup>111</sup> and it seemed to improve cost-effectiveness by screening high-risk patients only. We were unable to identify similar studies assessing HBV screening cost-effectiveness in rheumatic patients prior to immunosuppression.

Given the increased risk of HBVr associated with the immunosuppressive potency of the treatment used, we recommend universal screening in patients scheduled to start biologic and targeted synthetic DMARD therapy, as well as moderate to high doses of glucocorticoids for prolonged periods (>4 weeks duration). For conventional DMARDs, the risk of drug-related hepatotoxicity should be also of concern (especially for methotrexate and leflunomide) and, thus, screening should be performed in all patients.<sup>5</sup> A suggested algorithm for screening and interpretation of results is presented in Figure 1.

The authors support the inclusion of anti-HBs in the serological screening markers (HBsAg, anti-HBc) as suggested by recent American Association for the Study of Liver Diseases (AASLD) and European Association for the Study of the Liver (EASL) guidelines,<sup>10,11</sup> in order to identify cases that may benefit from initial or booster vaccination, but also cases of occult HBV infection in

anti-HBs positive patients that have not been vaccinated in the past and have lost their anti-HBc.

### Vaccination

Vaccination still remains the most effective intervention for prevention of acute and chronic HBV infection. The available vaccine containing recombinant HBsAg shows excellent efficacy in younger individuals (90% in those <40 years old); however, these responses could be attenuated in older persons.<sup>112,113</sup>

A screening algorithm that includes HBsAg, anti-HBc, and anti-HBs (Figure 1) is capable of identifying rheumatic patients not exposed to HBV [HBsAg/anti-HBc/anti-HBs(-)], and a complete series of vaccination should be offered to those at risk of HBV exposure according to their occupation, medical status, or sexual behavior. The complete vaccination schedule with the recombinant HBsAg vaccine consists of three doses (20 µg per dose) administered at 0, 1, and 6 months. Patients on dialysis should be vaccinated with high-dose (40 µg) vaccines with the same time intervals.<sup>114</sup>

A new two-dose HBV vaccine was recently approved by the United States Food and Drug Administration (FDA) and is already recommended by Advisory Committee on Immunization Practices (ACIP) for individuals >18 years. This vaccine includes a novel oligonucleotide adjuvant that activates the toll-like receptor 9 (TLR-9) pathway and has been shown to have higher vaccine efficacy compared with recombinant vaccine, even in difficult populations, such as patients with diabetes.<sup>115,116</sup> To date, data on the efficacy and safety of this novel vaccine in rheumatic patients are lacking.

### Management

The management of HBVr (Figure 2) should be based on individual HBVr risk according to the patient's HBV status (chronic or resolved infection), but also on the HBVr potential of the immunosuppressive treatment used (Table 4).

Patients who are at moderate or high risk of HBVr should be considered candidates for prophylactic antiviral treatment. Unlike in the oncology field, in the setting of rheumatic disease, immunosuppressive treatment is usually used for prolonged


periods of time, even lifelong. Thus, there is need for enduring and potent control of viral replication with negligible drug resistance. Given lamivudine's low genetic barrier and potential development of drug resistance (ranging from 24% in the first year up to 70% at 5 years),<sup>11</sup> currently the newer nucleos(t)ide analogues (NAs) with high barrier to genetic resistance, such as entecavir (ETV), tenofovir disoproxil fumarate (TDF), and tenofovir alafenamide (TAF) are the antiviral agents of choice (Figure 2).<sup>10,11</sup>

The ideal timing for commencing antiviral prophylaxis, especially in the setting of high baseline viral loads, is 1–2 weeks before starting immunosuppressive treatment.<sup>15</sup> If delaying the start of immunosuppression is not feasible, antivirals should be started as soon as possible. From the available literature data on the timing for cessation of prophylaxis, the suggestion is to continue with prophylactic treatment for at least 6 months after the end of antirheumatic treatment (12 months if that treatment is RTX). Reaching the therapeutic endpoint as per hepatology guidelines should guide the decision for stopping NAs when they are used for treating chronic hepatitis B rather than for prophylaxis of reactivation.<sup>10,11</sup>

For patients that are identified as never been exposed to HBV (HBsAg/Anti-HBc/Anti-HBs-), HBV vaccination should be considered and offered to the patient. This has greater value in patients in high risk groups, including patients with high-risk sexual activity, sexual partners and household contacts of HBV-infected individuals, injectable drug users, hemodialysis patients, health care workers, and patients from endemic areas. The vaccination should ideally take place before commencing immunosuppression, especially if this is B-cell depleting treatment.<sup>113</sup>

Patients with chronic HBV infection (HBsAg+) form the group with the highest HBVr risk. Decisions about managing these patients should be always taken after consulting a specialist with experience in treating HBV infection (gastroenterologist, hepatologist, infectious disease specialist, etc.).

In chronic active hepatitis B (elevated ALT and HBV DNA levels, or at least moderate liver necroinflammation) or cirrhosis, patients should be treated with antiviral therapy according to the most recent liver societies' guidelines,<sup>10,11</sup> regardless of their administered immunosuppressive therapy.



**Figure 2.** Suggested algorithm for the management of HBVr in rheumatic diseases.

\*In certain cases, close monitoring without antiviral therapy may be chosen after appropriate consultation. See text for details.

ALT, alanine aminotransferase; anti-HBc, antibody against Hepatitis B core antigen; anti-HBs, antibody against HBsAg; ETV, entecavir; HBsAg, hepatitis B surface antigen; HBV-DNA, HBV-deoxyribonucleic acid; HBVr, hepatitis B virus reactivation; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate.

**Table 4.** Risk of HBVr stratified per HBV status and agent used.

Medication	Chronic HBV infection [HBsAg (+)]	Resolved HBV infection [HBsAg (-) / anti-HBc (+)]
<b>GCs</b>		
GC >20 mg/day and >4 weeks	Moderate	Low
<b>Nonbiologics</b>		
MTX, AZA, MMF, LEF, HCQ	Low	Low
CYC	Moderate	Low
<b>Biologics</b>		
TNFi	Moderate	Low
RTX	Very high	Moderate
ABA	Moderate	Low
IL-6 inhibitors	Moderate	Low
IL-17 inhibitors	Moderate	Low
IL-12/23 inhibitors	Moderate	Low
<b>Oral targeted therapies</b>		
JAK inhibitors	Moderate	Low

ABA, Abatacept ; anti-HBc, antibody against Hepatitis B core antigen; AZA, azathioprine; CYC, Cyclophosphamide; GC, glucocorticoids; HBsAg, Hepatitis B surface antigen; HCQ, hydroxychloroquine; IL, interleukin; JAK, Janus kinase; LEF, leflunomide; MMF, mycophenolate mofetil; MTX, methotrexate; RTX, Rituximab ; TNFi, tumor necrosis factor inhibitors.

For inactive HBV carriers (normal ALT, low or undetectable HBV DNA, minimal liver necroinflammation, and no fibrosis), the exact risk with the different immunosuppressive regimens has been difficult to discern due to the lack of prospectively collected, high-quality data. AASLD and EASL recommend antiviral therapy for all patients with at least detectable HBV DNA levels.<sup>10,11</sup>

Taking into account the chronic nature of immunosuppression in most rheumatic patients, the unknown real risk conferred with these agents and the need for repeated laboratory monitoring (including HBV DNA measurement) in these patients, we believe that all such patients should receive long-term prophylactic antiviral therapy with the newer antivirals (as for patients with chronic hepatitis B). In individual patients (i.e. those scheduled for short term GCs: <1 month or treated with low risk nonbiologics such as AZA, SSZ, HCQ), close monitoring with frequent monitoring of ALT and HBV DNA levels may be chosen, after appropriate consultation with an HBV specialist.

The issue of antiviral prophylaxis in rheumatic patients with resolved HBV infection (HBsAg+, Anti-HBc-) treated with immunosuppressives remains currently the most controversial. The discrepancies in the literature relate to the underlying disease indication (hemato-oncology *versus* rheumatic) and to the geographical area studied (Asia *versus* Europe). For example, a recent meta-analysis of HBVr in resolved HBV infection patients found that the pooled HBVr rates were much lower in non-hematological (3.6%) *versus* hematological (10.9%) diseases.<sup>107</sup> In the studies described above, one can easily detect differences in reported HBVr incidence between the Western Asia/Pacific and European regions, with the former showing significantly higher rates. This could be explained by several factors, including divergence in HBV prevalence, variance in host immune responses, different viral genotypes and inconsistent monitoring intervals.

There is general agreement that rheumatic patients treated with non-biologics, as well as biologics other than B-cell-depleting agents (like RTX), do not require prophylactic antiviral therapy.<sup>10,11</sup> These patients should be monitored with ALT, HBsAg or HBV DNA, and treated with antivirals when HBsAg or HBV DNA are detected (Table 2, Figure 2).<sup>10,11</sup> There is also

consensus that, for patients with hematologic or oncologic diseases treated with B-cell-depleting agents like RTX, prophylactic antiviral therapy is also necessary.<sup>10,11</sup>

For rheumatic patients treated with RTX, such prophylactic antiviral therapy is either not recommended or not specifically mentioned.<sup>10,11</sup> Based on the literature data presented in this review and the worldwide accumulated experience with the use of RTX in large populations of rheumatic patients with resolved infection (ranging from 5% to 50% in the general population), we believe that the risk of RTX-induced HBVr is low and prophylactic antiviral therapy is not required.<sup>6</sup> Nevertheless, baseline screening with HBV DNA and close monitoring with ALT, HBsAg, or HBV DNA (every 3–6 months) is advised for all patients treated with RTX (Figure 2). Patients with detectable HBV DNA at baseline, or during monitoring, as well as those with seroreversion (HBsAg+) during monitoring, should be treated with oral antivirals as described above.

## Summary

In patients with rheumatic disease under immunosuppressive treatment and HBV infection, HBVr is a potentially severe and life-threatening complication that rheumatologists need to be aware of. HBVr rates depend on baseline HBV status of the patient and on the therapeutic agent used, with the highest prevalence connected to biologic and targeted synthetic drugs, especially those causing B-cell depletion. Screening for HBV prior to commencing treatment, and subsequent risk stratification, is of the utmost importance, as HBVr is a largely preventable complication *via* either the use of antiviral prophylaxis or careful and regular monitoring. Questions remaining unanswered in this field include the need for prophylaxis in certain risk groups, the optimal frequency of liver function and viral load monitoring and the potential use of alternative biomarkers for the management of HBVr.

## Funding


The authors received no financial support for the research, authorship, and/or publication of this article.

## Conflict of interest statement

The authors declare that there is no conflict of interest.



**ORCID iD**

Dimitrios Vassilopoulos  <https://orcid.org/0000-0003-2288-3863>

**References**

- Schweitzer A, Horn J, Mikolajczyk RT, *et al.* Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. *Lancet* 2015; 386: 1546–1555.
- World Health Organisation. *Global hepatitis report*, 2017. Geneva: World Health Organization, 2017.
- Dougados M, Soubrier M, Antunez A, *et al.* Prevalence of comorbidities in rheumatoid arthritis and evaluation of their monitoring: results of an international, cross-sectional study (COMORA). *Ann Rheum Dis* 2014; 73: 62–68.
- Moltó A, Etcheto A, van der Heijde D, *et al.* Prevalence of comorbidities and evaluation of their screening in spondyloarthritis: results of the international cross-sectional ASAS-COMOSPA study. *Ann Rheum Dis* 2016; 75: 1016–1023.
- Koutsianas C, Thomas K and Vassilopoulos D. Hepatitis B reactivation in rheumatic diseases: screening and prevention. *Rheum Dis Clin North Am* 2017; 43: 133–149.
- Koutsianas C, Thomas K and Vassilopoulos D. Prevention of HBV reactivation in patients treated with biologic agents. *Expert Rev Clin Pharmacol* 2016; 2433: 1–11.
- Vassilopoulos D and Calabrese LH. Management of rheumatic disease with comorbid HBV or HCV infection. *Nat Rev Rheumatol* 2012; 8: 348–357.
- Seto WK, Lo YR, Pawlotsky JM, *et al.* Chronic hepatitis B virus infection. *Lancet* 2018; 392: 2313–2324.
- Hoofnagle JH. Reactivation of hepatitis B. *Hepatology* 2009; 49: S156–S165.
- Terrault NA, Lok ASF, McMahon BJ, *et al.* Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology* 2018; 67: 1560–1599.
- Lampertico P, Agarwal K, Berg T, *et al.* EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol* 2017; 67: 370–398.
- Loomba R and Liang TJ. Hepatitis B reactivation associated with immune suppressive and biological modifier therapies: current concepts, management strategies, and future directions. *Gastroenterology* 2017; 152: 1297–1309.
- Yeo W, Chan PK, Zhong S, *et al.* Frequency of hepatitis B virus reactivation in cancer patients undergoing cytotoxic chemotherapy: a prospective study of 626 patients with identification of risk factors. *J Med Virol* 2000; 62: 299–307.
- Tohme RA, Bulkow L, Homan CE, *et al.* Rates and risk factors for hepatitis B reactivation in a cohort of persons in the inactive phase of chronic hepatitis B-Alaska, 2001–2010. *J Clin Virol* 2013; 58: 396–400.
- Hwang JP and Lok AS-F. Management of patients with hepatitis B who require immunosuppressive therapy. *Nat Rev Gastroenterol Hepatol* 2014; 11: 209–219.
- Moghoofei M, Mostafaei S, Ashraf-Ganjouei A, *et al.* HBV reactivation in rheumatic diseases patients under therapy: a meta-analysis. *Microb Pathog* 2018; 114: 436–443.
- Perrillo RP, Gish R and Falck-Ytter YT. American Gastroenterological Association Institute technical review on prevention and treatment of hepatitis b virus reactivation during immunosuppressive drug therapy. *Gastroenterology* 2015; 148: 221–244.e3.
- Galbraith RM, Eddleston AL, Williams R, *et al.* Fulminant hepatic failure in leukaemia and choriocarcinoma related to withdrawal of cytotoxic drug therapy. *Lancet* 1975; 2: 528–530.
- Tur-Kaspa R, Shaul Y, Moore DD, *et al.* The glucocorticoid receptor recognizes a specific nucleotide sequence in hepatitis B virus DNA causing increased activity of the HBV enhancer. *Virology* 1988; 167: 630–633.
- Perrillo RP, Schiff ER, Davis GL, *et al.* A randomized, controlled trial of interferon alfa-2b alone and after prednisone withdrawal for the treatment of chronic hepatitis B. The Hepatitis Interventional Therapy Group. *N Engl J Med* 1990; 323: 295–301.
- Liaw Y-F, Tsai S-L, Chien R-N, *et al.* Prednisolone priming enhances Th1 response and efficacy of subsequent lamivudine therapy in patients with chronic hepatitis B. *Hepatology* 2000; 32: 604–609.
- Hoofnagle JH, Davis GL, Pappas SC, *et al.* A short course of prednisolone in chronic type B hepatitis. Report of a randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 1986; 104: 12–17.

23. Cheng A-L, Hsiung CA, Su I-J, *et al.* Steroid-free chemotherapy decreases risk of hepatitis B virus (HBV) reactivation in HBV-carriers with lymphoma. *Hepatology* 2003; 37: 1320–1328.
24. Kim T-W, Kim M-N, Kwon J-W, *et al.* Risk of hepatitis B virus reactivation in patients with asthma or chronic obstructive pulmonary disease treated with corticosteroids. *Respirology* 2010; 15: 1092–1097.
25. Chen MH, Chen MH, Liu CY, *et al.* Hepatitis B virus reactivation in rheumatoid arthritis patients undergoing biologics treatment. *J Infect Dis* 2017; 215: 566–573.
26. Wong GL-H, Yuen BW-Y, Chan HL-Y, *et al.* Impact of dose and duration of corticosteroid on the risk of hepatitis flare in patients with chronic hepatitis B. *Liver Int* 2019; 39: 271–279.
27. Braun-Moscovici Y, Braun M, Saadi T, *et al.* Safety of corticosteroid treatment in rheumatologic patients with markers of hepatitis B viral infection. *J Clin Rheumatol* 2016; 22: 364–368.
28. Lin Y-C, Lee S-W, Yeh H-Z, *et al.* The prevalence and risk factors of hepatitis B flares in chronic hepatitis B patients receiving glucocorticoid pulse therapy. *Int J Clin Pharm* 2018; 40: 169–174.
29. Yang C-H, Wu T-S and Chiu C-T. Chronic hepatitis B reactivation: a word of caution regarding the use of systemic glucocorticosteroid therapy. *Br J Dermatol* 2007; 157: 587–90.
30. Xuan D, Yu Y, Shao L, *et al.* Hepatitis reactivation in patients with rheumatic diseases after immunosuppressive therapy - A report of long-term follow-up of serial cases and literature review. *Clin Rheumatol* 2014; 33: 577–586.
31. Wong GL-H, Wong VW-S, Yuen BW-Y, *et al.* Risk of hepatitis B surface antigen seroreversion after corticosteroid treatment in patients with previous hepatitis B virus exposure. *J Hepatol.* Epub ahead of print 6 September 2019. DOI: 10.1016/j.jhep.2019.08.023.
32. Hatano M, Mimura T, Shimada A, *et al.* Hepatitis B virus reactivation with corticosteroid therapy in patients with adrenal insufficiency. *Endocrinol Diabetes Metab* 2019; e00071.
33. Di Bisceglie AM, Lok AS, Martin P, *et al.* Recent US Food and Drug Administration warnings on hepatitis B reactivation with immune-suppressing and anticancer drugs: just the tip of the iceberg? *Hepatology* 2015; 61: 703–711.
34. Laohapand C, Arromdee E and Tanwandee T. Long-term use of methotrexate does not result in hepatitis B reactivation in rheumatologic patients. *Hepatol Int* 2015; 9: 202–208.
35. Tang K-T, Hung W-T, Chen Y-H, *et al.* Methotrexate is not associated with increased liver cirrhosis in a population-based cohort of rheumatoid arthritis patients with chronic hepatitis B. *Sci Rep* 2016; 6: 22387.
36. Tan J, Zhou J, Zhao P, *et al.* Prospective study of HBV reactivation risk in rheumatoid arthritis patients who received conventional disease-modifying antirheumatic drugs. *Clin Rheumatol* 2012; 31: 1169–1175.
37. Fukuda W, Hanyu T, Katayama M, *et al.* Incidence of hepatitis B virus reactivation in patients with resolved infection on immunosuppressive therapy for rheumatic disease: a multicentre, prospective, observational study in Japan. *Ann Rheum Dis* 2017; 76: 1051–1056.
38. Gong ZJ, De Meyer S, Clarysse C, *et al.* Mycophenolic acid, an immunosuppressive agent, inhibits HBV replication in vitro. *J Viral Hepat* 1999; 6: 229–236.
39. Ying C, De Clercq E and Neyts J. Ribavirin and mycophenolic acid potentiate the activity of guanine- and diaminopurine-based nucleoside analogues against hepatitis B virus. *Antiviral Res* 2000; 48: 117–124.
40. Maes BD, van Pelt JF, Peeters JC, *et al.* The effect of mycophenolate mofetil on hepatitis B viral load in stable renal transplant recipients with chronic hepatitis B. *Transplantation* 2001; 72: 1165–1166.
41. Ben-Ari Z, Zemel R and Tur-Kaspa R. The addition of mycophenolate mofetil for suppressing hepatitis B virus replication in liver recipients who developed lamivudine resistance-no beneficial effect. *Transplantation* 2001; 71: 154–156.
42. Li X, Tian J, Wu J, *et al.* A comparison of a standard-dose prednisone regimen and mycophenolate mofetil combined with a lower prednisone dose in chinese adults with idiopathic nephrotic syndrome who were carriers of hepatitis B surface antigen: a prospective cohort study. *Clin Ther* 2009; 31: 741–750.
43. Lin WT, Chen YM, Chen DY, *et al.* Increased risk of hepatitis B virus reactivation in systemic lupus erythematosus patients receiving immunosuppressants: a retrospective cohort study. *Lupus* 2018; 27: 66–75.
44. Schwaneck EC, Krone M, Kreissl-Kemmer S, *et al.* Management of anti-HBc-positive patients with rheumatic diseases treated with disease-modifying antirheumatic drugs—a single-center

- analysis of 2054 patients. *Clin Rheumatol* 2018; 37: 2963–2970.
45. Droz N, Gilardin L, Cacoub P, *et al.* Kinetic profiles and management of hepatitis B virus reactivation in patients with immune-mediated inflammatory diseases. *Arthritis Care Res* 2013; 65: 1504–1514.
  46. Carroll MB and Forgiione MA. Use of tumor necrosis factor alpha inhibitors in hepatitis B surface antigen-positive patients: a literature review and potential mechanisms of action. *Clin Rheumatol* 2010; 29: 1021–1029.
  47. Vassilopoulos D, Apostolopoulou A, Hadziyannis E, *et al.* Long-term safety of anti-TNF treatment in patients with rheumatic diseases and chronic or resolved hepatitis B virus infection. *Ann Rheum Dis* 2010; 69: 1352–1355.
  48. Pérez-Alvarez R, Díaz-Lagares C, García-Hernández F, *et al.* Hepatitis B Virus (HBV) Reactivation in Patients Receiving Tumor Necrosis Factor (TNF)-Targeted Therapy. *Medicine* 2011; 90: 359–371.
  49. Lan J-L, Chen Y-M, Hsieh T-Y, *et al.* Kinetics of viral loads and risk of hepatitis B virus reactivation in hepatitis B core antibody-positive rheumatoid arthritis patients undergoing anti-tumour necrosis factor alpha therapy. *Ann Rheum Dis* 2011; 70: 1719–1725.
  50. Zingarelli S, Frassi M, Bazzani C, *et al.* Use of tumor necrosis factor-alpha-blocking agents in hepatitis B virus-positive patients: reports of 3 cases and review of the literature. *J Rheumatol* 2009; 36: 1188–1194.
  51. Chung S, Kim JK, Park M-C, *et al.* Reactivation of hepatitis B viral infection in inactive HBsAg carriers following anti-tumor necrosis factor-alpha therapy. *J Rheumatol* 2009; 36: 2416–2420.
  52. Tamori A, Koike T, Goto H, *et al.* Prospective study of reactivation of hepatitis B virus in patients with rheumatoid arthritis who received immunosuppressive therapy: evaluation of both HBsAg-positive and HBsAg-negative cohorts. *J Gastroenterol* 2011; 46: 556–564.
  53. Ryu HH, Lee EYB, Shin K, *et al.* Hepatitis B virus reactivation in rheumatoid arthritis and ankylosing spondylitis patients treated with anti-TNF $\alpha$  agents: a retrospective analysis of 49 cases. *Clin Rheumatol* 2012; 31: 931–936.
  54. Ye H, Zhang X, Mu R, *et al.* Anti-TNF therapy in patients with HBV infection-analysis of 87 patients with inflammatory arthritis. *Clin Rheumatol* 2014; 33: 119–123.
  55. Mori S. Past hepatitis B virus infection in rheumatoid arthritis patients receiving biological and/or nonbiological disease-modifying antirheumatic drugs. *Mod Rheumatol* 2011; 21: 621–627.
  56. Caporali R, Bobbio-Pallavicini F, Atzeni F, *et al.* Safety of tumor necrosis factor alpha blockers in hepatitis B virus occult carriers (hepatitis B surface antigen negative/anti-hepatitis B core antigen positive) with rheumatic diseases. *Arthritis Care Res* 2010; 62: 749–754.
  57. Lee YH, Bae SC and Song GG. Hepatitis B virus (HBV) reactivation in rheumatic patients with hepatitis core antigen (HBV occult carriers) undergoing anti-tumour necrosis factor therapy. *Clin Exp Rheumatol* 2013; 31: 118–121.
  58. Barone M, Notarnicola A, Lopalco G, *et al.* Safety of long-term biologic therapy in rheumatologic patients with a previously resolved hepatitis B viral infection. *Hepatology* 2015; 62: 40–46.
  59. Evens a. M, Jovanovic BD, Su YC, *et al.* Rituximab-associated hepatitis B virus (HBV) reactivation in lymphoproliferative diseases: Metaanalysis and examination of FDA safety reports. *Ann Oncol* 2011; 22: 1170–1180.
  60. Mitka M. FDA: Increased HBV reactivation risk with ofatumumab or rituximab. *JAMA* 2013; 310: 1664.
  61. Pei S-N, Chen C-H, Lee C-M, *et al.* Reactivation of hepatitis B virus following rituximab-based regimens: a serious complication in both HBsAg-positive and HBsAg-negative patients. *Ann Hematol* 2010; 89: 255–262.
  62. Paul S, Saxena A, Terrin N, *et al.* Hepatitis B virus reactivation and prophylaxis during solid tumor chemotherapy: a systematic review and meta-analysis. *Ann Intern Med* 2016; 164: 30–40.
  63. Hsu C, Hsiung CA, Su IJ, *et al.* A revisit of prophylactic lamivudine for chemotherapy-associated hepatitis B reactivation in non-Hodgkin's lymphoma: a randomized trial. *Hepatology* 2008; 47: 844–853.
  64. Lau GKK, Yiu HHY, Fong DYT, *et al.* Early is superior to deferred preemptive lamivudine therapy for hepatitis B patients undergoing chemotherapy. *Gastroenterology* 2003; 125: 1742–1749.
  65. Huang H, Li X, Zhu J, *et al.* Entecavir vs lamivudine for prevention of hepatitis B virus reactivation among patients with untreated diffuse large B-cell lymphoma receiving R-CHOP chemotherapy. *JAMA* 2014; 312: 2521.

66. Pyrasopoulou A, Douma S, Vassiliadis T, *et al.* Reactivation of chronic hepatitis B virus infection following rituximab administration for rheumatoid arthritis. *Rheumatol Int* 2011; 31: 403–404.
67. Hui C-K, Cheung WWW, Zhang H-Y, *et al.* Kinetics and risk of de novo hepatitis B infection in HBsAg-negative patients undergoing cytotoxic chemotherapy. *Gastroenterology* 2006; 131: 59–68.
68. Seto W-K, Chan TSY, Hwang Y-Y, *et al.* Hepatitis B reactivation in patients with previous hepatitis B virus exposure undergoing rituximab-containing chemotherapy for lymphoma: a prospective study. *J Clin Oncol* 2014; 32: 3736–3743.
69. Hsu C, Tsou H-H, Lin S-F, *et al.* Chemotherapy-induced hepatitis B reactivation in lymphoma patients with resolved HBV infection: a prospective study. *Hepatology* 2014; 59: 2092–2100.
70. Koo YX, Tay M, Teh YE, *et al.* Risk of hepatitis B virus (HBV) reactivation in hepatitis B surface antigen negative/hepatitis B core antibody positive patients receiving rituximab-containing combination chemotherapy without routine antiviral prophylaxis. *Ann Hematol* 2011; 90: 1219–1223.
71. Huang Y-H, Hsiao L-T, Hong Y-C, *et al.* Randomized controlled trial of entecavir prophylaxis for rituximab-associated hepatitis B virus reactivation in patients with lymphoma and resolved hepatitis B. *J Clin Oncol* 2013; 31: 2765–2772.
72. Mitroulis I, Hatzara C, Kandili A, *et al.* Long-term safety of rituximab in patients with rheumatic diseases and chronic or resolved hepatitis B virus infection. *Ann Rheum Dis* 2013; 72: 308–310.
73. van Vollenhoven RF, Fleischmann RM, Furst DE, *et al.* Longterm safety of rituximab: final report of the rheumatoid arthritis global clinical trial program over 11 years. *J Rheumatol* 2015; 42: 1761–1766.
74. Varisco V, Viganò M, Batticciotto A, *et al.* Low risk of hepatitis B virus reactivation in HBsAg-negative/anti-HBc-positive carriers receiving rituximab for rheumatoid arthritis: a retrospective multicenter italian study. *J Rheumatol*. Epub ahead of print 15 February 2016. DOI: 10.3899/jrheum.151105.
75. Papalopoulos I, Fanouriakis A, Kougkas N, *et al.* Liver safety of non-tumour necrosis factor inhibitors in rheumatic patients with past hepatitis B virus infection: an observational, controlled, long-term study. *Clin Exp Rheumatol* 2018; 36: 102–109.
76. Tien Y-C, Yen H-H and Chiu Y-M. Incidence and clinical characteristics of hepatitis B virus reactivation in HBsAg-negative/HBcAb-positive patients receiving rituximab for rheumatoid arthritis. *Clin Exp Rheumatol* 35: 831–836.
77. Kim PS, Ho GY, Prete PE, *et al.* Safety and efficacy of abatacept in eight rheumatoid arthritis patients with chronic hepatitis B. *Arthritis Care Res* 2012; 64: 1265–1268.
78. Padovan M, Filippini M, Tincani A, *et al.* Safety of abatacept in rheumatoid arthritis with serologic evidence of past or present hepatitis B virus infection. *Arthritis Care Res* 2016; 68: 738–743.
79. Fanouriakis A, Vassilopoulos D, Repa A, *et al.* Hepatitis B reactivation following treatment with abatacept in a patient with past hepatitis B virus infection. *Rheumatology* 2014; 53: 195–196.
80. Germanidis G, Hytioglou P, Zakalka M, *et al.* Reactivation of occult hepatitis B virus infection, following treatment of refractory rheumatoid arthritis with abatacept. *J Hepatol* 2012; 56: 1420–1421.
81. Watanabe T, Fukae J, Fukaya S, *et al.* Incidence and risk factors for reactivation from resolved hepatitis B virus in rheumatoid arthritis patients treated with biological disease-modifying antirheumatic drugs. *Int J Rheum Dis* 2019; 22: 574–582.
82. Winthrop KL, Mariette X, Silva JT, *et al.* ESCMID Study Group for Infections in Compromised Hosts (ESGICH) Consensus Document on the safety of targeted and biological therapies: an infectious diseases perspective (Soluble immune effector molecules [II]: agents targeting interleukins, immunoglobuli. *Clin Microbiol Infect* 2018; 24: S21–S40.
83. Nagashima T and Minota S. Long-term tocilizumab therapy in a patient with rheumatoid arthritis and chronic hepatitis B. *Rheumatology* 2008; 47: 1838–1840.
84. Tsuboi H, Tsujii A, Nampei A, *et al.* A patient with rheumatoid arthritis treated with tocilizumab together with lamivudine prophylaxis after remission of infliximab-reactivated hepatitis B. *Mod Rheumatol* 2011; 21: 701–705.
85. Kishida D, Okuda Y, Onishi M, *et al.* Successful tocilizumab treatment in a patient with adult-onset Still's disease complicated by chronic



- active hepatitis B and amyloid A amyloidosis. *Mod Rheumatol* 2011; 21: 215–218.
86. Chen L-F, Mo Y-Q, Jing J, *et al.* Short-course tocilizumab increases risk of hepatitis B virus reactivation in patients with rheumatoid arthritis: a prospective clinical observation. *Int J Rheum Dis* 2017; 20: 859–869.
  87. Ahn SS, Jung SM, Song JJ, *et al.* Safety of tocilizumab in rheumatoid arthritis patients with resolved hepatitis B virus infection: data from real-world experience. *Yonsei Med J* 2018; 59: 452–456.
  88. Nakamura J, Nagashima T, Nagatani K, *et al.* Reactivation of hepatitis B virus in rheumatoid arthritis patients treated with biological disease-modifying antirheumatic drugs. *Int J Rheum Dis* 2014; 19(5): 470–475.
  89. Snast I, Atzmony L, Braun M, *et al.* Risk for hepatitis B and C virus reactivation in patients with psoriasis on biologic therapies: a retrospective cohort study and systematic review of the literature. *J Am Acad Dermatol* 2017; 77: 88–97.e5.
  90. Yanagihara S, Sugita K, Yoshida Y, *et al.* Psoriasis vulgaris in a hepatitis B virus carrier successfully treated with secukinumab and entecavir combination therapy. *Eur J Dermatol* 2017; 27: 185–186.
  91. Feaster B, Cline A and Feldman SR. Secukinumab for psoriasis in a patient with hepatitis B. *Dermatol Online J* 2018; 24: 4–6.
  92. Peccerillo F, Odorici G, Pellacani G, *et al.* Secukinumab: a positive outcome in a patient with severe psoriasis and HBV-HCV co-infection. *Dermatol Ther* 2018; 31: 1–2.
  93. Lora V, Graceffa D, De Felice C, *et al.* Treatment of severe psoriasis with ixekizumab in a liver transplant recipient with concomitant hepatitis B virus infection. *Dermatol Ther*; 32. Epub ahead of print 23 May 2019. DOI: 10.1111/dth.12909.
  94. Koike Y, Fujiki Y, Higuchi M, *et al.* An interleukin-17 inhibitor successfully treated a complicated psoriasis and psoriatic arthritis patient with hepatitis B virus infection and end-stage kidney disease on hemodialysis. *JAAD Case Reports* 2019; 5: 150–152.
  95. Chiu H, Hui R, Huang Y, *et al.* Safety Profile of secukinumab in treatment of patients with psoriasis and concurrent hepatitis B or C: a multicentric prospective cohort study. *Acta Derm Venereol* 2018; 98: 829–834.
  96. Chiu H-Y, Chen C-H, Wu M-S, *et al.* The safety profile of ustekinumab in the treatment of patients with psoriasis and concurrent hepatitis B or C. *Br J Dermatol* 2013; 169: 1295–1303.
  97. Ting S-W, Chen Y-C and Huang Y-H. Risk of hepatitis B reactivation in patients with psoriasis on ustekinumab. *Clin Drug Investig* 2018; 38: 873–880.
  98. Schwartz DM, Kanno Y, Villarino A, *et al.* JAK inhibition as a therapeutic strategy for immune and inflammatory diseases. *Nat Rev Drug Discov* 2017; 17: 78.
  99. Chen Y-M, Huang W-N, Wu Y-D, *et al.* Reactivation of hepatitis B virus infection in patients with rheumatoid arthritis receiving tofacitinib: a real-world study. *Ann Rheum Dis* 2018; 77: 780–782.
  100. Harigai M, Winthrop K, Takeuchi T, *et al.* FRI0077 Hepatitis b virus reactivation in patients with rheumatoid arthritis treated with baricitinib: post-hoc analysis from clinical trials. *Ann Rheum Dis* 2018; 77: 584–585.
  101. Bart PA, Jacquier P, Zuber PL, *et al.* Seroprevalence of HBV (anti-HBc, HBsAg and anti-HBs) and HDV infections among 9006 women at delivery. *Liver* 1996; 16: 110–116.
  102. De Paschale M, Manco MT, Belvisi L, *et al.* Prevalence of markers of hepatitis B virus infection or vaccination in HBsAg-negative subjects. *Blood Transfus* 2012; 10: 344–350.
  103. Cho Y, Yu SJ, Cho EJ, *et al.* High titers of anti-HBs prevent rituximab-related viral reactivation in resolved hepatitis B patient with non-Hodgkin's lymphoma. *J Med Virol* 2016; 88: 1010–1017.
  104. Kotake T, Satake H, Okita Y, *et al.* Prevalence and risk factors of hepatitis B virus reactivation in patients with solid tumors with resolved HBV infection. *Asia Pac J Clin Oncol* 2019; 15: 63–68.
  105. Paul S, Dickstein A, Saxena A, *et al.* Role of surface antibody in hepatitis B reactivation in patients with resolved infection and hematologic malignancy: a meta-analysis. *Hepatology* 2017; 66: 379–388.
  106. Tien Y-C, Hsue Y-T, Hung M-H, *et al.* Changes in hepatitis B virus surface antibody titer and risk of hepatitis B reactivation in HBsAg-negative/HBcAb-positive patients undergoing biologic therapy for rheumatic diseases: a prospective cohort study. *Arthritis Res Ther* 2018; 20: 246.
  107. Cholongitas E, Haidich AB, Apostolidou-Kiouti F, *et al.* Hepatitis b virus reactivation in HBsAg-negative, anti-HBc-positive patients receiving immunosuppressive therapy: a systematic review. *Ann Gastroenterol* 2018; 31: 480–490.



108. Sampedro B, Hernández-López C, Ramón J, *et al.* A computerized physician order entry-based system to prevent HBV reactivation in patients treated with biologic agents. The PRESCRIB Project. *Hepatology* 2014; 106–113.
109. Konijeti GG, Grandhe S, Konerman M, *et al.* Cost-effectiveness analysis of screening for hepatitis B virus infection in patients with solid tumors before initiating chemotherapy. *Clin Gastroenterol Hepatol*. Epub ahead of print 31 October 2019. DOI: 10.1016/j.cgh.2019.10.039.
110. Crespo J, Esteban R, Torres C, *et al.* Cost-effectiveness of a hepatitis B virus screening strategy to prevent reactivation in patients with hematologic neoplasms. *Rev Esp Enferm Dig* 2017; 109: 619–626.
111. Hwang JP, Huang D, Vierling JM, *et al.* Cost-effectiveness analysis of hepatitis B virus screening and management in patients with hematologic or solid malignancies anticipating immunosuppressive cancer therapy. *JCO Clin Cancer Informatics* 2019; 3: 1–12.
112. Mast EE, Weinbaum CM, Fiore AE, *et al.* A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP) Part II: immunization of adults. *MMWR Recomm reports* 2006; 55: 1–33; quiz CE1–4.
113. Thomas K and Vassilopoulos D. Immunization in patients with inflammatory rheumatic diseases. *Best Pract Res Clin Rheumatol* 2016; 30: 946–963.
114. Kim DK, Hunter P, Woods LD, *et al.* Recommended adult immunization schedule, United States, 2019. *Ann Intern Med* 2019; 170: 182–192.
115. Schillie S, Harris A, Link-Gelles R, *et al.* Recommendations of the Advisory Committee on Immunization Practices for Use of a Hepatitis B Vaccine with a Novel Adjuvant. *MMWR* 2018; 67: 455–458.
116. Jackson S, Lentino J, Kopp J, *et al.* Immunogenicity of a two-dose investigational hepatitis B vaccine, HBsAg-1018, using a toll-like receptor 9 agonist adjuvant compared with a licensed hepatitis B vaccine in adults. *Vaccine* 2018; 36: 668–674.

Visit SAGE journals online  
journals.sagepub.com/  
home/tab

 SAGE journals