

Placenta-Derived Cells for Acute Brain Injury

Cell Transplantation
2018, Vol. 27(1) 151–167
© The Author(s) 2018
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0963689717732992
journals.sagepub.com/home/ccl


Francesca Pischiutta¹, Eliana Sammali^{1,2}, Ornella Parolini^{3,4},
Hilary V. O. Carswell⁵, and Elisa R. Zanier¹

Abstract

Acute brain injury resulting from ischemic/hemorrhagic or traumatic damage is one of the leading causes of mortality and disability worldwide and is a significant burden to society. Neuroprotective options to counteract brain damage are very limited in stroke and traumatic brain injury (TBI). Given the multifaceted nature of acute brain injury and damage progression, several therapeutic targets may need to be addressed simultaneously to interfere with the evolution of the injury and improve the patient's outcome. Stem cells are ideal candidates since they act on various mechanisms of protection and repair, improving structural and functional outcomes after experimental stroke or TBI. Stem cells isolated from placenta offer advantages due to their early embryonic origin, ease of procurement, and ethical acceptance. We analyzed the evidence for the beneficial effects of placenta-derived stem cells in acute brain injury, with the focus on experimental studies of TBI and stroke, the engineering strategies pursued to foster cell potential, and characterization of the bioactive molecules secreted by placental cells, known as their secretome, as an alternative cell-free strategy. Results from the clinical application of placenta-derived stem cells for acute brain injury and ongoing clinical trials are summarily discussed.

Keywords

placenta-derived stem cells, amnion-derived stem cells, umbilical cord-derived stem cells, traumatic brain injury, stroke, clinical applications, regenerative medicine.

The Rationale for Using Placenta-Derived Cells for Acute Brain Injury

Acute brain injury, resulting from traumatic brain injury (TBI), ischemic stroke, or intracerebral hemorrhage (ICH), is associated with high rates of short-term mortality and long-term disability worldwide¹. Independent of its etiology (vascular occlusion, bleeding, or mechanical injury), brain damage is amplified after the primary insult, with the activation of genomic, cellular, and/or biochemical processes that interact in a complex network leading to delayed cellular dysfunction and death. Although the extensive effects of these events and the abundance of targets offer the potential for therapeutic interventions, so far there is no pharmacological treatment that can reverse the pathologic cellular cascades and improve the outcome. Thus, there is a pressing need for therapeutic approaches aimed at protecting and repairing the injured brain.

Stem cells, with their ability to act simultaneously on multiple targets, driving the damaged microenvironment from a toxic to a more protective/regenerative activation state, are a promising strategy^{2–6}. Stem cells have been isolated

from almost all body organs, raising the question of the best source for each specific pathology, since heterogeneous effects have been observed depending on the source.

¹ Department of Neuroscience, Laboratory of Acute Brain Injury and Therapeutic Strategies, IRCCS Istituto di Ricerche Farmacologiche Mario Negri, Milan, Italy

² Department of Cerebrovascular Diseases, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy

³ Centro di Ricerca E. Menni, Fondazione Poliambulanza Istituto Ospedaliero, Brescia, Italy

⁴ Institute of Anatomy and Cell Biology, Università Cattolica del Sacro Cuore, Rome, Italy

⁵ Strathclyde Institute of Pharmacy and Biomedical Sciences (SIPBS), University of Strathclyde, Glasgow, United Kingdom

Submitted: March 30, 2017. Revised: July 31, 2017. Accepted: August 01, 2017.

Corresponding Author:

Elisa R. Zanier, Department of Neuroscience, Laboratory of Acute Brain Injury and Therapeutic Strategies, IRCCS Istituto di Ricerche Farmacologiche Mario Negri, via Giuseppe La Masa 19, 20156 Milan, Italy. Email: elisa.zanier@marionegri.it






Figure 1. Specific subtypes of placenta-derived cells used for experimental acute brain injury.

According to their tissue origin, stem cells display specific differences at transcriptional and proteomic levels^{7,8}. Moreover, aging negatively affects stem/progenitor cell properties^{9,10}, reducing their proliferation and differentiation capacity^{11,12}, their immunomodulatory effects¹³, and their therapeutic potential^{14,15}. Thus, stem/progenitor cells from placenta are of special interest in a translational perspective, mainly on account of their reduced immunogenicity and high immunomodulatory potential and their broad plasticity^{16,17}.

The placenta is a fetomaternal organ. The fetal placental tissues include the amniotic and chorionic membranes, the umbilical cord, and the chorionic villi. The maternal component is known as decidua^{16,18}. All these placental tissues/fluids are easily available after birth, without invasive procedures, and their use is free from ethical issues.

This review examines the evidence for placenta-derived stem cell therapy for acute brain injury in experimental settings, with a focus on TBI, stroke, and ICH. We not only discuss efficacy but also the mechanisms of action and the engineering strategies used to amplify stem cell potency. One section will deal with the discussion of the bioactive molecules secreted by placental cells and the potential use of the secretome as a cell-free, better defined therapeutic strategy. Lastly, results from the clinical application of placenta-derived stem cells for acute brain injury and ongoing clinical trials will be discussed.

This is a narrative, nonsystematic, comprehensive review of the literature. The search string run in the PubMed database in January 2017 was as follows: (“traumatic brain injury” OR “brain trauma” OR “head trauma”) OR (“Brain Ischemia”[Mesh] OR “Stroke”[Mesh] OR stroke OR “brain ischemia”) AND (“Stem Cells”[Mesh] OR “stem cells” OR “stem cell” OR “stromal cells” OR “stromal cell”) AND (“placenta” OR “amnion” OR “amniotic” OR “cord blood” OR “umbilical” OR “chorion” OR “chorionic” OR “trophoblasts” OR “decidua”). We screened 301 papers for

title, abstract, and full text, finally selecting 139 original articles and 60 reviews. Five of the original articles were clinical studies and 134 experimental, of which 21 focused on TBI, 93 on cerebral ischemia, 16 on neonatal/infant stroke, and 4 on ICH. Among papers on stroke, those using models of heatstroke or diabetes were outside the remit of this review and were therefore not used. Thirty-five papers were added for analysis of the secretome, based on our own files or reference lists of previously reviewed articles.

In the majority of studies, cells were of human (h) origin, with very few using rat (r) or mouse (m) cells. Cells were isolated from different placental tissues and fluids including umbilical cord blood (UCB), umbilical cord tissue (UCT), amnion (A) and amniotic fluid (AF). The specific subtypes of cells used in the field of acute brain injury are listed in Figure 1. We found no study using cells from chorion, trophoblasts, or decidua.

Placenta-Derived Cell Transplantation for TBI

The literature concerning the use of placenta-derived stem cells in experimental models of TBI shows more widespread use of cells isolated from the UCB or tissue^{19–27} than cells from the amnion or AF^{28,29}. No papers were found using cells from chorion, trophoblasts, or decidua.

The first paper using human umbilical cord blood stem cells (hUCB-SCs) was published by Lu and colleagues¹⁹ who used 2 million hUCB-SCs, without selecting any specific population. Cells were intravenously infused in rats, 24 h after controlled cortical impact (CCI) brain injury and improved sensorimotor function from 2 wk post-TBI¹⁹. After this first experiment, the use of placenta-derived stem cells focused on investigation of the effects of specific subpopulations (endothelial, hematopoietic, or mesenchymal stromal cells [MSCs]) isolated from the hUCB.

The endothelial progenitors were selected with the aim of promoting vascular repair and stimulating neoangiogenic processes. Both systemic²² and local (intracerebroventricular [ICV] infusion)²³ transplantation of hUCB-derived endothelial progenitor cells (hUCB-EPCs) in traumatized rodents can increase microvascularization in the injured brain and angiogenic processes^{22,23}, preserving blood–brain barrier (BBB) integrity²³ and reducing brain edema²³. ICV-transplanted hUCB-EPCs integrated into the brain microvasculature²³, but when intravenously (IV) infused, the number of integrated hUCB-EPCs was scanty and the neovascularization was mostly due to the release of proangiogenic factors²².

Similar results on the vascular compartment have been obtained with hUCB-derived hematopoietic subpopulations (CD34⁺ or CD45⁺)^{21,24}, with an increase in cerebral perfusion pressure²¹, stimulation of vascular endothelial growth factor (VEGF) production²⁴, and increased angiogenesis²¹. Besides vascular effects, hUCB-CD34⁺ cells also stimulated neurogenesis, with increases in the amounts of newly generated neurons (BrdU⁺/NeuN⁺ cells), induced anti-inflammatory effects, lowering serum levels of proinflammatory cytokines (tumor necrosis factor α [TNF- α] and intercellular adhesion molecule 1 (Icam-1)) and increasing the anti-inflammatory ones (interleukin [IL] 10), and decreased toxic oxidizing radicals (NO_x)²¹. Overall functional improvement in both sensorimotor^{21–24} and cognitive^{22,23} deficits induced by hUCB endothelial or hematopoietic cells has been observed, together with a reduction in contusion volume^{21,22,24}. It is not clear, however, whether these nonvascular effects are directly promoted by the transplanted cells or are an indirect consequence of the vascular rescue.

The other subpopulation widely used for TBI are MSCs isolated from hUCB²⁰ or hUCT^{25–27,30}. Since the data available do not allow for a distinction of enhanced efficacy of the MSCs depending on the different regions of the umbilical cord, they are collectively referred to as umbilical-MSCs. Systemic^{25,27} and local^{20,26} transplantation of MSCs in TBI rodents was able to induce functional improvement in sensorimotor^{20,25–27} and cognitive^{20,26,27} deficits and reduce the contusion volume^{20,26}. Transplanted umbilical-MSCs were found in the lesion site and their migration toward the injured tissue seemed to be driven by the stromal cell-derived factor 1/ C-X-C chemokine receptor type 4 (SDF-1/CXCR4) pathway²⁵; however, the number of cells surviving after transplantation gradually drops²⁰.

The various protective mechanisms elicited by transplanted cells involve different pathways. MSCs reduce brain edema²⁶, lower the number of apoptotic cells in the damaged area²⁵, boost the expression of neurotrophic factors^{20,26}, stimulate endogenous neurogenesis^{25–27}, and have immunoregulatory effects, reducing serum markers of proinflammatory mediators (IL-1, interferon [IFN] γ , anti-brain antibody) and increasing anti-inflammatory ones (IL-10 and transforming growth factor β)³⁰. Umbilical-MSCs also shift activated microglial cells in the pericontusional

tissue toward a nonphagocytic phenotype with protective and healing functions²⁰.

The amniotic source for stem cell therapy has been less investigated for TBI, with only 2 papers published^{28,29}. Chen and colleagues used human amnion-derived multipotent progenitor cells (hAMPCs) and transplanted 2 million cells ICV immediately after penetrating ballistic-like brain injury in rats²⁸. Transplanted cells reduced axonal degeneration in the corpus callosum and in the thalamus. Labeled hAMPCs migrated from the subventricular zone to the corpus callosum but not to the thalamus, and there was no evidence of differentiation toward neuronal lineage, suggesting that the effects of these cells may result from a continuous supply of secreted bioactive factors²⁸.

Our group has provided evidence of the protective effects of human amnion membrane-derived MSCs (hAMSCs) in a mouse model of TBI modeled by controlled cortical impact (CCI)²⁹. We compared the effects of ICV and IV transplanted cells 24 h after TBI in mice. Only ICV transplanted cells were found in the brain at acute stages after injury, but ICV and IV transplanted hAMSCs induced functional improvement in sensorimotor deficits to similar extents. Functional improvement was associated with histological protective modifications, such as neuronal and vessel rescue in the pericontusional cortex and the stimulation of endogenous neurogenesis²⁹.

To summarize, the literature regarding the use of placenta-derived cell therapy for TBI varies widely, with different tissues of origin, isolation protocols, and subpopulations. Although it is hard to compare the therapeutic effects obtained in different laboratories, it is clear that infused cells have pleiotropic action on multiple targets, inducing protective/regenerative environmental changes and stimulating endogenous neuroprotective mechanisms by autocrine/paracrine factors. The presence of cells in the lesioned brain seems to be unnecessary to confer the protection, thus posing the basis for the development of a cell-free strategy using the stem cell secretome. This topic will be discussed in the section “Placenta-Derived Cell Secretome: Toward Cell-Free Therapy.”

Placenta-Derived Cell Transplantation for Stroke

The first preclinical use of placenta-derived cells for stroke dates back to 2001^{31,32}. Chen and colleagues reported that IV infused hUCB-SCs after middle cerebral artery occlusion (MCAO) in rats improved functional recovery³¹, and Okawa and colleagues described the permanence of hippocampal transplanted rat-derived AECs at the site of injection, 5 wk after stroke in gerbils³². Since then several studies have explored the efficacy and mechanisms of action of placenta-derived cells after brain ischemia. As with TBI, we also noted for stroke that cells derived from an umbilical source have been investigated far more than amniotic cells,

and we did not find any studies that used cells isolated from the chorion, trophoblast, or decidua.

The mechanisms of hUCB cell-induced recovery after stroke have been investigated with the aim of establishing whether transplanted cells must actually enter the brain to be effective. When hUCB-SCs were IV infused 24 to 48 h after MCAO, viable cells were found in the brain and improved functional recovery was reported^{31,33}. Efficiency was dose dependent and higher when compared to intracranial administration at the same time point³³. This dose-related effect was also found for hUCT-derived cells³⁴. The ability of hUCT-MSCs to differentiate into neural phenotypes has been shown *in vitro*³⁵; however, transdifferentiation of hUCT-MSCs into fully functional neurons after *in vivo* acute brain injury has never been demonstrated and mechanisms other than engraftment appear to mediate the restorative effects after hUCT-MSC infusion^{36,37}.

Despite the above evidence, along with the observation that hUCB-SCs have potent migratory capacity in response to ischemic brain injury³⁸, probably through chemokine SDF-1/CXCL12³⁹, there is evidence that systemically administered cells do not need to enter the brain to induce functional recovery after MCAO⁴⁰. Several studies indicate that hUCB cells induce restorative effects after stroke through the secretion of trophic factors⁴¹. In addition, hUCT-MSCs secrete human brain-derived neurotrophic factor (BDNF), neutrophil-activating protein-2 (NAP-2), angiopoietin-2 (Ang-2), CXCL16, platelet-derived growth factor-AA (PDGF-AA), basic fibroblast growth factor (hVEGFR-3), and human vascular endothelial growth factor receptor-3³⁷ (see also "Placenta-Derived Cell Secretome: Toward Cell-Free Therapy" section). Although most studies show these trophic factors induce angiogenesis, neurogenesis, and synaptogenesis at the site of injury after adult and perinatal ischemia^{34,42,43}, the role of hUCB-SCs in restoring spleen weight and splenic CD8⁺ T-cell counts after MCAO has been highlighted⁴⁴, indicating a role in dampening the systemic inflammatory response after stroke. The immunomodulatory effects of hUCB cells are established in other inflammatory conditions and are potential mechanisms for mediating restoration after stroke. hUCB-SCs were able to reduce ischemia-induced infiltration of granulocytes, monocytes, and CD3⁺ T cells and activation of astrocytes and microglia in adults^{45–47} and pups⁴⁸, in addition to reducing proinflammatory cytokines and nuclear factor κ B activity⁴⁶ after stroke, all of which may contribute to the resolution of the inflammatory response after stroke and facilitate postischemic plasticity.

In terms of the subpopulations, hUCB-CD34⁺ cells have been prominently investigated in stroke models and have been shown to reduce brain damage in newborns⁴⁹ and to induce behavioral recovery^{50,51} which is enhanced in CD34⁺ cells overexpressing glial cell line-derived neurotrophic factor (GDNF)⁵². hUCB-CD34⁺ cells have also been shown to be responsible for the angiogenic and neurogenic effects observed after hUCB-SC infusion⁴², although hUCB-

CD34⁻ cells were just as effective⁵⁰. Boltze and colleagues not only found that hUCB-CD34⁺ and CD34⁻ cells had similar behavioral effects, but also that human umbilical cord blood-derived mononuclear cells (hUCB-MNCs) were more effective⁵³. Among the different cells in hUCB-MNCs (immature T and B cells, monocytes, and stem cells), monocytes were found to have the most marked effects on improving functional recovery and reducing infarct⁵⁴. Several studies have also indicated the therapeutic benefits of hUCB- or hUCT-derived MSCs after stroke. hUCB-MSCs reduced infarction lesion size after cerebral ischemia in adult rats⁵⁵ and dogs⁵⁶ as well as rat neonates^{57,58}. hUCT-MSCs also improved functional recovery in rats after MCAO³⁷, possibly by reducing peripheral and cerebral proinflammatory cytokines, an observation reported for both hUCB-MSCs^{57,59} and hUCT-MSCs⁶⁰.

Despite this wealth of evidence on the comprehensive efficacy of hUCB cells in stroke, a number of studies show a lack of effects^{61–63}, so further research into different storage and processing conditions is still needed to confirm the therapeutic potential of hUCB- and hUCT-derived cells in stroke.

As regards amniotic cells, the epithelial population amniotic epithelial cells (AECs), isolated from rat³² or human⁶⁴ tissues, can migrate to the site of ischemic injury³² and improve functional recovery after MCAO⁶⁴. hAMSCs can also improve functional recovery after MCAO^{65,66}, accelerated by overexpression of BDNF⁶⁷.

AF stem cells were also shown to promote functional recovery after MCAO⁶⁸, reducing ischemic damage and promoting neurogenesis⁶⁹, and have been used for reprogramming into induced pluripotent stem cells (iPSCs)⁷⁰.

To summarize, the current evidence for the potential of placenta-derived cell therapy for stroke is promising. Despite the different protocols and subpopulations tested, and the consensus needed on optimum storage and processing conditions, the literature indicates that placenta-derived stem cells improve functional recovery after MCAO, most likely by promoting angiogenesis and neurogenesis and through potent immunomodulatory effects that contribute to the resolution of the inflammatory response after stroke and facilitate postischemic plasticity.

Placenta-Derived Cell Transplantation for ICH

In the experimental models of ICH, hUCB and hUCT are the only placental sources that have been investigated^{71–73}. hUCB-MNCs were tested in a dose-dependent study⁷¹ in order to study the effects of 4, 8, or 16 million IV infused cells 1 d after intrastriatal ICH. All transplanted groups showed similar improvement in sensorimotor function after 1 wk. Cells migrated selectively to the hemorrhagic area in the right striatum, with more cells found in the highest dose group. Similarly, the lesion volume decreased in all transplanted groups compared to control, with a dose-dependent effect.

Two studies investigated the effects of hUCB-MSCs⁷² or hUCT-MSCs⁷³. ICV infusion of 500,000 hUCB-MSCs, 2 d after intrastriatal ICH, improved sensorimotor function, with a reduction in the lesion volume 4 wk after transplant⁷². Cells were found in proximity to the hemorrhagic area 3 d after injection, but they disappeared at later time points (4 wk), with a nonsignificant tendency toward reduction in proinflammatory markers (TNF- α , cyclooxygenase-2 (COX-2) microglial activation, and neutrophil infiltration). Xie and colleagues compared the effect of intracerebrally (200,000 cells) or IV (2 million cells) infused hUCT-MSCs after intrastriatal ICH, observing similar protection on sensorimotor deficits and lesion volume⁷³. Four weeks later, intracerebrally transplanted hUCT-MSCs were still clustered in the injection site, near the hemorrhagic region, while only a small part of the IV transplanted cells could be found in the brain. Only rats that intracerebrally received hUCT-MSCs had enhanced vascular density in the peri-ICH regions of the ipsilateral hemisphere, which was attributed to the limited number of IV transplanted cells that migrated to the damaged tissue. Given the similarity of the functional benefit, it may well be that locally and systemically transplanted cells act through different mechanisms and that combining the two may exert synergistic effects.

Manipulation of Placenta-Derived Stem Cells and/or the Use of Survival-Promoting Scaffolds

To optimize the therapeutic efficacy of transplanted cells, different combinatory strategies have been employed, including the use of scaffolds (spatially guiding tissue regeneration), the genetic modification (to overexpress specific factors or promote survival), and delivery of different biomolecules (trophic factors, modulators of inflammatory molecules).

With the aim of enhancing cell engraftment and differentiation, *in vitro* neuronal commitment prior to *in vivo* transplantation has also been tried^{74,75}. hUCT-MSCs untransdifferentiated or transdifferentiated into neural-like cells (hUCT-MSC-NSCs) were resuspended in 10 μ L Matrigel and transplanted into the ipsilateral hippocampus of rats, 7 d after TBI induced by weight drop impact model⁷⁴. Only 6% to 7% of the transplanted cells were found in the brain 28 d after transplantation, with no differences in survival rates between hUCT-MSCs and hUCT-MSC-NSCs. Both treated groups showed a reduction in contusion volume and an improvement in cognitive deficits compared to control groups. However, untransdifferentiated hUCT-MSCs induced a better outcome than hUCT-MSC-NSCs. *In vitro* experiments showed that hUCT-MSCs secrete more neurotrophic factors (BDNF and neurotrophin-3 [NT-3]) than hUCT-MSC-NSCs, suggesting that transdifferentiation before use may not provide any advantage.

Two years later, the same authors did a similar analysis using hAMSCs undifferentiated or *in vitro* transdifferentiated

into neural stem-like cells (hAMSC-NSCs)⁷⁵. Cells were transplanted in TBI rats using a similar protocol, with the only exception that cells were infused earlier (4 d). Both transplanted groups showed improvement in sensorimotor and cognitive functions compared to controls, but hAMSC-NSC transplantation resulted in significant enhancement of sensorimotor function compared with hAMSC-treated rats. Unlike in the previous study, hAMSC-NSCs produced higher levels of neurotrophic factors (nerve growth factor [NGF], BDNF, NT-3, GDNF, ciliary neurotrophic factor [CNTF]) than hAMSCs, again suggesting that cell replacement is unlikely. Instead, a neuroprotective effect, associated with neurotrophic factors released by the grafted cells, may contribute to functional recovery.

In a similar study where 20,000 neural stem cells differentiated from hUCB (hUCB-NSCs) were intracerebrally infused 48 h after photothrombotic stroke, extensive migration of the cells into brain-damaged cortex in the first week after transplant was observed. However, differentiation was observed in very few cells and cell survival in the brain decreased gradually⁷⁶. Again, this suggests that the persistence of cells in the host brain is not required and differentiation does not seem the main mechanism of action.

Therapeutic hypothermia can reduce functional deficits and limit secondary damage in models of TBI⁷⁷. In order to potentially enhance therapeutic efficacy, Tu and colleagues combined hypothermia and hUCT-MSC treatments in a rat model of severe TBI⁷⁸. Temperature-sensitive hUCT-MSCs were generated in order to promote the survival and the activity of grafted cells in the hypothermic rats by infecting cells with a retrovirus carrying the temperature-sensitive tsA58 SV40 LT antigen gene. Hypothermia contributed to the survival, migration, and proliferation of ts-hUCT-MSCs up to 28 d after injection in the lesion cavity, promoting the improvement in sensorimotor and cognitive functions more than hUCT-MSC therapy alone⁷⁸.

Another approach used to establish an optimal environment to support lasting viability of stem cells *in situ* is the use of engineered scaffolds, which make it possible to deliver cells appropriately to the injured site, optimizing their therapeutic effects. Huang and colleagues combined hUCT-MSCs overexpressing CXCR4 fused to green fluorescent protein (hUCT-MSC^{CXCR4/GFP}), responsible for cell migration through SDF-1 signaling and usually present at low levels on MSC membrane, with a chitosan scaffold linked to recombinant BDNF⁷⁹. The scaffold containing hUCT-MSC^{CXCR4/GFP} was transplanted in the lesion cavity of TBI rats. CXCR4 overexpression stimulated hUCT-MSC migration toward the lesion site, and the BDNF released from scaffolds favored the differentiation of transplanted cells into neurons, contributing to the regeneration of brain tissue at the lesion boundary.

The same researchers developed another type of scaffold, RADA16-BDNF, a self-assembling peptide hydrogel that favored the differentiation of hUCT-MSCs into MAP-2-positive neuronal cells⁸⁰. When hUCT-MSCs were




Figure 2. Categories of molecules secreted by placenta-derived stem cells.

cocultured with lipopolysaccharide (LPS)-activated astrocytes, the percentage of MAP-2-positive cells was higher and typical neuron-like cells with neurites extending in 3-D directions differentiated from hUCT-MSCs. RADA16-BDNF scaffolds embedded with hUCT-MSC^{CXCR4/GFP} and LPS-activated astrocytes were then transplanted into the lesion cavity of rats subjected to moderate (2-mm-diameter) and large (5-mm-diameter) lesions. hUCT-MSC^{CXCR4/GFP} migrated from the scaffold to the injured tissue, with a lesion size-dependent efficacy in reducing the lesion size⁸⁰.

Transgenically modified hUCT-MSCs engineered to overexpress hepatocyte growth factor (hUCT-MSCs^{HGF}) have been used to enhance the therapeutic potential for ICH damage⁸¹. hUCT-MSCs or hUCT-MSCs^{HGF} (600,000) were transplanted ICV 7 d after ICH, modeled by collagenase VII injected into the left internal capsule. Both transplanted groups showed better motor performance than untreated animals, and hUCT-MSC^{HGF}-transplanted rats demonstrated a greater functional recovery than hUCT-MSC-transplanted rats. Neurological recovery was coupled with nerve fiber remyelination (upregulation of myelin basic protein (MBP)) and axonal regeneration (upregulation of growth-associated protein 43 (GAP-43)), demonstrating the pivotal role of growth factors in placenta-derived stem cell-induced protection.

To conclude, different strategies have been developed to enhance the therapeutic effects of transplanted cells. Understanding the mechanisms of protection induced by placenta-derived stem cells will help identify the best strategy for maximizing their effect.

Placenta-Derived Cell Secretome: Toward Cell-Free Therapy

The beneficial effects of cell therapy for acute brain injury are mediated by the release of soluble factors with multilevel

effects providing support for surviving host cells, offering protection from toxic stimuli from the damaged brain tissue, and stimulating endogenous protective/repairative mechanisms. The view of secreted factors as mediators of brain protection raises the possibility of a cell-free therapeutic strategy, overcoming important issues related to intrinsic cell heterogeneity and safety concerns. *In vitro*^{29,82–86} and *in vivo*^{86–88} evidence shows that the conditioned medium (CM) from placenta-derived cell cultures can induce effects similar to the cell counterparts, demonstrating the feasibility of this approach. The CM composition is influenced by different factors: besides the organ of origin and the donor's genetic background, several variables (i.e., cell density, cell passage, culture medium, days of collection, presence of preconditioning stimuli) may affect the release of soluble molecules. We recently provided evidence of comparable protective effects of hAMSC and CM treatments, indicating that cross talk between hAMSCs and the damaged tissue is not vital for the release of bioactive factors²⁹. However, whether the *in vivo* challenge to the transplanted cells enhances or otherwise changes the release of their bioactive factors calls for further investigations.

A number of cell-released soluble factors have been proposed as mediators of the protection, and different classes of molecules have been investigated (Figure 2). Among the different classes of molecules that could be released in the secretome, the proteic component is the most widely investigated. A recent review by Bai and colleagues provided an overview of the secretion properties of umbilical-MSCs⁸⁹, identifying more than 90 cytokines belonging to the families of ILs, TNFs, IFNs, colony-stimulating factors, growth factors, and chemokines. These proteins have important potential interest for brain pathologies, in view of their numerous physiological functions, executing anti-inflammatory and immunomodulatory effects; arousing angiogenesis, wound healing, and neurogenesis; and stimulating antiapoptotic,

antiscarring, and matrix remodeling effects⁸⁹. Pires and colleagues also identified other classes of proteins released by UCT-MSCs involved in protection against oxidative stress, anti-excitotoxicity effects, proteasomal degradation, and regulators of toxic protein deposition⁹⁰.

Nucleic acids are another class of molecules that have recently aroused great interest. MicroRNAs (miRs) are small noncoding sequences of RNA that can regulate the gene expression of target cells and are being studied for therapy as they play a key role in pathology. Packaged inside extracellular vesicles, miR can be released into the extracellular space and mediate nonhormonal intracellular communication⁹¹. The miR profile of UCT-MSCs was recently characterized, with distinct expression from fibroblasts⁹². Forty-two miRs have been found to be expressed differently in UCT-MSCs and fibroblasts, and among the 15 that were upregulated, miR-21, miR-146a, and miR-181 displayed the most significant expression. Analysis of the downstream targets unraveled the involvement of these miRs in the inflammatory pathways, regulating macrophage phenotype and inflammation ablation for tissue repair^{93–95}. In the brain, miR-21 mediates neuronal and microglial survival after ischemic injury, reducing the levels of the apoptosis-inducing factor Fas ligand^{96,97}; miR-181 regulates the expression of IL-10 anti-inflammatory cytokines in astrocytes⁹⁸ and reduces the neuronal apoptosis induced by microglia activated after an ischemic insult, suppressing TNF- α expression⁹⁹.

Another hUCB-MSC-released miR involved in neurorestorative effects is miR-126¹⁰⁰, which is vital for regulating the function of endothelial cells, angiogenesis, and vascular integrity¹⁰¹. Chen and colleagues showed that hUCB-MSC treatment of stroke in type 2 diabetes mellitus mice restored the serum and cerebral levels of miR-126, which was downregulated after injury¹⁰⁰. Cell therapy promotes vascular remodeling and angiogenesis and reduced brain hemorrhage and BBB leakage. These effects were coupled with white matter remodeling, anti-inflammatory effects, and a generally better functional outcome. All of these effects were significantly attenuated in mice treated with hUCB-MSCs in which miR-126 expression was inhibited, indicating an important role for this miR in hUCB-MSC-induced protection¹⁰⁰.

Metabolites are small molecules (molecular weight <2,000 Da) that include endogenous compounds (peptides, lipids, amino acids, nucleic acids, carbohydrates, organic acids, vitamins) or exogenous compounds (food additives, drugs, toxins, pollutants, and other chemicals that humans come into contact with)¹⁰². It is not easy to define the metabolomics profile of human cells because thousands of compounds belong to this class. In an *in vitro* model of acute brain injury obtained by exposing cortical organotypic brain slices to oxygen-glucose deprivation, we have recently demonstrated that the CM from hAMSCs was protective, and we identified a CM subfraction containing small metabolites (molecular weight <700 Da) with marked protective

properties²⁹. We further analyzed the CM active subfraction, by profiling 6 metabolic classes, and identified a pool of selectively enriched metabolites. These included molecules such as lysine, taurine, α -amino adipic acid, and spermidine, for all of which neuroprotective effects have been reported^{103–107}. However, our metabolomics analysis was targeted *a priori*; thus, we cannot exclude that additional metabolites with protective potential are present in the CM subfraction. Another metabolite that is attracting interest for the protection induced by stem cells is prostaglandin E2 (PGE2). PGE2 is highly secreted by placenta-derived stem cells^{7,108–110}, and there is increasing evidence of its immunosuppressive effects on lymphocytes^{109,111–113} and monocytes/macrophages^{114,115} and a role for brain protection^{116–119}.

To conclude, the secretome of placenta-derived stem cells holds great promise for the treatment of acute brain injury. Different molecules have been identified as possible drivers of beneficial effects. The exact mechanisms of action are still not clear and numerous factors, rather than any single one, probably contribute synergistically to the neuroprotective effects.

Clinical Use of Placenta-Derived Cells in Acute Brain Injury

Placenta-derived stem cells are still only at their dawn in the clinical setting compared to the more widely used bone marrow-derived stem cells, so only a few studies investigating the feasibility and safety of this approach for acute brain injury have been conducted so far (Table 1).

hUCB-derived stem cells for acute brain injury were first used as an experimental neurorestorative treatment in a 16-mo-old child in a permanent vegetative state resulting from a cardiac arrest-induced global hypoxic/ischemic brain injury when he was 9 mo old¹²⁰. The child had his own hUCB deposited in a blood bank; autologous hUCB cells were isolated and committed to neuronal lineage (hUCB-NSCs), then labeled with supermagnetic iron oxide (SPIO) and ICV transplanted. The child underwent 4 transplantations of 12 million hUCB-NSCs each, the first was when he was 16 mo old, followed by 3 consecutive monthly injections. Magnetic resonance imaging (MRI) monitoring 24 h posttransplant showed hUCB-NSCs extending along the ependyma of the lateral ventricle. The SPIO signal progressively decreased but was still visible 7 d, 1 mo, and 2 mo after transplant, and disappeared by 4 mo. There were no signs of brain edema, hemorrhage, or tumor in the 6 mo of observation¹²⁰ or in the follow-up at 33 mo¹²¹. The only side effect was transient moderate fever the day after transplant. Thus, the authors affirm the feasibility and the safety of intracerebral transplant of hUCB-NSCs in the brain.

Jiang and colleagues ran a pilot study to assess the safety of hUCT-MSCs delivered by a catheter through perilesional vessels to treat stroke of the MCA territory, in order to release more stem cells to the infarcted area compared to

Table 1.

Publication	Purpose	Type of Study	Pathology	Number of Patients	Age (years)	Cell Transplant	Time From Onset to Therapy	Follow-Up	Results
Wang et al ¹²⁵	To investigate the effects of transplantation with umbilical cord mesenchymal stem cells in patients with sequelae of traumatic brain injury	Randomized, open-label trial	Traumatic brain injury	40 (20 controls, 20 with hUCT-MSC)	5-57	Four stem cell transplantations (over an interval of 5-7 d) via lumbar puncture (2 mL suspension containing 10 ⁷ cells)	1-11 y	6 mo	<ul style="list-style-type: none"> No adverse effect during transplant No adverse effect during long-term follow-up Significant improvement in neurological function and self-care in hUCT-MSC-treated patients
Jozwiak et al ¹²⁰	To monitor the fate of SPIO-labeled autologous hUCB-derived cells after intracerebral transplantation in a child in a permanent vegetative state resulting from global hypoxic-ischemic brain injury	Pilot study	Global hypoxic/ischemic brain injury	1	1-5	Autologous neutrally committed cord blood cells	7 mo	6 mo	<ul style="list-style-type: none"> No sign of brain edema, hemorrhage, or tumor appearance after transplant Cells are found in the ventricle up to 2 mo posttransplant Neurological function improved
Jiang et al ¹²²	To evaluate the safety and efficacy of hUCT-MSC delivered by a catheter to a near lesion site for treatment of an infarction in the middle cerebral artery territory	Pilot study	Stroke	4 (3 ischemic and 1 hemorrhagic stroke)	40-59	One single dose of 2 × 10 ⁷ hUCT-MSC via catheterization in the M1 segment of the middle cerebral artery	11-50 d	6 mo	<ul style="list-style-type: none"> No major accident observed No fever or rash 2 of the 3 ischemic stroke patients showed improvement in muscle strength and modified Rankin scale
Chen et al ¹²³	To explore the possible role of multiple cells through different implant routes in stroke patients at chronic stage	Pilot study	Stroke	10 (6 ischemic and 4 hemorrhagic stroke)	42-87	Single or multiple doses of olfactory ensheathing cells, neural progenitor cells, umbilical cord mesenchymal cells or Schwann cells, infused intracranially, intrathecally, or intravenously	?	6 mo to 20 y	<ul style="list-style-type: none"> No adverse effect during transplant No adverse effect during long-term follow-up All patients showed neurological improvements
Chang et al ¹²⁴	To examine treatment of cerebral hemorrhages with bone marrow- or umbilical cord-derived mesenchymal stem cells and conventional surgical approaches	Retrospective analysis	Cerebral hemorrhage	24 (8 controls, 7 with BM-MSC, 9 with UC-MSC)	38-55	Transplantation in the hematoma cavity at 2 and 3 wk after hemorrhage	2 wk and 3 wk	5 y	<ul style="list-style-type: none"> Showed a shorter time of hematoma reabsorption Improvement in neurological function

Abbreviations: hUCB, human umbilical cord blood; SPIO, superparamagnetic iron oxide; hUCT-MSCs, human umbilical cord tissue-derived mesenchymal stromal cells; BM-MSCs, bone marrow-derived mesenchymal stromal cells; UC-MSCs, umbilical cord-derived mesenchymal stromal cells.

Table 2.

Registration Number	Trial Name	Purpose	Phase	Start Date	Status	Condition	Study Design	Intervention	Regimen	Sponsor
NCT01310114	Study of human placenta-derived cells (PDA001) to evaluate the safety and effectiveness for patients with ischemic stroke	To assess the safety and tolerability of human placenta-derived cells (PDA001) versus placebo administered IV in subjects following ischemic stroke	I/II	March 2011	Terminated	Stroke	Randomized, double blind placebo controlled	Human placenta-derived cells (PDA001)—Cenplace(L)	<ul style="list-style-type: none"> • 2×10^8 cells or placebo on day 1 • 4 units of 2×10^8 cells or placebo on day 1 	Celgene Corporation Tennessee, United States
NCT01673932	Safety and feasibility study of umbilical cord blood mononuclear cells transplant to treat ischemic stroke	To assess the safety and possible efficacy of umbilical cord blood mononuclear cells (UCBMC) for treatment of chronic ischemic stroke	I	October 2012	Recruiting	Ischemic stroke	Randomized open label	Umbilical cord blood mononuclear cells	<ul style="list-style-type: none"> • $10\text{--}40 \times 10^6$ cells into brain adjacent to infarcted site on day 0 • $10\text{--}40 \times 10^6$ cells into brain adjacent to infarcted site on month 6 	China Spinal Cord Injury Network, Hong Kong
NCT02378974	Evaluation of the safety and potential therapeutic effects after IV transplantation of Cordstem-ST in patients with cerebral infarction	To evaluate the safety and the potential therapeutic effects per dose of Cordstem-ST IV transplantation in cerebral infarction	I/II	February 2015	Active, not recruiting	Cerebral infarction	Randomized, double blind, placebo controlled	Cordstem-ST	<ul style="list-style-type: none"> • 2.0×10^8 cells or placebo on day 0 • 2.0×10^8 cells or placebo on day 0 and day 7 	CHA Biotech Co., Ltd. Korea
NCT02283879	Human umbilical cord mesenchymal stem cell in cerebral hemorrhage sequela	To evaluate the safety and efficacy of human umbilical cord mesenchymal stem cell (hUC-MSC) for cerebral hemorrhage sequela	I	March 2015	Active, not recruiting	Cerebral hemorrhage	Single-group assignment open label	Human umbilical cord mesenchymal stem cells	<ul style="list-style-type: none"> • Single dose of 2×10^7 hUC-MSC IV, repeated every weeks for 4 times 	Shenzhen Hornetcom Biotechnology Company, Ltd. China
NCT02433509	Phase I clinical safety study about human umbilical cord blood monocyte in the acute ischemic stroke	To determine the safety of human umbilical cord blood mononuclear cells by IV injection in acute ischemic stroke patients	I	May 2015	Recruiting	Acute ischemic stroke	Single-group assignment open label	Human umbilical cord blood mononuclear cells and 20% mannitol	<ul style="list-style-type: none"> • $200\text{--}500 \times 10^6$ cells IV infused within 72 h after stroke. 20% mannitol 200 mL IV administered twice after cord blood infusion 	China Medical University Hospital, Taiwan

(continued)

Table 2. (continued)

Registration Number	Trial Name	Purpose	Phase	Start Date	Status	Condition	Study Design	Intervention	Regimen	Sponsor
NCT02397018	Cord blood infusion for ischemic stroke	To investigate the safety of a single IV infusion of banked allogeneic umbilical cord blood in subjects following an acute ischemic stroke	I	May 2015	Active, not recruiting	Stroke	Single-group assignment open label	Allogenic umbilical cord blood cells	Single dose of $0.5-5 \times 10^7$ cells, IV infused, 3-10 d after stroke	Joanne Kurtzberg, MD, Texas (USA)
NCT02580019	Umbilical cord-derived mesenchymal stem cells treatment in ischemic stroke	To estimate the safety and efficacy of the IV injection of human umbilical cord mesenchymal stem cells for patients suffering from ischemic stroke in recent 3 mo	II	February 2016	Active, not recruiting	Stroke	Randomized, open label	Human umbilical cord mesenchymal stem cells	A single dose of 2×10^7 hUC-MSC to patients, IV. Repeat every weeks for 4 times	Affiliated hospital to academy of military medical sciences, China
NCT03004976	Study of allogeneic umbilical cord blood infusion for adults with ischemic stroke (CoBIS 2)	To determine the efficacy of a single IV infusion of unrelated donor cord blood in patients with ischemic stroke	II	January 2017	Active, not recruiting	Stroke	Randomized, double blind, placebo controlled	Umbilical cord blood cells	Single dose of $0.5-5 \times 10^7$ cells, IV infused, 3-10 d after stroke	Joanne Kurtzberg, MD, Texas (USA)

the more commonly used intravenous transplantation¹²². Four males (between 40 and 59 y) with stroke (3 ischemic and 1 hemorrhagic) were treated with 20 million hUCT-MSCs between 11 and 50 d after the stroke onset. Patients were monitored for the presence of additional ischemic strokes: angiography 20 min after cell delivery did not indicate any new blood flow impairment and MRI prior to discharge did not reveal any new infarct. After transplant, there was no sign of immune response (fever or rash), demonstrating the feasibility of intra-arterial delivery of hUCT-MSCs. The trial was not powered to assess hUCT-MSC efficacy and no conclusions could be drawn about the therapeutic potential.

Chen and colleagues employed a multiple cell transplantation approach for stroke patients in the chronic phase¹²³. Ten stroke patients (6 ischemic and 4 hemorrhagic), aged 42 to 87 y, were transplanted with different cell populations: olfactory ensheathing cells from aborted human fetal olfactory bulbs, Schwann cells from aborted human fetal sciatic nerve, neural progenitor cells from aborted human subependymal zone of fetal brain, and/or hUCT-MSC collected after birth. Patients received one or more doses of cells, ranging from 1 to 23 million cells, using different delivery routes (intracranial, intrathecal, or intravascular) and different intervals from stroke onset (from 6 mo to 20 y). These heterogeneous treatments/protocols led to no adverse events during or after the procedure or during the long-term follow-up. The authors state that all patients achieved some degree of improvement in neurological function, speech, muscle tension, balance, pain relief, and respiratory stability. Again, the small sample size limits the value of the findings on combinatory cell therapy, so the results must be interpreted with caution.

A retrospective analysis compared the effect of human bone marrow-derived mesenchymal stromal cells (hBM-MSCs) and hUCT-MSCs or a conventional surgical approach for ICH¹²⁴. A total of 24 patients were treated with conventional hematoma removal surgery within 6 h of the hemorrhage. Patients scheduled for MSC treatment received 2 transplants of hBM-MSCs ($n = 7$) or hUCT-MSCs ($n = 9$) into the hematoma cavity 2 and 3 wk after injury. All the transplanted patients had a shorter hematoma reabsorption time and a better outcome at 5 y than untreated patients. Importantly, patients receiving hUCT-MSCs had a better outcome than hBM-MSC-treated patients starting from 3 mo after injury, suggesting that placenta-derived stem cells have higher therapeutic potential than adult stem cells¹²⁴.

Only one clinical study using placenta-derived stem cells in TBI has been published¹²⁵. Forty patients with TBI at chronic stages (range: 1-11 y post-TBI) were randomly allocated to treatment with hUCT-MSCs or vehicle, and follow-up was obtained at 6 mo posttreatment. Twenty patients in the stem cell group received 4 hUCT-MSC transplants, each containing 10 million stem cells (over an interval of 5-7 d) by lumbar puncture. All patients were analyzed by Fugl-Meyer assessment (FMA)¹²⁶, a multi-item scale assessing

motor function, sensory function, balance, joint range of motion, and joint pain, and by Functional Independence Measure (FIM)¹²⁷, a multi-item rating scale assessing self-care, bowel and bladder management, mobility, communication, cognition, and psychosocial adjustment. During stem cell transplantation, patients were monitored for body temperature, heart rates, blood pressures, oxygen saturations, and respiratory rates, and no obvious abnormalities were found. Four (20%) patients experienced headache and mild dizziness within 48 h post lumbar puncture. At 6 mo, patients received head and spinal cord MRI examinations and no abnormalities related to the stem cell transplantation were found. Rating scales at 6 mo indicated that while the control group had FMA and FIM scores not significantly different from the baseline time point, the hUCT-MSC-treated patients had slightly better FMA and FIM scores¹²⁵. Thus, the preliminary findings of the therapeutic potential of hUCT-MSCs demonstrate the feasibility and safety of this approach for acute brain injury. Further research is now needed to validate and strengthen these results in order to offer cell therapy for patients with acute brain injury.

Currently, 8 ongoing phase I or II clinical trials are present in the world's largest registry clinicaltrials.gov using placenta-derived stem cells for acute brain injuries (Table 2). Seven trials target stroke, 1 cerebral hemorrhage, and none have been designed for TBI. All trials are single center and use UC-derived stem cells. Three trials are designed as single-group assignment open label, 2 as randomized open label, and 3 as randomized double-blind placebo-controlled trials. Thus, several conclusions will be drawn at the end of these trials, posing the bases for the construction of a larger phase 3 trial.

Conclusions

There is growing experimental evidence for the efficacy of placenta-derived stem cells for acute brain injury. The multi-target potential of cell therapy fulfills the need of the damaged brain in which numerous injury mechanisms are triggered. Infused cells can reprogram the local inflammatory microenvironment from detrimental to beneficial, favoring protective/pro-regenerative changes in the lesioned tissue and contributing to permanent improvement in neurological function. Even in case of predifferentiation into neuronal lineage, cell replacement is unlikely, whereas the main mechanisms of action through which infused cells confer protection is through the secretion of bioactive factors, suggesting the potential for a cell-free strategy based on the cell secretome. Different classes of molecules have been identified as mediators of protective effects. However, the pool of factors sufficient and necessary to promote brain protection/repair has not yet been pinpointed and further studies are needed to identify a controlled, standardized "secretome-based" cell-free strategy. Clinical studies are in their infancy, with few reports demonstrating the safety and preliminary efficacy of cell transplantation. The ongoing trials are crucial

to assess the efficacy of placenta-derived stem cells for acute brain injury.

Authors' Note

Francesca Pischiutta received a fellowship from “Fondazione Umberto Veronesi” (FUV, 2016).

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by the association “Amici del Mario Negri.”

References

- Stocchetti N, Taccone FS, Citerio G, Pepe PE, Le Roux PD, Oddo M, Polderman KH, Stevens RD, Barsan W, Maas AIR, et al. Neuroprotection in acute brain injury: an up-to-date review. *Crit Care*. 2015;19(1): 186.
- Harting MT, Baumgartner JE, Worth LL, Ewing-Cobbs L, Gee AP, Day MC, Cox CS Jr. Cell therapies for traumatic brain injury. *Neurosurg Focus*. 2008;24(3–4): E18.
- Zhang J, Chopp M. Cell-based therapy for ischemic stroke. *Expert Opin Biol Ther*. 2013;13(9):1229–1240.
- Eckert MA, Vu Q, Xie K, Yu J, Liao W, Cramer SC, Zhao W. Evidence for high translational potential of mesenchymal stromal cell therapy to improve recovery from ischemic stroke. *J Cereb Blood Flow Metab*. 2013;33(9):1322–1334.
- Gennai S, Monsel A, Hao Q, Liu J, Gudapati V, Barbier EL, Lee JW. Cell-based therapy for traumatic brain injury. *Br J Anaesth*. 2015;115(2):203–212.
- Moisan A, Favre I, Rome C, De Fraipont F, Grillon E, Coquery N, Mathieu H, Mayan V, Naegele B, Hommel M, et al. Intravenous injection of clinical grade human MSCs after experimental stroke: functional benefit and microvascular effect. *Cell Transplant*. 2016;25(12):2157–2171.
- Du WJ, Chi Y, Yang ZX, Li ZJ, Cui JJ, Song BQ, Li X, Yang SG, Han ZB, Han ZC. Heterogeneity of proangiogenic features in mesenchymal stem cells derived from bone marrow, adipose tissue, umbilical cord, and placenta. *Stem Cell Res Ther*. 2016; 7(1):163.
- Heo JS, Choi Y, Kim HS, Kim HO. Comparison of molecular profiles of human mesenchymal stem cells derived from bone marrow, umbilical cord blood, placenta and adipose tissue. *Int J Mol Med*. 2016;37(1):115–125.
- Efimenko AY, Kochegura TN, Akopyan ZA, Parfyonova YV. Autologous stem cell therapy: how aging and chronic diseases affect stem and progenitor cells. *Biores Open Access*. 2015; 4(1):26–38.
- Zhou S, Greenberger JS, Epperly MW, Goff JP, Adler C, Leboff MS, Glowacki J. Age-related intrinsic changes in human bone-marrow-derived mesenchymal stem cells and their differentiation to osteoblasts. *Aging Cell*. 2008;7(3):335–343.
- Stolzing A, Jones E, McGonagle D, Scutt A. Age-related changes in human bone marrow-derived mesenchymal stem cells: consequences for cell therapies. *Mech Ageing Dev*. 2008;129(3):163–173.
- Siegel G, Kluba T, Hermanutz-Klein U, Bieback K, Northoff H, Schäfer R. Phenotype, donor age and gender affect function of human bone marrow-derived mesenchymal stromal cells. *BMC Med*. 2013;11:146.
- Kizilay Mancini Ö, Lora M, Shum-Tim D, Nadeau S, Rodier F, Colmegna I. A proinflammatory secretome mediates the impaired immunopotency of human mesenchymal stromal cells in elderly patients with atherosclerosis. *Stem Cells Transl Med*. 2017;6(4):1132–1140.
- Fan M, Chen W, Liu W, Du GQ, Jiang SL, Tian WC, Sun L, Li RK, Tian H. The effect of age on the efficacy of human mesenchymal stem cell transplantation after a myocardial infarction. *Rejuvenation Res*. 2010;13(4):429–438.
- Efimenko A, Dzhoyashvili N, Kalinina N, Kochegura T, Akchurin R, Tkachuk V, Parfyonova Y. Adipose-derived mesenchymal stromal cells from aged patients with coronary artery disease keep mesenchymal stromal cell properties but exhibit characteristics of aging and have impaired angiogenic potential. *Stem Cells Transl Med*. 2014;3(1):32–41.
- Parolini O, Alviano F, Bagnara GP, Bilic G, Bühring HJ, Evangelista M, Hennerbichler S, Liu B, Magatti M, Mao N, et al. Concise review: isolation and characterization of cells from human term placenta: outcome of the first international workshop on placenta derived stem cells. *Stem Cells*. 2008;26(2): 300–311.
- Malek A, Bersinger NA. Human placental stem cells: biomedical potential and clinical relevance. *J Stem Cells*. 2011;6(2): 75–92.
- Caruso M, Evangelista M, Parolini O. Human term placental cells: phenotype, properties and new avenues in regenerative medicine. *Int J Mol Cell Med*. 2012;1(2):64–74.
- Lu D, Sanberg PR, Mahmood A, Li Y, Wang L, Sanchez-Ramos J, Chopp M. Intravenous administration of human umbilical cord blood reduces neurological deficit in the rat after traumatic brain injury. *Cell Transplant*. 2002;11(3):275–281.
- Zanier ER, Montinaro M, Viganò M, Villa P, Fumagalli S, Pischiutta F, Longhi L, Leoni ML, Rebullà P, Stocchetti N, et al. Human umbilical cord blood mesenchymal stem cells protect mice brain after trauma. *Crit Care Med*. 2011;39(11): 2501–2510.
- Chen SH, Wang JJ, Chen CH, Chang HK, Lin MT, Chang FM, Chio CC. Umbilical cord blood-derived CD34⁺ cells improve outcomes of traumatic brain injury in rats by stimulating angiogenesis and neurogenesis. *Cell Transplant*. 2014;23(8): 959–979.
- Zhang Y, Li Y, Wang S, Han Z, Huang X, Li S, Chen F, Niu R, Dong J, Jiang R, et al. Transplantation of expanded endothelial colony-forming cells improved outcomes of traumatic brain injury in a mouse model. *J Surg Res*. 2013;185(1):441–449.
- Huang XT, Zhang YQ, Li SJ, Li SH, Tang Q, Wang ZT, Dong JF, Zhang JN. Intracerebroventricular transplantation of ex vivo expanded endothelial colony-forming cells restores

- blood-brain barrier integrity and promotes angiogenesis of mice with traumatic brain injury. *J Neurotrauma*. 2013;30(24):2080–2088.
24. Ariens-Zakay H, Gincberg G, Nagler A, Cohen G, Liraz-Zaltsman S, Trembovler V, Alexandrovich AG, Matok I, Galski H, Elchalal U, et al. Neurotherapeutic effect of cord blood derived CD45+ hematopoietic cells in mice after traumatic brain injury. *J Neurotrauma*. 2014;31(16):1405–1416.
 25. Ma J, Liu N, Yi B, Zhang X, Gao BB, Zhang Y, Xu R, Li X, Dai Y. Transplanted hUCB-MSCs migrated to the damaged area by SDF-1/CXCR4 signaling to promote functional recovery after traumatic brain injury in rats. *Neurol Res*. 2015;37(1):50–56.
 26. Cheng T, Yang B, Li D, Ma S, Tian Y, Qu R, Zhang W, Zhang Y, Hu K, Guan F, et al. Wharton's jelly transplantation improves neurologic function in a rat model of traumatic brain injury. *Cell Mol Neurobiol*. 2015;35(5):641–649.
 27. Zhou HX, Liu ZG, Liu XJ, Chen QX. Umbilical cord-derived mesenchymal stem cell transplantation combined with hyperbaric oxygen treatment for repair of traumatic brain injury. *Neural Regen Res*. 2016;11(1):107–113.
 28. Chen Z, Tortella FC, Dave JR, Marshall VS, Clarke DL, Sing G, Du F, Lu XCM. Human amnion-derived multipotent progenitor cell treatment alleviates traumatic brain injury-induced axonal degeneration. *J Neurotrauma*. 2009;26(11):1987–1997.
 29. Pischiutta F, Brunelli L, Romele P, Silini A, Sammali E, Paracchini L, Marchini S, Talamini L, Bigini P, Boncoraglio GB, et al. Protection of brain injury by amniotic mesenchymal stromal cell-secreted metabolites. *Crit Care Med*. 2016;44(11):e1118–e1131.
 30. Yan H, Zhang HW, Wu QL, Zhang GB, Liu K, Zhi DS, Hu ZB, Zeng XW. Increased leakage of brain antigens after traumatic brain injury and effect of immune tolerance induced by cells on traumatic brain injury. *Chin Med J (Engl)*. 2012;125(9):1618–1626.
 31. Chen J, Sanberg PR, Li Y, Wang L, Lu M, Willing AE, Sanchez-Ramos J, Chopp M. Intravenous administration of human umbilical cord blood reduces behavioral deficits after stroke in rats. *Stroke*. 2001;32(11):2682–2688.
 32. Okawa H, Okuda O, Arai H, Sakuragawa N, Sato K. Amniotic epithelial cells transform into neuron-like cells in the ischemic brain. *Neuroreport*. 2001;12(18):4003–4007.
 33. Willing AE, Lixian J, Milliken M, Poulos S, Zigova T, Song S, Hart C, Sanchez-Ramos J, Sanberg PR. Intravenous versus intrastriatal cord blood administration in a rodent model of stroke. *J Neurosci Res*. 2003;73(3):296–307.
 34. Zhang L, Li Y, Zhang C, Chopp M, Gosiewska A, Hong K. Delayed administration of human umbilical tissue-derived cells improved neurological functional recovery in a rodent model of focal ischemia. *Stroke*. 2011;42(5):1437–1444.
 35. Fu YS, Shih YT, Cheng YC, Min MY. Transformation of human umbilical mesenchymal cells into neurons in vitro. *J Biomed Sci*. 2004;11(5):652–660.
 36. Koh SH, Kim KS, Choi MR, Jung KH, Park KS, Chai YG, Roh W, Hwang SJ, Ko HJ, Huh YM, et al. Implantation of human umbilical cord-derived mesenchymal stem cells as a neuroprotective therapy for ischemic stroke in rats. *Brain Res*. 2008;1229:233–248.
 37. Lin YC, Ko TL, Shih YH, Lin MYA, Fu TW, Hsiao HS, Hsu JYC, Fu YS. Human umbilical mesenchymal stem cells promote recovery after ischemic stroke. *Stroke*. 2011;42(7):2045–2053.
 38. Newman MB, Willing AE, Manresa JJ, Davis-Sanberg C, Sanberg PR. Stroke-induced migration of human umbilical cord blood cells: time course and cytokines. *Stem Cells Dev*. 2005;14(5):576–586.
 39. Rosenkranz K, Kumbruch S, Lebermann K, Marschner K, Jensen A, Dermietzel R, Meier C. The chemokine SDF-1/CXCL12 contributes to the “homing” of umbilical cord blood cells to a hypoxic-ischemic lesion in the rat brain. *J Neurosci Res*. 2010;88(6):1223–1233.
 40. Borlongan CV, Hadman M, Sanberg CD, Sanberg PR. Central nervous system entry of peripherally injected umbilical cord blood cells is not required for neuroprotection in stroke. *Stroke*. 2004;35(10):2385–2389.
 41. Xiao J, Nan Z, Motooka Y, Low WC. Transplantation of a novel cell line population of umbilical cord blood stem cells ameliorates neurological deficits associated with ischemic brain injury. *Stem Cells Dev*. 2005;14(6):722–733.
 42. Taguchi A, Soma T, Tanaka H, Kanda T, Nishimura H, Yoshikawa H, Tsukamoto Y, Iso H, Fujimori Y, Stern DM, et al. Administration of CD34+ cells after stroke enhances neurogenesis via angiogenesis in a mouse model. *J Clin Invest*. 2004;114(3):330–338.
 43. Rosenkranz K, Kumbruch S, Tenbusch M, Marcus K, Marschner K, Dermietzel R, Meier C. Transplantation of human umbilical cord blood cells mediated beneficial effects on apoptosis, angiogenesis and neuronal survival after hypoxic-ischemic brain injury in rats. *Cell Tissue Res*. 2012;348(3):429–438.
 44. Vendrame M, Gemma C, Pennypacker KR, Bickford PC, Davis Sanberg C, Sanberg PR, Willing AE. Cord blood rescues stroke-induced changes in splenocyte phenotype and function. *Exp Neurol*. 2006;199(1):191–200.
 45. Newcomb JD, Ajmo CT, Sanberg CD, Sanberg PR, Pennypacker KR, Willing AE. Timing of cord blood treatment after experimental stroke determines therapeutic efficacy. *Cell Transplant*. 2006;15(3):213–223.
 46. Vendrame M, Gemma C, de Mesquita D, Collier L, Bickford PC, Sanberg CD, Sanberg PR, Pennypacker KR, Willing AE. Anti-inflammatory effects of human cord blood cells in a rat model of stroke. *Stem Cells Dev*. 2005;14(5):595–604.
 47. Hocum Stone LL, Xiao F, Rotschafer J, Nan Z, Juliano M, Sanberg CD, Sanberg PR, Kuzmin-Nichols N, Grande A, Cheeran MCJ, et al. Amelioration of ischemic brain injury in rats with human umbilical cord blood stem cells: mechanisms of action. *Cell Transplant*. 2016;25(8):1473–1488.
 48. Wasielewski B, Jensen A, Roth-Härer A, Dermietzel R, Meier C. Neuroglial activation and Cx43 expression are reduced upon transplantation of human umbilical cord blood cells after perinatal hypoxic-ischemic injury. *Brain Res*. 2012;1487:39–53.
 49. Tsuji M, Taguchi A, Ohshima M, Kasahara Y, Sato Y, Tsuda H, Otani K, Yamahara K, Ihara M, Harada-Shiba M, et al. Effects of intravenous administration of umbilical cord blood

- CD34(+) cells in a mouse model of neonatal stroke. *Neuroscience*. 2014;263:148–158.
50. Boltze J, Kowalski I, Geiger K, Reich D, Gunther A, Buhrl C, Egger D, Kamprad M, Emmrich F. Experimental treatment of stroke in spontaneously hypertensive rats by CD34+ and CD34- cord blood cells. *Ger Med Sci*. 2005;3: Doc09.
 51. Nystedt J, Mäkinen S, Laine J, Jolkkonen J. Human cord blood CD34+ cells and behavioral recovery following focal cerebral ischemia in rats. *Acta Neurobiol Exp (Wars)*. 2006;66(4): 293–300.
 52. Ou Y, Yu S, Kaneko Y, Tajiri N, Bae EC, Chheda SH, Stahl CE, Yang T, Fang L, Hu K, et al. Intravenous infusion of GDNF gene-modified human umbilical cord blood CD34+ cells protects against cerebral ischemic injury in spontaneously hypertensive rats. *Brain Res*. 2010;1366:217–225.
 53. Boltze J, Reich DM, Hau S, Reymann KG, Strassburger M, Lobsien D, Wagner DC, Kamprad M, Stahl T. Assessment of neuroprotective effects of human umbilical cord blood mononuclear cell subpopulations in vitro and in vivo. *Cell Transplant*. 2012;21(4):723–737.
 54. Womble TA, Green S, Shahaduzzaman M, Grieco J, Sanberg PR, Pennypacker KR, Willing AE. Monocytes are essential for the neuroprotective effect of human cord blood cells following middle cerebral artery occlusion in rat. *Mol Cell Neurosci*. 2014;59:76–84.
 55. Lim JY, Jeong CH, Jun JA, Kim SM, Ryu CH, Hou Y, Oh W, Chang JW, Jeun SS. Therapeutic effects of human umbilical cord blood-derived mesenchymal stem cells after intrathecal administration by lumbar puncture in a rat model of cerebral ischemia. *Stem Cell Res Ther*. 2011;2(5):38.
 56. Chung DJ, Choi CB, Lee SH, Kang EH, Lee JH, Hwang SH, Han H, Lee JH, Choe BY, Lee SY, et al. Intraarterially delivered human umbilical cord blood-derived mesenchymal stem cells in canine cerebral ischemia. *J Neurosci Res*. 2009;87(16): 3554–3567.
 57. Xia G, Hong X, Chen X, Lan F, Zhang G, Liao L. Intracerebral transplantation of mesenchymal stem cells derived from human umbilical cord blood alleviates hypoxic ischemic brain injury in rat neonates. *J Perinat Med*. 2010;38(2):215–221.
 58. Kim ES, Ahn SY, Im GH, Sung DK, Park YR, Choi SH, Choi SJ, Chang YS, Oh W, Lee JH, et al. Human umbilical cord blood-derived mesenchymal stem cell transplantation attenuates severe brain injury by permanent middle cerebral artery occlusion in newborn rats. *Pediatr Res*. 2012;72(3):277–284.
 59. Zhu Y, Guan Y, Huang H, Wang Q. Human umbilical cord blood mesenchymal stem cell transplantation suppresses inflammatory responses and neuronal apoptosis during early stage of focal cerebral ischemia in rabbits. *Acta Pharmacol Sin*. 2014;35(5):585–591.
 60. Cheng Q, Zhang Z, Zhang S, Yang H, Zhang X, Pan J, Weng L, Sha D, Zhu M, Hu X, et al. Human umbilical cord mesenchymal stem cells protect against ischemic brain injury in mouse by regulating peripheral immunoinflammation. *Brain Res*. 2015;1594:293–304.
 61. Mäkinen S, Kekarainen T, Nystedt J, Liimatainen T, Huhtala T, Närvänen A, Laine J, Jolkkonen J. Human umbilical cord blood cells do not improve sensorimotor or cognitive outcome following transient middle cerebral artery occlusion in rats. *Brain Res*. 2006;1123(1):207–215.
 62. Riegelsberger UM, Deten A, Pösel C, Zille M, Kranz A, Boltze J, Wagner DC. Intravenous human umbilical cord blood transplantation for stroke: impact on infarct volume and caspase-3-dependent cell death in spontaneously hypertensive rats. *Exp Neurol*. 2011;227(1):218–223.
 63. Weise G, Lorenz M, Pösel C, Maria Riegelsberger U, Störbeck V, Kamprad M, Kranz A, Wagner DC, Boltze J. Transplantation of cryopreserved human umbilical cord blood mononuclear cells does not induce sustained recovery after experimental stroke in spontaneously hypertensive rats. *J Cereb Blood Flow Metab*. 2014;34(1): e1–e9.
 64. Liu T, Wu J, Huang Q, Hou Y, Jiang Z, Zang S, Guo L. Human amniotic epithelial cells ameliorate behavioral dysfunction and reduce infarct size in the rat middle cerebral artery occlusion model. *Shock*. 2008;29(5):603–611.
 65. Li F, Miao ZN, Xu YY, Zheng SY, Qin MD, Gu YZ, Zhang XG. Transplantation of human amniotic mesenchymal stem cells in the treatment of focal cerebral ischemia. *Mol Med Rep*. 2012;6(3):625–630.
 66. Bigini P, Zanier ER, Saragozza S, Maciotta S, Romele P, Bonassi Signoroni P, Silini A, Pischiutta F, Sammali E, Balducci C, et al. Internalization of nanopolymeric tracers does not alter characteristics of placental cells. *J Cell Mol Med*. 2016;20(6):1036–1048.
 67. Tao J, Ji F, Liu B, Wang F, Dong F, Zhu Y. Improvement of deficits by transplantation of lentiviral vector-modified human amniotic mesenchymal cells after cerebral ischemia in rats. *Brain Res*. 2012;1448:1–10.
 68. Rehni AK, Singh N, Jaggi AS, Singh M. Amniotic fluid derived stem cells ameliorate focal cerebral ischaemia-reperfusion injury induced behavioural deficits in mice. *Behav Brain Res*. 2007;183(1):95–100.
 69. Tajiri N, Acosta S, Glover LE, Bickford PC, Jacotte Simancas A, Yasuhara T, Date I, Solomita MA, Antonucci I, Stuppia L, et al. Intravenous grafts of amniotic fluid-derived stem cells induce endogenous cell proliferation and attenuate behavioral deficits in ischemic stroke rats. *PLoS One*. 2012;7(8):e43779.
 70. Qin M, Chen R, Li H, Liang H, Xue Q, Li F, Chen Y, Zhang X. Direct reprogramming of human amniotic fluid stem cells by OCT4 and application in repairing of cerebral ischemia damage. *Int J Biol Sci*. 2016;12(5):558–568.
 71. Seghatoleslam M, Jalali M, Nikravesh MR, Hamidi Alamdari D, Hosseini M, Fazel A. Intravenous administration of human umbilical cord blood-mononuclear cells dose-dependently relieve neurologic deficits in rat intracerebral hemorrhage model. *Ann Anat*. 2013;195(1):39–49.
 72. Kim K, Park HW, Moon HE, Kim JW, Bae S, Chang JW, Oh W, Yang YS, Paek SH. The effect of human umbilical cord blood-derived mesenchymal stem cells in a collagenase-induced intracerebral hemorrhage rat model. *Exp Neurobiol*. 2015;24(2):146–155.
 73. Xie J, Wang B, Wang L, Dong F, Bai G, Liu Y. Intracerebral and intravenous transplantation represents a favorable

- approach for application of human umbilical cord mesenchymal stromal cells in intracerebral hemorrhage rats. *Med Sci Monit.* 2016;22:3552–3561.
74. Hong SQ, Zhang HT, You J, Zhang MY, Cai YQ, Jiang XD, Xu RX. Comparison of transdifferentiated and undifferentiated human umbilical mesenchymal stem cells in rats after traumatic brain injury. *Neurochem Res.* 2011;36(12):2391–2400.
75. Yan ZJ, Zhang P, Hu YQ, Zhang HT, Hong SQ, Zhou HL, Zhang MY, Xu RX. Neural stem-like cells derived from human amnion tissue are effective in treating traumatic brain injury in rat. *Neurochem Res.* 2013;38(5):1022–1033.
76. Kozłowska H, Jabłonka J, Janowski M, Jurga M, Kossut M, Domańska-Janik K. Transplantation of a novel human cord blood-derived neural-like stem cell line in a rat model of cortical infarct. *Stem Cells Dev.* 2007;16(3):481–488.
77. Marion D, Bullock MR. Current and future role of therapeutic hypothermia. *J Neurotrauma.* 2009;26(3):455–467.
78. Tu Y, Chen C, Sun HT, Cheng SX, Liu XZ, Qu Y, Li X, Zhang S. Combination of temperature-sensitive stem cells and mild hypothermia: a new potential therapy for severe traumatic brain injury. *J Neurotrauma.* 2012;29(14):2393–2403.
79. Huang C, Zhao L, Gu J, Nie D, Chen Y, Zuo H, Huan W, Shi J, Chen J, Shi W. The migration and differentiation of hUC-MSCs(CXCR4/GFP) encapsulated in BDNF/chitosan scaffolds for brain tissue engineering. *Biomed Mater.* 2016;11(3):035004.
80. Shi W, Huang CJ, Xu XD, Jin GH, Huang RQ, Huang JF, Chen YN, Ju SQ, Wang Y, Shi YW, et al. Transplantation of RADA16-BDNF peptide scaffold with human umbilical cord mesenchymal stem cells forced with CXCR4 and activated astrocytes for repair of traumatic brain injury. *Acta Biomater.* 2016;45:247–261.
81. Liu AM, Lu G, Tsang KS, Li G, Wu Y, Huang ZS, Ng HK, Kung HF, Poon WS. Umbilical cord-derived mesenchymal stem cells with forced expression of hepatocyte growth factor enhance remyelination and functional recovery in a rat intracerebral hemorrhage model. *Neurosurgery.* 2010;67(2):357–365; discussion 365–366.
82. Uchida S, Inanaga Y, Kobayashi M, Hurukawa S, Araie M, Sakuragawa N. Neurotrophic function of conditioned medium from human amniotic epithelial cells. *J Neurosci Res.* 2000;62(4):585–590.
83. Salgado AJ, Fraga JS, Mesquita AR, Neves NM, Reis RL, Sousa N. Role of human umbilical cord mesenchymal progenitors conditioned media in neuronal/glial cell densities, viability, and proliferation. *Stem Cells Dev.* 2010;19(7):1067–1074.
84. Ribeiro CA, Fraga JS, Grãos M, Neves NM, Reis RL, Gimble JM, Sousa N, Salgado AJ. The secretome of stem cells isolated from the adipose tissue and Wharton jelly acts differently on central nervous system derived cell populations. *Stem Cell Res Ther.* 2012;3(3):18.
85. Fraga JS, Silva NA, Lourenço AS, Gonçalves V, Neves NM, Reis RL, Rodrigues AJ, Manadas B, Sousa N, Salgado AJ. Unveiling the effects of the secretome of mesenchymal progenitors from the umbilical cord in different neuronal cell populations. *Biochimie.* 2013;95(12):2297–2303.
86. Teixeira FG, Carvalho MM, Neves-Carvalho A, Panchalingam KM, Behie LA, Pinto L, Sousa N, Salgado AJ. Secretome of mesenchymal progenitors from the umbilical cord acts as modulator of neural/glial proliferation and differentiation. *Stem Cell Rev.* 2015;11(2):288–297.
87. Deng-Bryant Y, Chen Z, van der Merwe C, Liao Z, Dave JR, Rupp R, Shear DA, Tortella FC. Long-term administration of amnion-derived cellular cytokine suspension promotes functional recovery in a model of penetrating ballistic-like brain injury. *J Trauma Acute Care Surg.* 2012;73(2 suppl 1):S156–S164.
88. Deng-Bryant Y, Readnow RD, Leung LY, Cunningham TL, Shear DA, Tortella FC. Treatment with amnion-derived cellular cytokine solution (ACCS) induces persistent motor improvement and ameliorates neuroinflammation in a rat model of penetrating ballistic-like brain injury. *Restor Neurol Neurosci.* 2015;33(2):189–203.
89. Bai L, Li D, Li J, Luo Z, Yu S, Cao S, Shen L, Zuo Z, Ma X. Bioactive molecules derived from umbilical cord mesenchymal stem cells. *Acta Histochem.* 2016;118(8):761–769.
90. Pires AO, Mendes-Pinheiro B, Teixeira FG, Anjo SI, Ribeiro-Samy S, Gomes ED, Serra SC, Silva NA, Manadas B, Sousa N, et al. Unveiling the differences of secretome of human bone marrow mesenchymal stem cells, adipose tissue-derived stem cells, and human umbilical cord perivascular cells: a proteomic analysis. *Stem Cells Dev.* 2016;25(14):1073–1083.
91. Yu X, Odenthal M, Fries JWU. Exosomes as miRNA carriers: formation-function-future. *Int J Mol Sci.* 2016;17(12). pii: E2028.
92. Ti D, Hao H, Fu X, Han W. Mesenchymal stem cells-derived exosomal microRNAs contribute to wound inflammation. *Sci China Life Sci.* 2016;59(12):1305–1312.
93. Etzrodt M, Cortez-Retamozo V, Newton A, Zhao J, Ng A, Wildgruber M, Romero P, Wurdinger T, Xavier R, Geissmann F, et al. Regulation of monocyte functional heterogeneity by miR-146a and Relb. *Cell Rep.* 2012;1(4):317–324.
94. Das A, Ganesh K, Khanna S, Sen CK, Roy S. Engulfment of apoptotic cells by macrophages: a role of microRNA-21 in the resolution of wound inflammation. *J Immunol.* 2014;192(3):1120–1129.
95. Dan C, Jinjun B, Zi-Chun H, Lin M, Wei C, Xu Z, Ri Z, Shun C, Wen-Zhu S, Qing-Cai J, et al. Modulation of TNF- α mRNA stability by human antigen R and miR181s in sepsis-induced immunoparalysis. *EMBO Mol Med.* 2014;7(2):140–157.
96. Buller B, Liu X, Wang X, Zhang RL, Zhang L, Hozeska-Solgot A, Chopp M, Zhang ZG. MicroRNA-21 protects neurons from ischemic death. *FEBS J.* 2010;277(20):4299–4307.
97. Zhang L, Dong LY, Li YJ, Hong Z, Wei WS. miR-21 represses FasL in microglia and protects against microglia-mediated neuronal cell death following hypoxia/ischemia. *Glia.* 2012;60(12):1888–1895.
98. Hutchison ER, Kawamoto EM, Taub DD, Lal A, Abdelmohsen K, Zhang Y, Wood WH, Lehrmann E, Camandola S, Becker KG, et al. Evidence for miR-181 involvement in neuroinflammatory responses of astrocytes. *Glia.* 2013;61(7):1018–1028.

99. Zhang L, Dong LY, Li YJ, Hong Z, Wei WS. The microRNA miR-181c controls microglia-mediated neuronal apoptosis by suppressing tumor necrosis factor. *J Neuroinflammation*. 2012;9:211.
100. Chen J, Ning R, Zacharek A, Cui C, Cui X, Yan T, Venkat P, Zhang Y, Chopp M. MiR-126 contributes to human umbilical cord blood cell-induced neurorestorative effects after stroke in type-2 diabetic mice. *Stem Cells*. 2016; 34(1):102–113.
101. Wang S, Aurora AB, Johnson BA, Qi X, McAnally J, Hill JA, Richardson JA, Bassel-Duby R, Olson EN. The endothelial-specific microRNA miR-126 governs vascular integrity and angiogenesis. *Dev Cell*. 2008;15(2):261–271.
102. Wishart DS, Jewison T, Guo AC, Wilson M, Knox C, Liu Y, Djoumbou Y, Mandal R, Aziat F, Dong E, et al. HMDB 3.0—the human metabolome database in 2013. *Nucleic Acids Res*. 2013;41(Database issue):D801–D807.
103. Løvaas E. Antioxidative and metal-chelating effects of polyamines. *Adv Pharmacol*. 1997;38:119–149.
104. Menzie J, Prentice H, Wu JY. Neuroprotective mechanisms of taurine against ischemic stroke. *Brain Sci*. 2013;3(2): 877–907.
105. Kondoh T, Kameishi M, Mallick HN, Ono T, Torii K. Lysine and arginine reduce the effects of cerebral ischemic insults and inhibit glutamate-induced neuronal activity in rats. *Front Integr Neurosci*. 2010;4:18.
106. Vécsei L, Szalárdy L, Fülöp F, Toldi J. Kynurenines in the CNS: recent advances and new questions. *Nat Rev Drug Discov*. 2013;12(1):64–82.
107. Yan EB, Frugier T, Lim CK, Heng B, Sundaram G, Tan M, Rosenfeld JV, Walker DW, Guillemin GJ, Morganti-Kossmann MC. Activation of the kynurenine pathway and increased production of the excitotoxin quinolinic acid following traumatic brain injury in humans. *J Neuroinflammation*. 2015;12:110.
108. Satoh K, Yasumizu T, Kawai Y, Ozaki A, Wu T, Kinoshita K, Sakamoto S. In vitro production of prostaglandins E, F, and 6-keto prostaglandin F1 alpha by human pregnant uterus, decidua and amnion. *Prostaglandins Med*. 1981; 6(4):359–368.
109. Rossi D, Pianta S, Magatti M, Sedlmayr P, Parolini O. Characterization of the conditioned medium from amniotic membrane cells: prostaglandins as key effectors of its immunomodulatory activity. *PLoS One*. 2012;7(10): e46956.
110. Lee BC, Kim HS, Shin TH, Kang I, Lee JY, Kim JJ, Kang HK, Seo Y, Lee S, Yu KR, et al. PGE2 maintains self-renewal of human adult stem cells via EP2-mediated autocrine signaling and its production is regulated by cell-to-cell contact. *Sci Rep*. 2016;6:26298.
111. Najar M, Raicevic G, Boufker HI, Fayyad Kazan H, De Bruyn C, Meuleman N, Bron D, Toungouz M, Lagneaux L. Mesenchymal stromal cells use PGE2 to modulate activation and proliferation of lymphocyte subsets: combined comparison of adipose tissue, Wharton's jelly and bone marrow sources. *Cell Immunol*. 2010;264(2):171–179.
112. Kang JW, Koo HC, Hwang SY, Kang SK, Ra JC, Lee MH, Park YH. Immunomodulatory effects of human amniotic membrane-derived mesenchymal stem cells. *J Vet Sci*. 2012;13(1):23–31.
113. Rozenberg A, Rezk A, Boivin MN, Darlington PJ, Nyirenda M, Li R, Jalili F, Winer R, Artsy EA, Uccelli A, et al. Human mesenchymal stem cells impact Th17 and Th1 responses through a prostaglandin E2 and myeloid-dependent mechanism. *Stem Cells Transl Med*. 2016; 5(11):1506–1514.
114. Maggini J, Mirkin G, Bognanni I, Holmberg J, Piazzón IM, Nepomnaschy I, Costa H, Cañones C, Raiden S, Vermeulen M, et al. Mouse bone marrow-derived mesenchymal stromal cells turn activated macrophages into a regulatory-like profile. *PLoS One*. 2010;5(2): e9252.
115. Spaggiari GM, Abdelrazik H, Becchetti F, Moretta L. MSCs inhibit monocyte-derived DC maturation and function by selectively interfering with the generation of immature DCs: central role of MSC-derived prostaglandin E2. *Blood*. 2009; 113(26):6576–6583.
116. Cazeville C, Muller A, Meynier F, Dutrait N, Bonne C. Protection by prostaglandins from glutamate toxicity in cortical neurons. *Neurochem Int*. 1994;24(4):395–398.
117. Foraker JE, Oh JY, Ylostalo JH, Lee RH, Watanabe J, Prockop DJ. Cross-talk between human mesenchymal stem/progenitor cells (MSCs) and rat hippocampal slices in LPS-stimulated cocultures: the MSCs are activated to secrete prostaglandin E2. *J Neurochem*. 2011;119(5): 1052–1063.
118. Leclerc JL, Ahmad AS, Singh N, Soshnik-Schierling L, Greene E, Dang A, Doré S. Intracerebral hemorrhage outcomes following selective blockade or stimulation of the PGE2 EP1 receptor. *BMC Neurosci*. 2015;16:48.
119. Mohan S, Narumiya S, Doré S. Neuroprotective role of prostaglandin PGE2 EP2 receptor in hemin-mediated toxicity. *Neurotoxicology*. 2015;46:53–59.
120. Jozwiak S, Habich A, Kotulska K, Sarnowska A, Kropiwnicki T, Janowski M, Jurkiewicz E, Lukomska B, Kmiec T, Walecki J, et al. Intracerebroventricular transplantation of cord blood-derived neural progenitors in a child with severe global brain ischemic injury. *Cell Med*. 2010;1(2): 71–80.
121. Janowski M, Walczak P, Kropiwnicki T, Jurkiewicz E, Domanska-Janik K, Bulte JWM, Lukomska B, Roszkowski M. Long-term MRI cell tracking after intraventricular delivery in a patient with global cerebral ischemia and prospects for magnetic navigation of stem cells within the CSF. *PLoS One*. 2014;9(2): e97631.
122. Jiang Y, Zhu W, Zhu J, Wu L, Xu G, Liu X. Feasibility of delivering mesenchymal stem cells via catheter to the proximal end of the lesion artery in patients with stroke in the territory of the middle cerebral artery. *Cell Transplant*. 2013;22(12):2291–2298.
123. Chen L, Xi H, Huang H, Zhang F, Liu Y, Chen D, Xiao J. Multiple cell transplantation based on an intraparenchymal

- approach for patients with chronic phase stroke. *Cell Transplant*. 2013;22(suppl 1):S83–S91.
124. Chang Z, Mao G, Sun L, Ao Q, Gu Y, Liu Y. Cell therapy for cerebral hemorrhage: five year follow-up report. *Exp Ther Med*. 2016;12(6):3535–3540.
125. Wang S, Cheng H, Dai G, Wang X, Hua R, Liu X, Wang P, Chen G, Yue W, An Y. Umbilical cord mesenchymal stem cell transplantation significantly improves neurological function in patients with sequelae of traumatic brain injury. *Brain Res*. 2013;1532:76–84.
126. Fugl-Meyer AR, Jääskö L, Leyman I, Olsson S, Steglind S. The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance. *Scand J Rehabil Med*. 1975;7(1):13–31.
127. Granger CV, Gresham GE. *Functional Assessment in Rehabilitation Medicine*. Baltimore (MD): Williams and Wilkins; 1984.