




ORIGINAL ARTICLE

Risk and outcomes of COVID-19 in patients with multiple sclerosis

Irene Moreno-Torres¹ | Virginia Meca Lallana²  | Lucienne Costa-Frossard³ |
Celia Oreja-Guevara⁴ | Clara Aguirre² | Elda María Alba Suárez⁵ |
Mayra Gómez Moreno⁶  | Laura Borrega Canelo⁷ | Julia Sabín Muñoz⁸ |
Yolanda Aladro⁹ | Alba Cárcamo⁹ | Elena Rodríguez García¹⁰ | Juan Pablo Cuello¹¹ |
Enric Monreal³ | Susana Sainz de la Maza³ | Fernando Pérez Parra¹² |
Francisco Valenzuela Rojas¹³ | Carlos López de Silanes de Miguel¹⁴ | Ignacio Casanova¹⁴ |
María Luisa Martínez Gines¹¹ | Rosario Blasco⁸ | Aida Orviz García¹ |
Luisa María Villar-Guimerans³ | Guillermo Fernández-Dono¹⁵ | Víctor Elvira¹⁶ |
Carmen Santiuste³ | Mercedes Espiño³ | José Manuel García Domínguez¹¹ 

¹Demyelinating Diseases Unit, Jiménez Díaz Foundation University Hospital, Madrid, Spain

²Demyelinating Diseases Unit, La Princesa University Hospital, Madrid, Spain

³Multiple Sclerosis Unit, Ramón y Cajal University Hospital, Madrid, Spain

⁴Department of Neurology, Department of Medicine, Faculty of Medicine, San Carlos Clinical Hospital, Complutense University of Madrid and San Carlos Institute for Health Research, Madrid, Spain

⁵Neuroimmunology and Multiple Sclerosis Unit, La Paz University Hospital, Madrid, Spain

⁶Infanta Leonor University Hospital, Madrid, Spain

⁷Alcorcón Foundation University Hospital, Madrid, Spain

⁸Neuroimmunology Unit, Puerta de Hierro University Hospital, Madrid, Spain

⁹Demyelinating Diseases Unit, University Hospital of Getafe, Madrid, Spain

¹⁰Severo Ochoa University Hospital, Madrid, Spain

¹¹Demyelinating Diseases Unit, Gregorio Marañón General University Hospital, Madrid, Spain

¹²Henares University Hospital, Madrid, Spain

¹³Central Defense Hospital "Gómez Ulla," Madrid, Spain

¹⁴Torrejón University Hospital, Madrid, Spain

¹⁵Complutense University of Madrid, Madrid, Spain

¹⁶University of Edinburgh, Edinburgh, Scotland

Correspondence

José Manuel García Domínguez, Unidad de Enfermedades Desmielinizantes, Hospital General Universitario Gregorio Marañón, Calle Doctor Esquerdo, 46, 28007 Madrid, Spain.
Email: jgarcia Dominguez@salud.madrid.org

Funding information

No funding was received for the study design, collection of data, or statistical analysis of the results. Medical writing assistance was funded by Merck Spain.

Abstract

Background and purpose: Limited information is available on incidence and outcomes of COVID-19 in patients with multiple sclerosis (MS). This study investigated the risks of SARS-CoV-2 infection and COVID-19-related outcomes in patients with MS, and compared these with the general population.

Methods: A regional registry was created to collect data on incidence, hospitalization rates, intensive care unit admission, and death in patients with MS and COVID-19. National government outcomes and seroprevalence data were used for comparison. The study was conducted at 14 specialist MS treatment centers in Madrid, Spain, between February and May 2020.

Results: Two-hundred nineteen patients were included in the registry, 51 of whom were hospitalized with COVID-19. The mean age \pm standard deviation was 45.3 ± 12.4 years, and the mean duration of MS was 11.9 ± 8.9 years. The infection incidence rate was lower in patients with MS than the general population (adjusted incidence rate ratio = 0.78, 95% confidence interval [CI] = 0.70–0.80), but hospitalization rates were higher (relative risk = 5.03, 95% CI = 3.76–6.62). Disease severity was generally low, with only one admission to an intensive care unit and five deaths. Males with MS had higher incidence rates and risk of hospitalization than females. No association was found between the use of any disease-modifying treatment and hospitalization risk.

Conclusions: Patients with MS do not appear to have greater risks of SARS-CoV-2 infection or severe COVID-19 outcomes compared with the general population. The decision to start or continue disease-modifying treatment should be based on a careful risk-benefit assessment.

KEYWORDS

COVID-19, incidence, multiple sclerosis, severe acute respiratory syndrome coronavirus 2, Spain

INTRODUCTION

Patients with multiple sclerosis (MS) are at increased risk of infection and infectious complications, due to the effects of both the disease and disease-modifying treatments (DMTs) [1]. Infections are an important cause of hospitalization in patients with MS [2,3], with almost 8% of those hospitalized requiring intensive care [3]. Several factors are associated with an increased risk of infection-related hospitalization in patients with MS, including older age, longer disease duration, comorbidities, immunosuppression, prior hospital admission, obesity, and diabetes [2,4,5].

In December 2019, the first cases of pneumonia caused by an unknown virus were reported in Wuhan, China [6]. The pathogen causing the infection was subsequently identified and named SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), and on 11 March 2020 the World Health Organization declared a pandemic [7].

SARS-CoV-2 is a zoonotic RNA virus similar to SARS-CoV (severe acute respiratory syndrome coronavirus) and MERS-CoV (Middle East respiratory syndrome coronavirus). In infections caused by those coronaviruses, immunosuppression does not appear to be associated with an increased risk of severe disease or death [8]. To date, no increase in the development of clinical or severe lung disease has been reported in patients with SARS-CoV-2 infection who have undergone solid organ transplantation or received chemotherapy [8,9]. Specific data on outcomes in treated patients with MS and coronavirus infection are, however, scarce.

We performed an observational study to quantify the risks of infection, hospitalization, admission to intensive care, and death due to SARS-CoV-2 infection (henceforth referred to as COVID-19, except where the virus itself is being discussed) among patients with MS relative to the general population, and to identify factors associated with risk of hospitalization.

METHODS

Design

The study was conducted in the Madrid region of Spain. A regional COVID-19 registry was set up by members of the Study Group for Demyelinating Diseases, representing 14 MS centers across the region.

Between 9 February and 29 May 2020, all patients who were treated for MS at one of these centers and in whom COVID-19 was confirmed or highly suspected were included in the registry, regardless of whether they were admitted to hospital. Clinical and demographic data, laboratory test results, MS-specific information, and details of treatment for COVID-19 were entered into the registry.

This study was conducted according to the principles of the Declaration of Helsinki. The study protocol was approved by the ethics committee at the Jiménez Díaz Foundation University Hospital, Madrid; all patients gave informed consent to be included in the registry. Patient data were anonymized, and measures were implemented to prevent reidentification or access by unauthorized parties. Follow-up was until recovery from COVID-19 or death.

Definitions and outcomes

Confirmed COVID-19 was defined as a positive reverse transcriptase–polymerase chain reaction (RT-PCR) nasopharyngeal swab or serologic (enzyme-linked immunosorbent assay [ELISA]) test for SARS-CoV-2. COVID-19 was highly suspected in patients who had at least two cardinal symptoms of COVID-19 (e.g., persistent dry cough, difficulty breathing, anosmia/ageusia,

and fever) and/or positive radiologic findings, and a high circumstantial probability of infection in a pandemic setting (e.g., a positive test in a close family member), in the absence of a confirmatory test result.

The outcomes of interest were number of cases of COVID-19, COVID-19-related hospitalization, admission to an intensive care unit (ICU), and death.

Statistical analysis

Data on clinical variables, laboratory test results, COVID-19 treatment, and MS-related information were analyzed descriptively. Laboratory data from patients with MS were compared with corresponding data from matched (hospitalized) patients with COVID-19 but without MS; the latter were from a registry created by Dr. Luisa Maria Villar and colleagues at the Ramón y Cajal University Hospital, Madrid.

Binary and categorical variables were analyzed using contingency 2×2 tables, and significance was calculated using the chi-squared test. For continuous variables, a nonparametric Wilcoxon rank sum test was performed to compare subgroups (hospitalized vs. nonhospitalized patients; hospitalized MS patients vs hospitalized non-MS patients from the Villar et al. registry). Binary uni- and multivariate logistic regression was used to identify factors predictive of hospitalization. We use a penalized version of the multivariate logistic regression model, with L1 penalty (Lasso regression), to avoid the separation problem [10]. The penalized model also helps to enhance the interpretability of the results by excluding irrelevant variables. Odds ratios (ORs) and 95% confidence intervals (CIs) were computed for each variable, with $p < 0.05$ considered significant.

In addition, we calculated the incidence rate (IR; defined as the number of cases divided by the total population at risk) of COVID-19 for both the MS population and the general Spanish population aged 15–79 years; this range was chosen because it corresponds to the age distribution of the MS population. IRs for patients with MS were derived from registry and local data on the MS population, whereas data from the ongoing National Study of Serological Epidemiology of COVID-19 (NSSE-C19) [11] were used to generate IRs for the general Spanish population. In NSSE-C19, the percentage of the Spanish population infected with SARS-COV-2 was estimated by measuring the prevalence of antibodies to the new virus. The size of the MS population for the 14 centers was estimated from local pharmacy registries, and increased by 10% to account for untreated patients. IRs were then used to calculate IR ratios (IRRs), defined as the IR for the MS population divided by the IR for the general population, to allow for comparison between populations.

The risk of hospitalization in patients with MS and COVID-19, relative to the general COVID-19 population in Spain [12], was calculated by comparing the ratios of hospitalizations to cases in both populations. Lastly, the case-fatality rate (number of deaths divided

by number of reported cases) for the registry (COVID-19/MS) population was compared with national data [12].

RESULTS

Study population

Of 5641 patients with MS treated at the 14 participating centers, 219 (3.9%) had confirmed or highly suspected COVID-19, and were entered into the registry. COVID-19 was confirmed in 101 (46.1%) patients (RT-PCR, 88 patients [40.2%]; ELISA, 13 patients [5.9%]), and highly suspected in 118 (53.9%) patients.

The demographic and MS-related characteristics of patients are summarized in Table 1. Most (82.2%) patients had relapsing–remitting MS (RRMS), and most had stable disease; 182 (83.0%) patients experienced no relapses in the previous year. The mean age of MS cases was 45.3 years. Compared with the Spanish MS population [11,13], the registry had a higher percentage of patients aged 30–50 years; however, no significant difference was found in the percentage of cases by age group in the MS and general populations (Figure S1 in Appendix S1). Consistent with this, comorbidities of interest were uncommon in the COVID-19/MS population; among patients with PCR-confirmed disease, diabetes, pulmonary disease, and cardiovascular disease were all significantly less prevalent than in the corresponding general Spanish COVID-19 population (Table 2). Lymphopenia was absent in most patients for whom data were available (152/177, 85.9%). Patients with MS had broadly similar symptoms of COVID-19 to the general population, but had a significantly higher incidence of pneumonia, colds, and diarrhea (Table 2).

In our study population, 63% of COVID-19 cases were in females, similar to the general Spanish COVID-19 population ($p = 0.12$). However, after adjusting for the approximately threefold greater prevalence of MS in females than in males [14], the risk of developing COVID-19 was found to be 1.67-fold higher in males versus females with MS.

Outcome of COVID-19 in MS patients

Fifty-one patients (23.3%) were hospitalized due to COVID-19. One patient (0.45%) was admitted to ICU, but did not require mechanical ventilation. This patient, a 36-year-old male, had been diagnosed with RRMS 2 years previously, had an Expanded Disability Status Scale (EDSS) score of 2.0, and was receiving alemtuzumab.

Sequential Organ Failure Assessment (SOFA) and CURB-65 scores were available for 71 and 57 patients, respectively; only one patient had a SOFA score of greater than 4, and two patients had CURB-65 scores of 3 or greater.

Five patients died: four males aged 48–61 years and one female aged 57 years (Table S1 in Appendix S1). Their median EDSS score

TABLE 1 Baseline characteristics of the study population (N = 219)

Parameter	Value
Male sex, n (%)	81 (37.0)
Age, years	45.3 ± 12.4
Time since MS diagnosis, years	11.9 ± 8.9
Type of MS, n (%)	
Relapsing–remitting	180 (82.2)
Primary progressive	15 (6.8)
Secondary progressive	23 (10.5)
Clinically isolated syndrome	1 (0.5)
EDSS score	2.74 ± 2.28
Radiological disease activity in past year, n (%)	26 (11.9)
≥1 relapse in past year, n (%) ^a	33 (15.1)
Previous MS treatments, n	0.80 ± 0.93
DMT, n (%)	
Teriflunomide	33 (15.1)
Dimethyl fumarate	32 (14.6)
Natalizumab	27 (12.3)
Fingolimod	18 (8.2)
Interferon-β	16 (7.3)
Ocrelizumab	16 (7.3)
Rituximab	13 (5.9)
Alemtuzumab	11 (5.0)
Glatiramer acetate	10 (4.6)
Cladribine	9 (4.1)
Other drugs, including clinical trial medications	4 (1.8)
No DMT	30 (13.7)
Time since last treatment cycle, days	
Alemtuzumab	644 ± 525
Cladribine	334 ± 127
Rituximab	197 ± 66
Ocrelizumab	164 ± 57
Natalizumab	26 ± 18

Note: Continuous variables are expressed as mean ± SD unless otherwise indicated.

Abbreviations: DMT, disease-modifying treatment; EDSS, Expanded Disability Status Scale; MS, multiple sclerosis.

^aOne hundred eighty-two (83.0%) patients experienced no relapses in the previous year. Data were missing for four patients.

was 8.0 (range = 4.0–9.0), and time since diagnosis ranged from 3 to 27 years. All of those who died had secondary progressive MS (SPMS), and all except one had known bilateral pneumonia and lymphopenia. Only one of the deceased patients was taking DMT (rituximab).

The outcome of COVID-19 by DMTs is shown in Figure 1. Most of the MS patients who were hospitalized for COVID-19 were receiving teriflunomide, ocrelizumab, rituximab, or no treatment. MS patients receiving anti-CD20 monoclonal antibodies or not receiving

DMT were more likely to have SPMS or primary progressive MS (PPMS) compared with those receiving other treatments (Figure 1).

Factors influencing hospitalization risk

Compared with nonhospitalized patients, hospitalized patients in our registry were significantly older (mean age ± SD = 51.2 ± 13.7 vs. 43.4 ± 11.4, $p < 0.01$) and had higher EDSS scores (mean ± SD score = 4.26 ± 2.87 vs. 2.27 ± 1.85, $p < 0.01$), as shown in Figure 2 and Table S2 in Appendix S1. In addition, hospitalized patients had significantly lower lymphocyte counts, and significantly higher C-reactive protein and fibrinogen levels, than nonhospitalized patients. Compared with patients in the Villar et al. registry (hospitalized patients with COVID-19, but without MS), hospitalized patients from the study population were significantly younger and had lower C-reactive protein levels.

In the univariate logistic regression analysis, the relapsing–remitting phenotype was associated with a reduced risk of hospitalization, whereas progressive forms of MS, an EDSS of greater than 3.0, and a history of smoking, diabetes, or arterial hypertension were associated with an increased risk of hospitalization (Figure 2 and Table S3 in Appendix S1). However, in multivariate logistic regression, the only variable confirmed to reduce the risk of hospitalization was the relapsing–remitting phenotype (Figure 2; OR = 0.18, 95% CI = 0.08–0.41, $p < 0.001$). No variable or specific DMT was associated with an increased risk of hospitalization in the multivariate analysis.

Incidence of COVID-19 in MS patients

Using estimated IRs obtained from NSSE-C19 [11], the IRR for the registry population was 0.78 (95% CI = 0.70–0.80). IRRs were less than 1 for all subgroups defined by age, except for patients aged 15–29 years (Table 3; IRR = 1.04, 95% CI = 0.79–1.28). Compared with the general population, infection probability was lower for female patients with MS and higher for male patients with MS.

The likelihood of hospitalization due to COVID-19 was significantly higher in MS patients compared with the general Spanish population (Table 4; relative risk = 5.03, 95% CI = 3.76–6.62). Among those with MS, males were 1.5–3.0 times more likely to be hospitalized than females.

Five patients with MS died, giving a case–fatality ratio of 0.023. This was significantly lower than in the Madrid region overall (0.062; 4258 deaths/68,984 cases; $p = 0.024$), but not compared with Spain overall (0.041; 7678 deaths/187,703 cases; $p < 0.23$).

DISCUSSION

We present outcomes and incidence data from a registry of patients with MS diagnosed with COVID-19 at 14 centers in Madrid, one of the cities most affected during the first wave of the 2020

TABLE 2 Comorbidities, risk factors, and symptoms^a

	MS Patients with COVID-19, Madrid		General Spanish COVID-19 population [12]		p
	n ^b	Total (%)	n ^b	Total (%)	
Comorbidity or risk factor					
Diabetes	83	5 (6.0)	149,636	24,222 (16.2)	0.014 ^c
Pulmonary disease	85	2 (2.4)	149,636	16,454 (11.0)	0.018 ^c
Cardiovascular disease	85	2 (2.4)	149,636	43,420 (29.0)	<0.01 ^c
Arterial hypertension	76	11 (14.5)	149,636	31,800 (21.3)	0.072
Cancer	83	4 (4.8)	NA	NA	NA
Active smoker	62	7 (11.3)	NA	NA	NA
Ex-smoker	63	7 (11.1)	NA	NA	NA
Symptom					
Fever	87	65 (74.7)	101,523	74,057 (72.9)	0.70
Pneumonia	74	53 (71.6)	125,022	67,293 (53.8)	<0.01 ^c
Dry cough	86	53 (61.6)	93,105	64,281 (69.0)	0.18
Myalgia	85	44 (51.8)	NA	NA	NA
Fatigue	86	42 (48.8)	NA	NA	NA
Dyspnea	86	37 (43.0)	87,744	41,727 (47.6)	0.55
Sore throat	85	23 (27.1)	38,299	8430 (22.0)	0.17
Anosmia	79	19 (24.1)	NA	NA	NA
Ageusia	79	16 (20.3)	NA	NA	NA
Cold	87	14 (16.1)	38,003	8843 (23.3)	0.04 ^c
Diarrhea	86	12 (14.0)	39,799	10,678 (26.8)	0.04 ^c
Productive cough	87	10 (11.5)	NA	NA	NA

Abbreviations: MS, multiple sclerosis; NA, not available.

^aTable includes patients with COVID-19 confirmed by reverse transcriptase–polymerase chain reaction.

^bNumber of patients for whom data were available.

^cStatistically significant.

pandemic. Patients with MS in Madrid are in regular contact with neurologists, and are instructed to seek medical advice whenever they suspect infection. This allows for greater detection of positive cases. Additionally, the organization of MS care in Madrid, with specialist clinics in each hospital, allowed us to accurately estimate the incidence of MS in our population.

In the general population, older age is a clear risk factor for diagnosis of COVID-19 [15–17]. However, data from NSSE-C19 show that the distribution of antibodies to SARS-CoV-2 by age group is consistent with the population pyramid for Spain [11]. Thus, increasing age does not augment the risk of SARS-CoV-2 infection per se; however, it does increase the risk of a more severe clinical course, hospitalization, and death [16]. In agreement with this, we found that patients with MS who were hospitalized with COVID-19 were significantly older than those who were not hospitalized.

Because of a lack of published data specific to the Spanish population, we were not able to adequately adjust our results for both age and comorbidities. Patients in our population had a lower prevalence of relevant comorbidities compared with the general Spanish COVID-19 population, probably because they were, on

average, younger. These factors may partly explain the low rates of severe infection in our population, as measured by SOFA and CURB-65 scores; only one patient in our registry was admitted to an ICU. Interestingly, however, patients with MS were more likely to be hospitalized because of COVID-19 versus the general population. We postulate that this may be due to the use of admission criteria that prioritized patients with chronic comorbidities and/or taking immunosuppressive medication. Thus, despite having low disease severity, patients with MS may have been hospitalized as a precaution rather than because of risk to life, although the higher rates of pneumonia among MS patients may also have been a factor.

In agreement with the observations of other researchers [18–21], we also believe that the low disease severity in many of our patients may be related to a protective effect of therapeutic immunosuppression against cytokine storm. Cytokine storm is associated with increased morbidity and mortality in COVID-19 [15,22], and has been described as a consequence of lymphocyte dysregulation leading to overactivity in other parts of the immune response [22,23]. Most DMTs used in MS either deplete or modulate these overactive cells,




FIGURE 1 Outcome of COVID-19 in multiple sclerosis (MS) patients according to disease-modifying treatments (DMTs). The bars show the number of patients who were not hospitalized (blue), were hospitalized (orange), were admitted to an intensive care unit (ICU; green), or died (red) for each treatment. On the right, age, Expanded Disability Status Scale (EDSS) and MS phenotype are shown for each treatment. Continuous variables are presented as mean ± SD. Hosp, hospital; PPMS, primary progressive multiple sclerosis; RRMS, relapsing–remitting multiple sclerosis; SPMS, secondary progressive multiple sclerosis [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.com)]

and there is therefore a mechanistic basis for protection against cytokine storm in DMT-treated patients with MS.

Some authors have proposed that DMT-treated patients with MS are at a higher risk of infection with SARS-CoV-2 or of having a more severe outcome due to their immune status [24–27]. However, in our study, none of the DMTs used in MS was associated with an increased risk of hospitalization. Our results are similar to those reported by the Dutch group [28], and indicate that no apparent difference exists in DMT use and COVID-19 disease course.

In contrast to the Italian group [29], we did not find an association between the use of anti-CD20 monoclonal antibodies, interferon, or corticosteroids and the outcome of COVID-19 in MS patients. With regard to anti-CD20 monoclonal antibodies, we found a higher percentage of hospitalized patients taking these drugs, but these patients were also more likely to have SPMS or PPMS. For this reason, we did not find an increased risk of hospitalization for patients taking anti-CD20 monoclonal antibodies in multivariate analysis. We believe that treatment in this case acts as a dependent factor, and that the true factor involved in the risk of hospitalization is the MS phenotype.

The small number of patients who died or were admitted to ICU does not permit correlation analysis, but our findings suggest that the course of COVID-19 in patients with MS is, overall, no worse than in the general population.

Some DMTs can induce sustained lymphopenia. In our study, there were no differences in the need for hospitalization between patients who had or did not have lymphopenia before SARS-CoV-2 infection. The only characteristic found to have a significant impact on hospitalization risk in our population was relapsing–remitting disease, which was associated with a lower risk of hospitalization for COVID-19. This finding may be explained by better functional status and less disability in RRMS patients compared to those with

progressive forms of the disease. All five of the deaths in our registry were in patients with SPMS; patients with this form of MS have a longer duration of disease, and might be more immunologically senescent than patients with RRMS. This may predict a worse outcome of COVID-19.

Comparing our data with estimates based on data from the ongoing NSSE-C19 study allows for a more realistic calculation of the risk of COVID-19 in the MS population. In our analysis, patients with MS had a lower COVID-19 infection risk than the general Spanish population. However, to confirm this, we believe there is a need for an antibody prevalence study in the MS population to establish the exact risk of SARS-CoV-2 infection in MS patients compared with the general population.

Although men and women appear to have an equal risk of SARS-CoV-2 infection [30,31], the probability of a severe disease outcome or death is higher for men, both in Spain [12] and globally [30–32]. Our results in patients with MS are consistent with these findings. We found that the probability of COVID-19 (relative to the general population) was higher for male than female patients with MS. Additionally, the risk of hospitalization for COVID-19 (again, relative to the general population) was higher for males with MS than for females. Four of the five deaths in our study population were in men. The reasons for sex-based differences in disease course and outcome are not yet clear.

Our study has limitations that are common to many registry studies. Definitive cause-and-effect relationships can only be verified through well-designed prospective studies, and we cannot discount the influence of confounding factors on our findings. Moreover, although the study was conducted across multiple sites within the region, the results cannot necessarily be generalized to other settings within Spain or to other countries. Nevertheless, our findings make an important contribution to knowledge at this early stage in our




FIGURE 2 Risk factors for hospitalization in multiple sclerosis (MS) patients with COVID-19. (a, b) Comparison of median age (a) and Expanded Disability Status Scale (EDSS; b) between nonhospitalized patients and hospitalized patients (p -values from Wilcoxon rank sum test). Horizontal lines within each colored box indicate the median and interquartile range, whereas the whiskers indicate the range of values. Outliers are indicated by diamond symbols. (c, d) Risk factors for hospitalization, with odds ratios and 95% confidence intervals. (c) Univariate logistic regression. (d) Multivariate logistic regression with L1 penalty (Lasso regression). Complete logistic regression data are shown in Table S3 in Appendix S1. (d) Lymphocyte counts are $\times 10^9/L$; alemtuzumab, cladribine, fingolimod, glatiramer acetate, ocrelizumab, and rituximab were penalized in the Lasso regression, and are thus excluded. (c) Lymphocyte counts are in cells/ μ L. PPMS, primary progressive MS; RRMS, relapsing–remitting MS; SPMS, secondary progressive MS [Colour figure can be viewed at wileyonlinelibrary.com]

understanding of COVID-19 and its interaction with MS, and should be viewed in that context.

CONCLUSIONS

The risk of COVID-19 appears to be no greater in patients with MS than in the general Spanish population. We found a low risk of severe outcomes (ICU admission and death) in patients with MS and COVID-19, and a lower risk of COVID-19-related hospitalization for patients with RRMS versus other forms of MS. Our findings also suggest that outcomes of COVID-19 may be more severe in men, patients with SPMS, and those with greater disability. We did not find an association between the use of specific DMTs and an increased

risk of hospitalization. MS is a debilitating disease, and the potential risks and benefits of starting or continuing DMTs must be carefully considered, alongside other factors such as age and comorbidities, in the context of the current pandemic.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in Mendeley at <http://dx.doi.org/10.17632/cczx45n32y.1> [33].

ACKNOWLEDGMENTS

We would like to thank Richard Crampton of Springer Healthcare Communications for editorial assistance with the first and

TABLE 3 IR and IRR of COVID-19 in MS patients

	General population of Spain, estimated cases, % (95% CI) ^a	MS patients, Madrid			
		Cases	MS population ^c	IR	IRR (95% CI) ^b
Age group, years					
15-29	0.043 (0.034-0.055)	22	483	0.04555	1.04 (0.79-1.28)
30-39	0.042 (0.033-0.052)	54	1487	0.03631	0.86 (0.67-1.06)
40-49	0.055 (0.047-0.063)	80	2160	0.03704	0.67 (0.57-0.77)
50-59	0.059 (0.050-0.070)	38	994	0.03822	0.64 (0.53-0.74)
60-69	0.060 (0.050-0.072)	15	299	0.05016	0.83 (0.70-1.00)
70-79	0.065 (0.052-0.080)	10	218	0.04587	0.71 (0.57-0.88)
Total	0.050 (0.047-0.054)	219	5641	0.03882	0.78 (0.70-0.80)
Sex					
Female	0.051 (0.041-0.055)	139	4230	0.03286	0.64 (0.58-0.77)
Male	0.050 (0.046-0.054)	80	1411	0.05670	1.13 (1.04-1.22)
Total	0.050 (0.047-0.054)	219	5641	0.03882	0.78 (0.70-0.80)

Abbreviations: CI, confidence interval; IR, incidence rate; IRR, IR ratio; MS, multiple sclerosis.

^aData from National Study of Serological Epidemiology of COVID-19 in Spain, 13 May 2020 [11].

^bIRR for MS patients versus the overall Spanish population [11].

^cMS population of the 14 participating hospitals.

TABLE 4 RR of hospitalization for COVID-19 in MS patients

	General population, Spain			MS Patients, Madrid			
	Hospitalizations ^b	Total population ^c	Risk	Hospitalizations	MS population ^d	Risk	RR (95% CI) ^a
Age group, years							
15-29	1634	7,194,110	0.00022	3	483	0.00621	27.17 (8.76-84.11)
30-39	3825	6,322,990	0.00060	6	1487	0.00403	6.67 (3.04-14.81)
40-49	8709	7,832,487	0.00111	18	2160	0.00833	7.43 (4.65-11.76)
50-59	14,202	6,904,789	0.00206	12	994	0.01207	5.81 (3.31-10.24)
60-69	17,396	5,197,906	0.00335	5	299	0.01672	4.93 (2.14-11.75)
70-79	20,787	3,793,879	0.00548	7	218	0.03211	5.70 (2.75-11.75)
Total	66,553	37,276,161	0.00179	51	5641	0.00904	5.03 (3.76-6.62)
Sex							
Female	27,473	18,449,909	0.00149	22	4230	0.00520	3.47 (2.33-5.31)
Male	39,080	18,826,252	0.00208	29	1411	0.02055	9.68 (6.79-13.85)
Total	66,553	37,276,161	0.00179	51	5641	0.00904	5.03 (3.76-6.62)

Abbreviations: CI, confidence interval; MS, multiple sclerosis; RR, relative risk.

^aRR of hospitalization for MS patients versus the overall Spanish population are shown.

^bOfficial Spanish government data, 29 May 2020 [12].

^cData from Instituto Nacional de Estadística (www.ine.es).

^dMS population of the 14 participating hospitals.

subsequent drafts of the manuscript. This medical writing assistance was funded by Merck Spain.

CONFLICT OF INTEREST

I.M.-T. has received honoraria for lecturing or assistance for conference attendance from Bayer, Biogen-Idec, Merck, Teva, Novartis, Roche, and Genzyme, and research grants from Novartis. V.M.L.

has received consulting or speaking fees from Almirall, Biogen-Idec, Genzyme, Merck Serono, Novartis, Roche, Terumo, Sanofi-Genzyme, Teva, and Celgene. L.C.-F. has received funding for research projects or in the form of conference fees, mentoring, and assistance for conference attendance from Bayer, Biogen-Idec, Roche, Merck, Novartis, Almirall, Celgene, Biopas, Ipsen, and Sanofi-Genzyme. C.O.-G. has received speaker fees and/or has served

on the advisory boards of Biogen-Idec, Celgene, Merck, Novartis, Roche, Sanofi-Genzyme, and Teva. E.M.A.S. has received funding for conference fees and assistance for conference attendance from Sanofi-Genzyme, and participates as a subinvestigator and coordinator of a Roche-sponsored clinical trial. L.B.-C. has received honoraria for lecturing, consulting, or travel expenses from Bayer, Biogen-Idec, Merck, Teva, Novartis, Roche, Almirall, and Genzyme. J.S.M. has received honoraria for lecturing, consulting, or travel expenses from Biogen-Idec, Merck, Teva, Novartis, Roche, and Sanofi-Genzyme. Y.A. has received funding for research projects, conference fees, mentoring, and assistance for conference attendance from Teva, Merck, Biogen-Idec, Roche, Merck, Novartis, Almirall, and Sanofi-Genzyme. E.R.G. has received honoraria for lecturing or assistance for conference attendance from Bayer, Biogen-Idec, Merck, Teva, Novartis, Roche, Almirall, and Sanofi-Genzyme. J.P.C. has received funding for research projects from Biogen-Idec, and assistance for conference attendance from Biogen-Idec, Roche, and Sanofi-Genzyme. E.M. has received honoraria for lecturing or assistance for conference attendance from Roche, Sanofi-Genzyme, Almirall, Biogen-Idec, and Merck, and research grants from Merck. S.S.d.I.M. has received funding for research projects or support in the form of conference fees, mentoring, and assistance for conference attendance from Sanofi-Genzyme, Biogen-Idec, Roche, Merck-Serono, Novartis, Teva, Bayer, Almirall, and Mylan. F.V.R. has received support for scientific meetings and courses, and honoraria for advisory work, from Sanofi-Genzyme, Merck-Serono, Biogen-Idec, Novartis, Bial, Daichii-Sankyo, and Roche. I.C. has received funding for travel or speaker honoraria from Bayer, Biogen-Idec, Genzyme, Merck, Novartis, Roche, Sanofi-Genzyme, and Teva, and research support from Teva. M.L.M.G. has received compensation for consulting services and speaking fees from Merck, Biogen-Idec, Novartis, Sanofi-Genzyme, Almirall, Bayer, Roche, and Teva. A.O.G. has received honoraria for lecturing, consulting or travel expenses from Merck, Biogen-Idec, Sanofi-Genzyme, Novartis, Bayer, Roche, Almirall, and Teva. J.M.G.D. has received speaker fees or advisor honoraria from Biogen-Idec, Celgene, AbbVie, Sanofi-Genzyme, Almirall, Novartis, Roche, Teva, and Merck. None of the other authors has any conflict of interest to disclose.

AUTHOR CONTRIBUTIONS

Irene Moreno-Torres: Conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal). **Virginia Meca Lallana:** Conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal). **Lucienne Costa-Frossard:** Conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal). **Celia Oreja-Guevara:** Investigation (equal). **Clara Aguirre:** Investigation (equal). **Elda María Alba Suárez:** Investigation (equal). **Mayra Gómez Moreno:** Investigation (equal). **Laura Borrega Canelo:** Investigation (equal). **Julia Sabín Muñoz:** Investigation (equal). **Yolanda Aladro:** Investigation (equal). **Alba Cárcamo:** Investigation (equal). **Elena Rodríguez García:** Investigation (equal). **Juan Pablo Cuello:** Investigation (equal). **Enric Monreal:** Investigation (equal). **Susana Sainz de la Maza:** Investigation (equal).

Fernando Pérez Para: Investigation (equal). **Francisco Valenzuela Rojas:** Investigation (equal). **Carlos López de Silanes de Miguel:** Investigation (equal). **Ignacio Casanova:** Investigation (equal). **Marisa Luisa Martínez Gines:** Investigation (equal). **Rosario Blasco:** Investigation (equal). **Aida Orviz García:** Investigation (equal). **Luisa María Villar-Guimerans:** Investigation (equal). **Guillermo Fernández-Dono:** Formal analysis (equal), software (equal). **Víctor Elvira:** Formal analysis (equal), software (equal). **Carmen Santiuste:** Investigation (equal). **Mercedes Espiño:** Investigation (equal). **José Manuel García Domínguez:** Conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal).

ORCID

Virginia Meca Lallana  <https://orcid.org/0000-0001-9418-4127>

Mayra Gómez Moreno  <https://orcid.org/0000-0002-4009-4561>

José Manuel García Domínguez  <https://orcid.org/0000-0002-5205-3652>

REFERENCES

- Williamson EM, Berger JR. Infection risk in patients on multiple sclerosis therapeutics. *CNS Drugs*. 2015;29:229-244. <https://doi.org/10.1007/s40263-015-0226-2>
- Al-Sakran LH, Marrie RA, Blackburn DF, Knox KB, Evans CD. Predictors of hospitalization in a Canadian MS population: a matched cohort study. *Mult Scler Relat Disord*. 2020;41:102028. <https://doi.org/10.1016/j.msard.2020.102028>
- Maia C, Costa A, Abreu P, et al. All-cause hospitalizations in multiple sclerosis patients. *Rev Neurol*. 2019;68:229-235. <https://doi.org/10.33588/rn.6806.2018281>
- Evans C, Kingwell E, Zhu F, et al. Hospital admissions and MS: temporal trends and patient characteristics. *Am J Manag Care*. 2012;18:735-742.
- World Health Organization. Seasonal influenza factsheets. <http://www.who.int/mediacentre/factsheets/fs211/en/>. 2018. Accessed June 22, 2020.
- World Health Organization. Novel coronavirus – China. www.who.int/csr/don/12-january-2020-novel-coronavirus-china/en/. 2020. Accessed May 22, 2020.
- World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. Accessed June 23, 2020.
- D'Antiga L. Coronaviruses and immunosuppressed patients: the facts during the third epidemic. *Liver Transpl*. 2020;26(6):832-834. <https://doi.org/10.1002/lt.25756>
- Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*. 2020;395:1033-1034. [https://doi.org/10.1016/S0140-6736\(20\)30628-0](https://doi.org/10.1016/S0140-6736(20)30628-0)
- Zorn C. A solution to separation in binary response models. *Political Analysis*. 2005;13:157-170. <https://doi.org/10.1093/pan/mpi009>
- Gobierno de España. Estudio ENE-COVID19: primera ronda Estudio Nacional de Sero-epidemiología de la Infección por SARS-CoV-2 en España: informe preliminar 13 de mayo de 2020. https://d3cra5ec8gdi8w.cloudfront.net/uploads/documentos/2020/05/13/_estudio_f63dc898.pdf. Accessed June 24, 2020.
- Ministerio de Sanidad. Análisis de los casos de COVID-19 notificados a la RENAVE hasta el 10 de mayo en España. Informe COVID-19 no. 33. 29 de mayo de 2020. <https://www.isciii.es/QueHacemos/>

- Servicios/VigilanciaSaludPublicaRENAVE/EnfermedadesTransmisibles/Paginas/InformesCOVID-19.aspx. Accessed July 6, 2020.
13. Otero-Romero S, Roura P, Sola J, et al. Increase in the prevalence of multiple sclerosis over a 17-year period in Osona, Catalonia, Spain. *Multiple Scler*. 2013;19(2):245-248. <https://doi.org/10.1177/1352458512444751>
 14. Wallin MT, Culpepper WJ, Campbell JD, et al. The prevalence of MS in the United States: a population-based estimate using health claims data. *Neurology*. 2019;92(10):e1029-e1040. <https://doi.org/10.1212/WNL.0000000000007035>
 15. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-1720. <https://doi.org/10.1056/NEJMoa2002032>
 16. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular disease, drug therapy, and mortality in Covid-19. *N Engl J Med*. 2020;382:e102. <https://doi.org/10.1056/NEJMoa2007621>
 17. Natale F, Ghio D, Tarchi D, et al. COVID-19 cases and case fatality rate by age. https://ec.europa.eu/knowledge4policy/sites/known4pol/files/jrc120420_covid_risk_and_age.pdf. 2020. Accessed August 7, 2020.
 18. Novi G, Mikulska M, Briano F, et al. COVID-19 in a MS patient treated with ocrelizumab: does immunosuppression have a protective role? *Mult Scler Relat Disord*. 2020;42:102120. <https://doi.org/10.1016/j.msard.2020.102120>
 19. Siddiqi HK, Mehra MR. COVID-19 illness in native and immunosuppressed states: a clinical–therapeutic staging proposal. *J Heart Lung Transplant*. 2020;39:405-407. <https://doi.org/10.1016/j.healun.2020.03.012>
 20. Willis MD, Robertson NP. Multiple sclerosis and the risk of infection: considerations in the threat of the novel coronavirus, COVID-19/SARS-CoV-2. *J Neurol*. 2020;267(5):1567-1569. <https://doi.org/10.1007/s00415-020-09822-3>
 21. Louapre C, Collongues N, Stankoff B, et al. Clinical characteristics and outcomes in patients with coronavirus disease 2019 and multiple sclerosis. *JAMA Neurol*. 2020;77:1079-1088. <https://doi.org/10.1001/jamaneurol.2020.2581>
 22. Qin C, Zhou L, Hu Z, et al. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clin Infect Dis*. 2020;71:762-768. <https://doi.org/10.1093/cid/ciaa248>
 23. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *JAMA*. 2020;323(15):1488. <https://doi.org/10.1001/jama.2020.3204>
 24. Amor S, Baker D, Houry SJ, et al. SARS-CoV-2 and multiple sclerosis: not all immune depleting DMTs are equal or bad. *Ann Neurol*. 2020;87:794-797. <https://doi.org/10.1002/ana.25770>
 25. Berger JR, Brandstadter R, Bar-Or A. COVID-19 and MS disease-modifying therapies. *Neurol Neuroimmunol Neuroinflamm*. 2020;7:e761. <https://doi.org/10.1212/NXI.0000000000000761>
 26. Costa-Frossard L, Moreno-Torres I, Meca-Lallana V, et al. EMCAM (Multiple Sclerosis Autonomous Community of Madrid) document for the management of patients with multiple sclerosis during the SARS-CoV-2 pandemic. *Rev Neurol*. 2020;70:329-340. <https://doi.org/10.33588/rn.7009.2020155>
 27. Mansoor S, Kelly S, Murphy K, et al. COVID-19 pandemic and the risk of infection in multiple sclerosis patients on disease modifying therapies: “what the bleep do we know?”. *Egypt J Neurol Psychiatr Neurosurg*. 2020;56:44. <https://doi.org/10.1186/s41983-020-00177-0>
 28. Loonstra FC, Hoitsma E, van Kempen ZL, Killestein J, Mostert JP. COVID-19 in multiple sclerosis: The Dutch experience. *Mult Scler*. 2020;26(10):1256-1260. <https://doi.org/10.1177/1352458520942198>
 29. Sormani MP, De Rossi N, Schiavetti I, et al. Disease modifying therapies and COVID-19 severity in multiple sclerosis. *Ann Neurol*. 2021;89(4):780-789. <https://doi.org/10.1002/ana.26028>
 30. La Vignera S, Cannarella R, Condorelli RA, et al. Sex-specific SARS-CoV-2 mortality: among hormone-modulated ACE2 expression, risk of venous thromboembolism and hypovitaminosis D. *Int J Mol Sci*. 2020;21:2948. <https://doi.org/10.3390/ijms21082948>
 31. Wenham C, Smith J, Morgan R, et al. COVID-19: the gendered impacts of the outbreak. *Lancet*. 2020;395:846-848. [https://doi.org/10.1016/S0140-6736\(20\)30526-2](https://doi.org/10.1016/S0140-6736(20)30526-2)
 32. Jin JM, Bai P, He W, et al. Gender differences in patients with COVID-19: focus on severity and mortality. *Front Public Health*. 2020;8:152. <https://doi.org/10.3389/fpubh.2020.00152>
 33. Moreno-Torres I, Meca Lallana V, Costa-Frossard L, et al. Database COVID-19 EMCAM Registry; Mendeley; Version 1. 2020. <https://doi.org/10.17632/cczx45n32y1>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Moreno-Torres I, Meca Lallana V, Costa-Frossard L, et al. Risk and outcomes of COVID-19 in patients with multiple sclerosis. *Eur J Neurol*. 2021;28:3712-3721. <https://doi.org/10.1111/ene.14990>