

SHORT COMMUNICATION

Genomic potential for nitrogen assimilation in uncultivated members of *Prochlorococcus* from an anoxic marine zone

This article has been corrected since Advance Online Publication and an erratum is also printed in this issue

Marcia Astorga-Eló^{1,2}, Salvador Ramírez-Flandes^{1,3}, Edward F DeLong^{4,5,6} and Osvaldo Ulloa¹

¹Departamento de Oceanografía and Instituto Milenio de Oceanografía, Universidad de Concepción, Concepción, Chile; ²Programa de Magíster en Bioquímica y Bioinformática, Facultad de Ciencias Biológicas, Universidad de Concepción, Concepción, Chile; ³Programa de Doctorado en Ingeniería de Sistemas Complejos, Universidad Adolfo Ibáñez, Santiago, Chile; ⁴Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA; ⁵Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA and ⁶Daniel K. Inouye Center for Microbial Oceanography: Research and Education, University of Hawaii, Honolulu, HI, USA

Cyanobacteria of the genus *Prochlorococcus* are the most abundant photosynthetic marine organisms and key factors in the global carbon cycle. The understanding of their distribution and ecological importance in oligotrophic tropical and subtropical waters, and their differentiation into distinct ecotypes, is based on genetic and physiological information from several isolates. Currently, all available *Prochlorococcus* genomes show their incapacity for nitrate utilization. However, environmental sequence data suggest that some uncultivated lineages may have acquired this capacity. Here we report that uncultivated low-light-adapted *Prochlorococcus* from the nutrient-rich, low-light, anoxic marine zone (AMZ) of the eastern tropical South Pacific have the genetic potential for nitrate uptake and assimilation. All genes involved in this trait were found syntenic with those present in marine *Synechococcus*. Genomic and phylogenetic analyses also suggest that these genes have not been acquired recently, but perhaps were retained from a common ancestor, highlighting the basal characteristics of the AMZ lineages within *Prochlorococcus*.

The ISME Journal (2015) 9, 1264–1267; doi:10.1038/ismej.2015.21; published online 20 February 2015

Cyanobacteria of the genus *Prochlorococcus* are the most abundant photosynthetic microorganisms inhabiting the oceans, key factors in the carbon cycle and a model organism in environmental microbiology (Partensky and Garczarek, 2010). They can be broadly classified into high-light and low-light (LL)-adapted ecotypes (Rocap *et al.*, 2002). These ecotypes exhibit distinct distributions both vertically in the water column and geographically across oligotrophic tropical and subtropical waters (Bouman *et al.*, 2006; Johnson *et al.*, 2006; Zwirgmaier *et al.*, 2008).

In past years, the genomes of over a dozen isolates of *Prochlorococcus* have been fully sequenced (for example, Kettler *et al.*, 2007) and over a hundred

single-cell-amplified partial genomes have been described (Malmstrom *et al.*, 2013; Kashtan *et al.*, 2014). All of them have revealed that they cannot use nitrate as a nitrogen source. However, new uncultivated lineages of *Prochlorococcus* have been identified in the environment using culture-independent techniques based on the sequencing of the 16S rRNA gene and related genomic regions (Lavin *et al.*, 2010; West *et al.*, 2011; Mühling, 2012; Malmstrom *et al.*, 2013). On the other hand, nitrate assimilation rates were reported for uncultivated deep populations of *Prochlorococcus* in the Western Atlantic Ocean (Casey *et al.*, 2007). In addition, genes necessary for nitrate assimilation associated to *Prochlorococcus* were identified in the global ocean sampling metagenomic database (Martiny *et al.*, 2009) and in metagenomes of flow-cytometry-sorted *Prochlorococcus* populations (Batmalle *et al.*, 2014).

Important uncultivated *Prochlorococcus* lineages include those thriving in anoxic marine zones (AMZs), where oxygen concentrations fall below the detection limit of modern sensors, light is scarce,

Correspondence: O Ulloa, Departamento de Oceanografía and Instituto Milenio de Oceanografía, Universidad de Concepción, Casilla 160-C, Concepción 4070386, Chile.

E-mail: oulloa@udec.cl

Received 29 August 2014; revised 19 December 2014; accepted 31 December 2014; published online 20 February 2015




Figure 2 Phylogenetic trees for nitrate assimilation and uptake genes. Maximum-likelihood phylogenetic trees of (a) *narB*- and (b) *napA*-predicted amino acid sequences found in contig 51148. Evolutionary history was inferred using neighbour joining (NJ), maximum parsimony (MP) and maximum likelihood (ML). Bootstrap support values for 100 replications are shown at the nodes (NJ/MP/ML).

60 m depth within the AMZ region of the eastern tropical South Pacific (Supplementary Figure S1), where dissolved oxygen was undetectable and inorganic nutrients were abundant (Supplementary Figure S2a; Thamdrup *et al.*, 2012). The microbial community was enriched in *Prochlorococcus*, shown to comprise ~10% of cell abundance, versus ~0.7% of *Synechococcus*, assessed by flow cytometry (Supplementary Figure S2b). Blast analysis of the taxonomic affiliation of sequences matching the *rpoC* region 1, a taxonomic marker for cyanobacteria based on a single-copy gene (Palenik, 1994), showed an *rpoC* gene relative abundance of 86% for *Prochlorococcus* and 14% for *Synechococcus* (Supplementary Table S1), supporting the flow cytometry results. Moreover, of the 15% protein-coding sequences assigned to cyanobacteria, 10% binned with *Prochlorococcus* and 5% with *Synechococcus* (Supplementary Figure S3). Of those assigned to *Prochlorococcus*, 90% were related to the LL ecotypes MIT9313 and MIT9303, the closest reported relatives to the AMZ lineages with genomes fully sequenced (Lavin *et al.*, 2010). General statistics of this AMZ metagenome are shown in Supplementary Tables S2 and S3.

Analysis of *de novo*-assembled contigs revealed the presence of several large contigs that binned with *Prochlorococcus*. In particular, a single contig was found to encode genes related to urea and nitrate uptake and assimilation (contig 51148, GenBank accession number KM282015; 10 300 bp; Figure 1), in synteny with those in *Synechococcus* WH8102. The genes in the urease gene cluster (*ureABCD*) presented high identity to those described for *Prochlorococcus* MIT9313 and MIT9303 (Rocap *et al.*, 2003; Supplementary Figure S4). Notably, the nitrate/nitrite transporter *napA* and assimilatory nitrate reductase *narB* were also found within the same contig

(Figure 1a), as well as the genes *moeA* and *mobA* (Supplementary Figure S5) involved in the biosynthesis of the Mo-cofactor and necessary for the *narB* function (Flores *et al.*, 2005). None of these genes have been found in any of the genomes of *Prochlorococcus* sequenced and described so far. However, homologues that presumably come from uncultivated relatives of *Prochlorococcus* have been found in the global ocean sampling database (Martiny *et al.*, 2009) and in metagenomes of uncultured, sorted *Prochlorococcus* populations (Batmalle *et al.*, 2014).

The GC content of contig 51148 was ~51.1% (Figure 1b) and similar to that of LL *Prochlorococcus* and some marine *Synechococcus* (Kettler *et al.*, 2007). Likewise, the *narB* gene had a GC content of 52%, which is less than the ~60% of those in the marine *Synechococcus* strains WH8102 and WH7803 (to which it presented the highest nucleotide identity), but significantly higher than the ~40% GC of the global ocean sampling high-light *Prochlorococcus narB* (Supplementary Figure S6). Analysis of codon usage patterns (Yu *et al.*, 2012) and tetranucleotide frequencies (see Supplementary Material and Methods) showed that the cyanobacterial portion of the metagenome and contig 51148 exhibit the highest similarity with LL *Prochlorococcus* MIT9303 (Figures 1d and e). Additionally, nucleotide identities and phylogenetic analysis confirmed that the urease genes of contig 51148 were associated more closely with *Prochlorococcus* than *Synechococcus* (Supplementary Table S4 and Supplementary Figure S4).

The homogeneous GC content of contig 51148, the differences in codon usage bias with *Synechococcus* and phylogenetic analyses of AMZ *narB* and *napA* (Figures 2a and b) all suggest that the genetic potential for nitrate uptake and assimilation was not obtained recently by horizontal gene transfer, but instead potentially were retained from a common ancestor

with *Synechococcus*. Mapping the presence/absence of the different nitrate utilization genes onto the cyanobacteria 16S rRNA phylogenetic tree is consistent with this hypothesis (Supplementary Figure S7).

In summary, our results indicate that AMZ *Prochlorococcus* lineages have the genetic potential for urea and nitrate assimilation, likely an adaptation to the unique nutrient-rich environment where they thrive. Additional genomic characteristics that could explain their high abundance in the oxygen-deficient and very-LL waters of AMZs remain to be assessed.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank the officers, crew and scientific party on board *R/V Vaedderen* during the Galathea-3 Expedition. This study was supported by the Chilean National Commission for Scientific and Technological Research (CONICYT), grant Fondecyt 1130784, and the Agouron Institute grant AI-MO5_08_3. MA-E was supported by a graduate fellowship from CONICYT. Additional support was provided by the Millennium Scientific Initiative, grant IC 120019.

References

- Batmalle CS, Chiang HI, Zhang K, Lomas MV, Martiny AC. (2014). Development and bias assessment of a method for targeted metagenomic sequencing of marine cyanobacteria. *Appl Environ Microbiol* **80**: 1116–1125.
- Bouman HA, Ulloa O, Scanlan DJ, Zwirgmaier K, Li WKW, Platt T *et al.* (2006). Oceanographic basis of the global surface distribution of *Prochlorococcus* ecotypes. *Science* **312**: 918–921.
- Casey JR, Lomas MW, Mandecki J, Walker DE. (2007). *Prochlorococcus* contributes to new production in the Sargasso Sea deep chlorophyll maximum. *Geophys Res Lett* **34**: L10604.
- Flores E, Frías JE, Rubio LM, Herrero A. (2005). Photosynthetic nitrate assimilation in cyanobacteria. *Photosynth Res* **83**: 117–133.
- Goericke R, Olson RJ, Shalapyonok A. (2000). A novel niche for *Prochlorococcus* sp. in low-light suboxic environments in the Arabian Sea and the Eastern Tropical North Pacific. *Deep Sea Res I Oceanogr Res Pap* **47**: 1183–1205.
- Johnson ZI, Zinser ER, Coe A, McNulty NP, Woodward EMS, Chisholm SW. (2006). Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. *Science* **311**: 1737–1740.
- Kashtan N, Roggensack SE, Rodrigue S, Thompson JW, Biller SJ, Coe A *et al.* (2014). Single-cell genomics reveals hundreds of coexisting subpopulations in wild *Prochlorococcus*. *Science* **344**: 416–420.
- Kettler GC, Martiny AC, Huang K, Zucker J, Coleman ML, Rodrigue S *et al.* (2007). Patterns and implications of gene gain and loss in the evolutions of *Prochlorococcus*. *PLoS Genet* **3**: e231.
- Lavin P, González B, Santibáñez JF, Scanlan DJ, Ulloa O. (2010). Novel lineages of *Prochlorococcus* thrive within the oxygen minimum zone of the eastern tropical South Pacific. *Environ Microbiol Rep* **2**: 728–738.
- Malmstrom RR, Rodrigue S, Huang KH, Kelly L, Kern SE, Thompson A *et al.* (2013). Ecology of uncultured *Prochlorococcus* clades revealed through single-cell genomics and biogeographic analysis. *ISME J* **7**: 184–198.
- Martiny AC, Kathuria S, Berube PM. (2009). Widespread metabolic potential for nitrite and nitrate assimilation among *Prochlorococcus* ecotypes. *Proc Natl Acad Sci USA* **106**: 10787–10792.
- Mühling M. (2012). On the culture-independent assessment of the diversity and distribution of *Prochlorococcus*. *Environ Microbiol* **14**: 567–579.
- Palenik B. (1994). Cyanobacterial community structure as seen from RNA polymerase gene sequence analysis. *Appl Environ Microbiol* **60**: 3212–3219.
- Partensky F, Garczarek L. (2010). *Prochlorococcus*: advantages and limits of minimalism. *Ann Rev Mar Sci* **2**: 305–331.
- Rocap G, Larimer FW, Lamerdin J, Malfatti S, Chain P, Ahlgren NA *et al.* (2003). Genome divergence in two *Prochlorococcus* ecotypes reflects oceanic niche differentiation. *Nature* **424**: 1042–1047.
- Rocap G, Distel DL, Waterbury JB, Chisholm SW. (2002). Resolution of *Prochlorococcus* and *Synechococcus* ecotypes by using 16S-23S ribosomal DNA internal transcribed spacer sequences. *Appl Environ Microbiol* **68**: 1180–1191.
- Thamdrup B, Dalsgaard T, Revsbech NP. (2012). Widespread functional anoxia in the oxygen minimum zone of the Eastern South Pacific. *Deep Sea Res Part I Oceanogr Res Pap* **65**: 36–45.
- Ulloa O, Canfield DE, DeLong EF, Letelier RM, Stewart FJ. (2012). Microbial oceanography of anoxic oxygen minimum zones. *Proc Natl Acad Sci USA* **109**: 15996–156003.
- West NJ, Lebaron P, Strutton PG, Suzuki MT. (2011). A novel clade of *Prochlorococcus* found in high nutrient low chlorophyll waters in the South and Equatorial Pacific Ocean. *ISME J* **5**: 933–944.
- Yu T, Li J, Yang Y, Qi L, Chen B, Zhao F *et al.* (2012). Codon usage patterns and adaptive evolution of marine unicellular cyanobacteria *Synechococcus* and *Prochlorococcus*. *Mol Phylogenet Evol* **62**: 206–213.
- Zwirgmaier K, Jardillier L, Ostrowski M, Mazard S, Garczarek L, Vaultot D *et al.* (2008). Global phylogeography of marine *Synechococcus* and *Prochlorococcus* reveals a distinct partitioning of lineages among oceanic biomes. *Environ Microbiol* **10**: 147–161.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)