

Parental stress before, during, and after pediatric stem cell transplantation: a review article

C. M. J. Vrijmoet-Wiersma · R. M. Egeler ·
H. M. Koopman · A. Lindahl Norberg ·
M. A. Grootenhuis

Received: 16 December 2008 / Accepted: 18 June 2009 / Published online: 2 July 2009
© The Author(s) 2009. This article is published with open access at Springerlink.com

Abstract

Goals of work Pediatric stem cell transplantation (SCT) is a stressful treatment for children with relapsed or high-risk malignancies, immune deficiencies and certain blood diseases. Parents of children undergoing SCT can experience ongoing stress related to the SCT period. The aim of this article was to present a literature review of articles on parental distress and adaptation before, during, and after SCT and to identify risk and protective factors.

Materials and methods The review was conducted systematically by using PubMed, Web of Science, PsychInfo, and Picarta databases. Eighteen articles met our inclusion criteria: publishing date between January 1, 1990 and January 1, 2009; studies concerning parents of children undergoing SCT; studies examining the psychological adjustment and/or stress reactions of parents as primary outcomes and studies available in English.

Main results Highest levels of parental stress are reported in the period preceding SCT and during the acute phase.

Stress levels decrease steadily after discharge in most parents. However, in a subgroup of parents, stress levels still remain elevated post-SCT. Parents most at risk in the longer term display highest levels of stress during the acute phase of the SCT.

Conclusions Psychosocial assessment before SCT, during the acute phase and in the longer term, is necessary to identify parents in need for support and follow-up care.

Keywords Pediatric SCT · Parental stress · Adaptation · Review

Introduction

Stem cell transplantation (SCT) is an invasive treatment for seriously ill children who have hematological, oncological, or metabolic diseases. Recently, for some high-risk leukemia protocols, SCT has become close to being a first-choice treatment. SCT is a perilous treatment, associated with significant mortality and morbidity [5]. It involves a lengthy hospital admission in an isolated environment to prevent infections and treatment with high doses of chemotherapy and/or radiation, followed by infusions of donor stem cells [22]. During the acute phase of SCT, children report high levels of somatic distress, mood disturbance [30], nausea and pain [7], and fatigue and malaise [29]. In the first 4–6 months post-SCT, children are still susceptible to infections and need to live with restrictions. SCT has a profound impact on the lives of children and parents, both during the acute phase and afterwards. Parents are faced with the need to provide both physical and emotional care for their ill child during a long and stressful period. Furthermore, they have to deal with their own emotions, especially with the realistic fear of

C. M. J. Vrijmoet-Wiersma (✉) · R. M. Egeler
Pediatric Department, Leiden University Medical Center,
P.O. Box 9600, 2300 RC Leiden, The Netherlands
e-mail: c.m.j.vrijmoet-wiersma@lumc.nl

H. M. Koopman
Medical Psychology, Leiden University Medical Center,
Leiden, The Netherlands

A. L. Norberg
Childhood Cancer Research Unit,
Department of Woman and Child Health, Karolinska Institutet,
Stockholm, Sweden

M. A. Grootenhuis
Pediatric Psychosocial Department, Emma Children's Hospital,
Academic Medical Center Amsterdam,
Amsterdam, The Netherlands

losing their child and they have to make complicated decisions about the treatment together with the multidisciplinary team. Some parents are also faced with supporting one of their other children who will be acting as a sibling donor and increasingly, parents are acting as haploidentical donors themselves in case no appropriate donor has been found.

Despite improved survival rates, SCT remains a high-risk procedure. The result of the transplantation depends on several risk factors, including type and status of the underlying disease [5]. After treatment, parents and children are faced with the risk of recurrence, acute or chronic graft-versus-host disease, and numerous possible late effects such as pulmonary disease, growth problems, and infertility [4, 21]. Even in the longer term, the child's illness and the SCT may influence parents' everyday lives [8]. SCT treatment protocols have changed in the past decades, one of the most important differences between treatment now and in the 1990s is the shortened admission period. On the one hand this is an improvement because parents may have fewer concerns about the practical issues during admission, e.g. being away from home for a long period and dealing with work-related stress. On the other hand, caring for a child at home post-SCT can be a heavy burden on parents and families. In addition, increasing survival rates entail increasing numbers of survivors with possible long-term side effects. Moreover, the fear of losing the child is still realistic. Accordingly, changed treatment protocols may not make any difference for parental stress levels.

The field of parental adaptation to their child's SCT has gained more attention in the past decade; most studies have been conducted in the past 8 years [18, 31]. The majority of studies have focused on parental stress and adjustment pre-SCT, during the acute phase, and 3, 6, or 18 months post-SCT, e.g. [5, 18]. These time points seem to cover the SCT time frame well: 3 months post-SCT, many children still suffer from the after effects of the SCT, whereas after 6 months, most of the children can return to school and pick up their old lives, even if their health is suboptimal [1]. Twelve to eighteen months post-SCT, the majority of children report to have a health-related quality of life (HRQoL) comparable to healthy peers [2, 7]. However, certain aspects of HRQoL seem to be affected in the long-term in many survivors, for example cognitive functions and pain, which has been demonstrated recently by Löf et al. [15].

No review articles have been published in this specific area so far. In related areas, however, review studies have focused on adjustment and coping of parents of children with cancer [9, 40]; on the quality of life and/or emotional adjustment of children after SCT [2, 38]; on the psychological adjustment of families of adult SCT patients [14] and on the psychosocial impact of SCT of adult patients [11, 23]. The aim of our article was to conduct a systematic

review of the current literature (1990 to 2008) on parental distress and adaptation to their child's SCT and to identify risk and protective factors.

Materials and methods

Search strategy for identification of studies

Several research engines were used to obtain the studies included in this review: Pubmed, Web of Science, PsychInfo, and Picarta. These databases were searched for one of the words: BMT, bone marrow transplantation, SCT, or stem cell transplantation combined with the following words used in headings, keywords, subjects, or abstracts: pediatric, paediatric, parent, child, mother, father, AND/OR stress, distress, post-traumatic stress disorder (PTSD), post-traumatic stress symptoms (PTSS), anxiety, depression, parental stress, parental distress, psychological, adjustment, and coping. Subsequently, reference lists of the relevant articles were examined to identify additional papers that met the search criteria and a hand search was conducted in several relevant academic journals.

Criteria for considering studies for the review

Criteria for inclusion were: publishing date between January 1, 1990 and January 1, 2009; studies concerning parents of children undergoing SCT; studies examining the psychological adjustment and/or stress reactions of parents as a primary outcome; and studies available in English. Exclusion criteria were: reviews, guidelines, protocols, commentaries, and other descriptive articles; studies focusing on psychological adjustment of pediatric SCT patients only; studies focusing on other critical illnesses or including other treatments; and studies focusing solely on intervention programs.

Description of the studies

Eighteen studies were selected for this review. Table 1 contains a descriptive summary of the articles. We included the aim of the study, number and characteristics of parents and children, methodological features (e.g. study design, timing of measurement); measures; and main results. The indications for SCT varied among the different studies. Most studies reported the underlying diagnosis, type of transplant and type of donor, transplant risk, and disease risk category [17]. On average 80% of the children undergoing SCT suffered from a malignant disease and around 60% of the children in the studies underwent allogeneic SCT (transplantation with bone marrow from a foreign donor) as opposed to autologous SCT (transplantation with own body material).

Table 1 Summary of the studies included in the review

No.	Year, author	Aim	N parents, N and age range/ mean age children	Methodological features	Measures	Results
1	1990, Dermatis	Determine the nature and prevalence of the psychological symptomatology in parents of children undergoing SCT. Investigate relationship of certain psychosocial factors to parental distress associated with the informed consent process. Examine the stress responses of mothers during their child's hospitalization for SCT. Determine the relationships between mothers' stress responses and the resources for coping and social support.	46 M, 15 F 61 children, range 1–17 yrs	Single centered Cross-sectional	BSI, WOC, newly constructed scale on the quality of the physician–parent communication	47% of fathers and 60% of mothers exhibited significant psychological distress of a generalized nature. Mothers reported more severe levels of depression and anxiety than fathers did.
2	1997, Nelson	Examine the stress responses of mothers during their child's hospitalization for SCT.	50 M	Single-centered	STAI, CES-D, HAS, IES, SSS	Maternal anxiety and depressive scores decreased significantly over time
3	2000, Streisand	Document levels of stress in mothers of children undergoing SCT. Pilot a psychological intervention program.	50 children, mean age 9.3 yrs 11 M	Longitudinal, time points: time of admission (T1), second (T2), tenth (T3), and twentieth (T4) day post-SCT Single-centered	DSI, PSI, SSINT	The coping style defined as 'active reviewing of feelings or information associated with the situation' significantly explained variance in scores for anxiety, depressive symptomatology, somatic complaints and sleep behavior. Most stress was reported pre-admission.
4	2001, Manne	Examine anxiety and depressive symptoms among mothers of children undergoing SCT.	11 children, range 2–16 yrs, mean age 8.8 yrs 115 M	Longitudinal, time points: pre-admission to 3 weeks post-SCT Multi-centered	BAI, BDI, SCID-NP	Mothers reported using more stress management techniques post-intervention than mothers in the standard care condition. The analyses revealed no significant differences in stress between intervention and control mothers. 20% of mothers were diagnosed with a MDD, a GAD, or a PD. There was evidence of comorbidity between anxiety and depressive disorders. Mothers with lower incomes, who were Caucasian, had received prior psychiatric care and were caring for female SCT patients may be at higher risk for adverse psychological reactions.
5	2002, Manne	Investigate the role of cognitive and social processing in post-traumatic stress symptoms and disorder (PTSD) among mothers of children undergoing SCT.	115 children, range 4 months–20 yrs, mean age 9.2 yrs 90 M	Cross-sectional; 85% of mothers on day–7 to day –1; 15% of the mothers 10-days post-SCT Multi-centered	SCID-NP-PTSD, PCL-C, BAI, BDI, fear network, CSI, LSCM	Emotional distress, SCT-related fears, and negative responses of family and friends assessed at the time of SCT hospitalization were predictive of later PTSD symptoms. Cognitive processing (the appraisal of threat) at the time of transplantation played the most important role in later PTSD symptoms. Parents expressed intense distress and disorientation and sometimes difficult relations with their child. Many parents expressed having an ambivalent relation with care providers.
6	2002, Oppenheim	Understand parents' perception of children treated in an SCT unit.	90 children, range 9 months–20 yrs, mean age 8.8 yrs 40 pairs of parents No details given	Longitudinal, time points: time of SCT, 3 and 6 months past SCT Single-centered Cross-sectional	Interviews	Acceptance and humor were associated with reductions in maternal depressive symptoms. Planning and alcohol/substance use were associated with increases in maternal depressive symptoms. Active problem solving and use of instrumental support did not predict changes in depressive symptoms.
7	2003a, Manne	Evaluate the role of maternal coping strategies in depressive symptoms experienced by mothers of children undergoing SCT.	207 M 207 children, mean age 8.3 yrs	Multi-centered Longitudinal, time points: at SCT, 3 and 6 months post-SCT	COPE, BDI, appraisal of fear/worry, medical risk	Fear structure, distress, and unsupportive responses by
8	2003b, Manne	Examine the role of perceived	148 M	Multi-centered	SCID-NP-PTSD, PCL-C,	

Table 1 (continued)

No.	Year, author	Aim	N parents, N and age range/ mean age children	Methodological features	Measures	Results
9	2003, Nelson	partner criticism and avoidance in anxiety and depressive symptoms of mothers of children undergoing SCT.	148 children, range 4 months–17 yrs, mean age 8.5 yrs	Longitudinal, time points: at SCT, 3 and 6 months post-SCT	BAI, BDI, fear network, CSI, CSI, LSCM	family and friends measured at transplantation were predictive of PTSD symptom severity at 6 months after SCT. Partner criticism was associated with higher average depressive symptoms. ICU transfers and number of days of hospitalization 6 months post-SCT were risk factors.
10	2004, DuHamel	Examine the relationships between maternal anxiety and depressive symptoms and resources during their child's SCT.	23 M 23 children, mean age 8.1 yrs	Single-centered longitudinal, time points: admission and 10 days post-SCT	STAI, CES-D, SPSS, SSS	The majority of mothers reported moderate to high anxiety levels and were at risk of developing depression. Most of the mothers indicated low or moderate satisfaction with the perceived social support. A relationship was found between a negative problem solving orientation and emotional responses.
11	2004, Forinder	Investigate the role of cognitive processing in maternal adjustment to a life-threatening pediatric medical procedure.	91 M 91 children, range 9–19 yrs, mean age 8.7 yrs	Multi-centered Longitudinal, time points: 3 days prior to SCT and 3 months post-SCT	Structured interviews Fear network IES, BAI, BDI	Mothers' fear network, intrusions and avoidance played a primary role in their adjustment to their child's transplantation, during and after hospitalization. The article shows a cognitive processing model of psychological distress.
12	2004, Manne	To get in-depth knowledge of the parents' situation during the SCT-process.	20 pairs of parents 20 children, no details given	Single-centered Longitudinal, time points: 4 to 8 yrs post-transplant and 4 yrs after first time point	2 semi-structured qualitative interviews Jalowiec Coping Scale	The child's illness and treatment played an important role in the parents' lives for many years. Those parents who managed to put reason before emotion rated their coping as better. A sense of participation was also a useful coping strategy.
13	2004, Phipps	Examine the prevalence and predictors of anxiety, depression and PTSD among mothers of children who underwent HSCT.	111 M 111 children, range 1–18, mean age 8.2 yrs	Multi-centered Longitudinal, time points: at time of SCT and 18 months post-SCT	BAI, BDI, TSS, ISSB, WOC, COPE 18 month follow up: SCID-NP	Approximately 20% of mothers had clinically significant distress reactions. Mothers who were most at risk were younger and reported anxiety and depressive symptoms at the time of transplantation. The prevalence of depressive disorders declined after 18 months.
14	2004a, Rini	Examine changes in parental distress across the acute phase of SCT.	136 M, 9 F, 6 others 136 children, range 1–20 yrs, mean age 8.9 yrs	Single-centered Longitudinal, 13 time points: weekly from week -1 to week+6, after that on a monthly basis through month+6	POMS, PSS, CBS, BASES-P, BASES-C	Parents demonstrated modest, but significant elevations in distress, particularly during the early period from admission through week +3. Parental distress was unrelated to child age, gender, diagnosis, or type of transplant, but was significantly related to parental SES. Moderate correlations were observed between measures of parent and child distress.
15	2004b, Rini	Examine the relationship of parental distress to child distress during the SCT process.	100 M Range 9 months–20 yrs, mean age 8 yrs	Multi-centered Longitudinal, time points: at admission and 1 yr post-SCT	WAS, TSS, LES, SF36	Prior trauma and negative events were associated with basic beliefs during hospitalization and with changes in basic beliefs in the subsequent year, with distress mediating some of these relations. Relations were found between basic beliefs and maternal physical and mental functioning.
16	2004b, Rini	Examination of children's medical risk and mother's dispositional risk	144 M 144 children, range 9 months–	Multi-centered Longitudinal, time points:	LOF, SF-36 (MHSS), 2 newly created items for	Predictors of benefit finding differed systematically across assessments, with optimism and medical risk

	optimism and socio-demographic resources as predictors of benefit finding at admission (T1) and 6 months later (T2).	20 yrs, mean age 8 yrs	at admission and 6 months after the first time point	benefit finding	predicting benefit finding at both time points. Socio-demographic resources predicted only T2 benefit finding. T1 benefit finding was positively associated with T2 adaptation only for mothers who scored high in optimism.
16	2005, Phipps	139 M, 9 F, 3 others	Single-centered	POMS, PSS, CBS, PIES, CBCL, FES, ISSB, WOC	Significant changes were observed in parental distress across the course of SCT, with relatively high levels of parental distress at admission, slightly increasing and peaking at week+2.
17	2007, DuHamel	151 children, range 1–20 yrs, mean age 8.9 yrs	Longitudinal, time points: weekly basis through week+6 post-SCT, monthly until +6	LOT, LES, TSS, newly created items for fear appraisals	Predictors of stress: prior parent and patient illness-related distress, pre-morbid child internalizing behavior problems, the family relationship dimensions of the family environment and parental avoidant coping behaviors. Lower optimism and a greater number of negative life events were independently associated with greater maternal fear appraisals. Lifetime history of trauma was not associated with maternal fear appraisals. Mothers' fear appraisals during their child's hospitalization were associated with their fear appraisals up to 6 months later.
18	2008, Vrijmoet-Wiersma	19 M	Single-centered	PSI	Compared to parents of healthy children, parenting stress was higher post-SCT. Post-SCT, parenting stress levels were higher than pre-SCT, both total parenting stress and the perceived demandingness of the child. High levels of parenting stress were predictive of poor parental ratings of child HRQoL post-SCT.

Abbreviations: M mothers, F fathers, yrs years, BAI Beck Anxiety Inventory, BASES-P/C Behavioral, Affective, and Somatic Experiences Scales—Parent Version/Child Version, BDI Beck Depression Inventory, BSI Brief Symptom Inventory, CBCL Child Behavior Check List, CBS Caregiver Burden Scale, CES-D Center of Epidemiologic Studies Depression Scale, CSI Cancer Support Inventory, DSI Daily Stress Inventory, Faces III Family Adaptability and Cohesion Evaluation Scale, FES Family Environment Scale, HAS Health Assessment Scale, IES Impact of Events Scale, ISSB Inventory for Socially Supportive Behaviors, LES Life Experiences Survey, LOF Life Orientation Test, LSCM Lepore's Social Constraints Measure, MHSS-SF36 Mental Health Summary Scale of the Short Form-36, PCL-C Post-traumatic Symptom Disorder Checklist-Civilian version, PIES Prior Illness Experience Scale, POQOLS Pediatric Oncology Quality of Life Scale, POMS Profile of Mood States, PSI Parenting Stress Index, PSS Perceived Stress Scale, SCID-NP Structured Clinical Interview for DSM-IV, Non-Patient version, SSINT Semi-structured Interview, SSS Stress Support Scale, STAI State-Trait Anxiety Inventory, TSS Traumatic Stress Schedule, VABS Vineland Adaptive Behavior Scales, WAS World Assumptions Scale, WOC Ways of Coping Checklist

The studies in our review did not distinguish between the experiences of parents of children undergoing SCT once and parents of children who had multiple SCT experiences, who will undoubtedly have to face a unique set of stressors.

In this review article, we will first discuss the methodological qualities of the studies, next we will summarize the main results of the studies, and lastly we will discuss risk and protective factors of parental adaptation to their child's SCT.

Methodological qualities of the selected studies

Study design and timing of assessment

Three of the included studies used a cross-sectional study design [3, 20, 27] and one study was descriptive/retrospective [8]. The other 14 studies used a prospective longitudinal design with repeated measures, ranging from two to 13 measurement points. However, only a few of these studies followed a particular aspect of parental distress over time, e.g. [18, 32]. Exactly half of the studies included in the review used a multi-centered design, the other nine studies recruited participants from one medical center. Until now, only one intervention study has been published in this area. It included mothers of children undergoing SCT [36] and was based on a stress inoculation model.

In most longitudinal studies, two measurement points were used. The first time point was between 47 to 1 day(s) pre-admission and a few days post-SCT. The time point for the second assessment varied strongly between the studies, ranging from 1 week post-SCT to 18 months post-SCT. Phipps et al. [31, 32] used up to 13 time points in both studies. Only one of the studies assessed long-term parental distress, 4 to 8 years after stem cell transplantation [8].

Participants

The majority of the studies (13 in total) used only mothers as respondents. Sample size in the studies varied from 11 [36] to 207 parents [18]. Eleven of the studies included more than 90 parents. All studies described the recruitment process. Phipps et al. [31, 32] assessed one of the caregivers, resulting in 90% mothers. In only three studies [3, 27, 39], both parents were used as respondents. The age range of the children was most often 1 to 20 years of age with a mean age at first assessment between 8 and 9 years. In two studies, the age of the children was not mentioned [8, 27].

Outcome measures

The conceptualization of stress or distress varied widely between the studies. The distinction between the assess-

ment of (subclinical) levels of distress on the one hand and clinical psychiatric diagnoses was not always clearly made, which makes comparisons difficult. Anxiety and depression were studied as manifestations of parental distress in nearly all of the studies. Other manifestations of parental distress or psychiatric disorders were disturbed and obsessive-compulsive thinking [3], post-traumatic stress symptoms [16, 17], generalized anxiety disorder (GAD), panic disorder (PD), and major depressive disorder (MDD) [20]. Somatic complaints and changes in sleep behavior were added by some researchers [25, 34, 35] as symptoms of parental distress. Variables influencing parental stress levels were mostly operationalized as 'coping' [18, 24, 25, 32], 'family functioning' [32], 'parenting stress issues' [39], and 'social support', both positively and negatively perceived [19, 25, 32].

In most studies multiple measures were used, most often self-report questionnaires combined with, or additional to, interviews as a way of collecting data. In the majority of studies standardized questionnaires were used to assess parental distress reactions (e.g. Beck Depression Inventory (BDI), Beck Anxiety Inventory (BAI), State-Trait Anxiety Inventory (STAI), and Parental Stress Index (PSI)), e.g. [17, 20]. Disease- or context-specific measures were used less frequently. Phipps et al. [31, 32] developed and used the Prior Illness Experiences Scale (PIES) to assess previous parent and child experiences with cancer therapy and inpatient hospitalization. Rini et al. [35] created two items to measure benefit finding in their study and DuHamel and her study group developed a scale to assess maternal fear appraisals [6].

Results

Parental stress

Feelings of anxiety and depression, post-traumatic stress symptoms, disturbed or obsessive compulsive thinking, and somatic complaints are the most common stress reactions parents report before, during, and after SCT. Parental stress levels were reported to be higher than norm groups pre-transplant [3, 32, 36] and during the acute phase of the SCT, peaking at 2–3 weeks post-SCT [32]. The most common observation was that parental distress levels decreased over time after SCT [5, 36], with the sharpest declines between 3–6 months post-transplant [28]. A recent study, however, showed that parents, on average 10 months post-SCT, reported higher parenting stress levels and, specifically, felt less competent as a parent [39]. In the longer term, 4 to 8 years post-SCT [8], many parents reported that their child's illness and subsequent treatment played an important role in their lives for years, ranging from parents still struggling on a

daily basis to parents who put this ordeal more or less behind them. Studies comparing mothers and fathers are few in this area, since most studies included mothers only. Dermatis and Lesko [3] found higher levels of depression and phobic anxiety in mothers than in fathers.

Table 1 depicts the results of the 18 studies included in the review.

Risk and protective factors

Many factors have been identified to influence levels of parental distress. The most frequently identified risk factor for parental distress in the longer term is the manner in which the parent is able to handle stress during the acute phase. We grouped the risk factors into three clusters, based on a manual count of the determinants described in the included studies:

1. Disease factors, i.e. if the child had been transferred to the ICU and if it had had a higher number of hospitalizations 6 months post-SCT, parents reported more anxious and depressive symptoms [19]. Higher transplant risk was also associated with higher parental distress during the child's post hospital stage of recovery [6].
2. Psychological factors and parental coping: mother's appraisal of threat to her child's life [5, 16], a greater number of negative life events [6], prior parent and patient experiences of distress associated with the child's illness [32], avoidance and intrusions [18, 32], alcohol/substance abuse [18], perceived partner criticism [19], and an unsupportive family environment [32] all added to parental (i.e. mostly maternal) stress levels. Furthermore, mothers experiencing depressive symptoms during the acute SCT phase had a higher probability to be diagnosed with a psychiatric disorder 18 months post-SCT [17]. Parent distress has also been associated with child distress: child mood disturbance at admission was predictive of parent global distress over time [28].
3. Socio-demographic risk factors: younger maternal age [17] and lower social economical status (SES) [31] were associated with higher levels of stress throughout the SCT process.

Factors that did not appear to influence parental stress were: the age of the child, type of diagnosis, type of transplant, or the nature of the germ-free environment in which the child was placed [3, 5, 32]. Other objective medical aspects of the child's condition (i.e. disease risk, treatment course, and current disease status) than the factors mentioned under the first cluster (see above) were not related to parental stress levels or fear appraisals. It seems that the subjective appraisal of these factors is a better

predictor of parental stress than the objective disease characteristics [6].

Protective factors or predictors of good psychological adaptation have been identified in terms of 'benefit finding' [35]: mothers who were optimistic by nature reported most benefit finding both during the acute SCT phase and 6 months later. Benefit finding is defined as 'an attempt to restore positive basic beliefs about the self and the world that have been challenged by a traumatic experience' [35, 37]. Acceptance and humor as coping mechanisms were associated with reductions in maternal depressive symptoms [18] and 'putting reason before emotion' was identified as another coping mechanism associated with positive outcomes [8]. Lastly, a supportive family environment was associated with lower distress levels throughout the transplant process [32].

Discussion

Having a child undergo stem cell transplantation is a stressful event for any parent. Feelings of anxiety and depression, post-traumatic stress symptoms, disturbed or obsessive compulsive thinking, and somatic complaints are the most common stress reactions parents report before, during, and after SCT. The process of SCT is comprised of several phases and distress levels seem most elevated in the pre-SCT phase and the acute phase during hospitalization, but can stay elevated after discharge. Most parents return to healthy levels of psychosocial functioning 18 months post-SCT, but a subset of parents reports persistent symptoms years later in terms of anxiety, depressive feelings, and post-traumatic stress symptoms (arousal, avoidance, and recurring memories). Certain maternal coping strategies (e. g. acceptance, humor, putting reason before emotion, and having positive cognitive appraisals) during the acute phase of SCT have been identified as protective factors.

It has been shown that dispositional factors and prior experiences influence the way an individual appraises an event such as SCT and adjusts to it. For example, mothers who were more optimistic by nature reported lower fear appraisals at the time of their child's hospitalization for SCT [6] and later on, post-SCT. Optimism seems to be a more or less stable trait that can serve as a coping mechanism and a buffer [10], like positive reframing [18] and benefit finding [35]. Mothers who have encountered more negative life events in the months before their child's SCT reported more fear appraisals. It is hypothesized that traumatic events can trigger increased arousal, cognitions that one's life is difficult and traumatic [17], and a tendency to interpret potentially harmful new events more negatively [6]. Pre-existing factors should be assessed and used as starting points for psychosocial interventions.

The present review reveals potential areas of improvement in future research. In the 18 studies included in this review a variety of definitions of the core elements of the psychological stress process have been used, often described together and simply referred to as ‘stress’. It is important to clarify what is meant by ‘stress’ and to specify both the temporal course of a stressor [13] and to identify SCT-specific stressors. Post-traumatic stress reactions imply an existential challenge, but findings suggest that the complex situation of SCT involves several different stressors for the parents. To facilitate communication and collaboration it is necessary to be more specific in the terminology used to describe the psychological reactions of both parents and patients and to make a clear distinction between stress as a predictor variable and psychological stress as an outcome.

In most of the research in this area, no distinction was made between subclinical manifestations of parental distress versus psychiatric states. This is unfortunate, because in the latter approach, parents tend to be ‘pathologized’ [33] instead of assuming that the majority of families with a seriously ill child are competent and adaptively organized families, without any elevations in their a priori risk (as a group) for psychopathology [12]. Furthermore, it seems that in multidisciplinary SCT teams often there is no consensus of what is ‘normal’ distress or ‘adequate coping’ in this context. For example, young and inexperienced nurses can get worried about a parent in tears whereas an experienced social worker or psychologist may feel that a certain level or manifestation of distress is ‘normal’. This issue points to the need for adequate psychosocial screening by pediatric psychologists pre-admission and during the acute phase of SCT, in order to target those parents most in need for psychosocial guidance and intervention.

Family functioning, an area of increasing importance in the pediatric psychology literature, is still understudied in parents of children undergoing SCT. The experience of fathers is another area of neglect. In many studies on parental stress of parents of pediatric cancer patients, higher stress levels have been found for mothers than for fathers [40], but recent research has shown that the experience of the child’s illness often is as stressful for fathers as for mothers [26]. This finding points to the need to include fathers in future studies.

Strengths of the studies included in our review are the large number of longitudinal designs and multi-centered studies and the majority of studies with 90 participants or more. We have found only a minority of studies in which disease- or context-specific measures were used and this is unfortunate, because SCT is a highly complex treatment with very specific issues to deal with for parents. A combination of generic and disease-specific instruments could further our understanding of parental distress trajectories during the course of the SCT.

Conclusions

The authors conclude that the majority of parents of pediatric SCT patients are resilient, 18 months post-SCT and beyond. The most frequently identified risk factor for parental distress in the longer term is the manner in which the parent is able to handle stress during the acute phase. Parents (mostly mothers) with the most severe stress reactions and fear appraisals during the acute phase, continue to experience heightened levels of anxiety, depressive symptoms, and PTSS later on.

The next step is to develop and systematically examine feasible, limited, brief interventions for sub-clinical manifestations of psychological distress prior to and during the acute phase of SCT in parents who have been identified as ‘risk’ group. Follow-up care is needed for parents, especially when their child recovers and when control visits to the hospital become less frequent. Intervention research is a growing area in medical psychology and despite the many methodological challenges, efforts should be made to implement and evaluate existing intervention programs in this parent group. This can only be done through sound SCT-specific-assessment, well-funded (inter) national cooperation, and well-developed study designs. Lastly, the ethical domain of conceiving designed children as donors is an area of interest that deserves to be studied in the future, as well as the issue of stress in parents of children who need to undergo more than one stem cell transplantation or whose children suffer from more serious late effects, such as chronic graft-versus-host disease or other health problems.

Acknowledgements We would like to thank our psychology master student Saskia de Ridder for her contribution to the literature search and for setting up the table and we owe Robbert Bredius thanks for his important comments on the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Reference

1. Barrera M, Boyd-Pringle LA, Sumbler K, Saunders F (2000) Quality of life and behavioral adjustment after pediatric bone marrow transplantation. *Bone Marrow Transplant* 26(4):427–435
2. Clarke SA, Eiser C, Skinner R (2008) Health-related quality of life in survivors of BMT for paediatric malignancy: a systematic review of the literature. *Bone Marrow Transplant* 42:73–80
3. Dermatis H, Lesko LM (1990) Psychological distress in parents consenting to child’s bone marrow transplantation. *Bone Marrow Transplant* 6(6):411–417

4. Duell T, van Lint MT, Ljungman P, Tichelli A, Socie G, Apperley JF et al (1997) Health and functional status of long-term survivors of bone marrow transplantation. *Ann Intern Med* 126 (3):184–192
5. DuHamel KN, Manne S, Nereo N, Ostroff J, Martini R, Parsons S et al (2004) Cognitive processing among mothers of children undergoing bone marrow/stem cell transplantation. *Psychosom Med* 66(1):92–103
6. DuHamel KN, Rini C, Austin J, Ostroff J, Parsons S, Martini R et al (2007) Optimism and life events as predictors of fear appraisals in mothers of children undergoing hematopoietic stem cell transplantation. *Psychooncology* 16(9):821–833
7. Felder-Puig R, di Gallo A, Waldenmair M, Norden P, Winter A, Gadner H et al (2006) Health-related quality of life of pediatric patients receiving allogeneic stem cell or bone marrow transplantation: results of a longitudinal, multi-center study. *Bone Marrow Transplant* 38(2):119–126
8. Forinder U (2004) Bone marrow transplantation from a parental perspective. *J Child Health Care* 8(2):134–148
9. Grootenhuis MA, Last BF (1997) Adjustment and coping by parents of children with cancer: a review of the literature. *Support Care Cancer* 5(6):466–484
10. Grootenhuis MA, Last BF (1997) Predictors of parental emotional adjustment to childhood cancer. *Psychooncology* 6(2):115–128
11. Hoodin F, Weber S (2003) A systematic review of psychosocial factors affecting survival after bone marrow transplantation. *Psychosomatics* 44(3):181–195
12. Kazak AE, Rourke MT, Alderfer MA, Pai A, Reilly AF, Meadows AT (2007) Evidence-based assessment, intervention and psychosocial care in pediatric oncology: a blueprint for comprehensive services across treatment. *J Pediatr Psychol* 32 (9):1099–1110
13. La Greca AM, Lemanek KL (1996) Assessment as a process in pediatric psychology. *J Pediatr Psychol* 21(2):137–151
14. Lesko LM (1994) Bone marrow transplantation: support of the patient and his/her family. *Support Care Cancer* 2(1):35–49
15. Löf C, Winiarski J, Giesecke A, Ljungman P, Forinder U (2009) Health-related quality of life in adult survivors after paediatric allo-SCT. *Bone Marrow Transplant* 43(6):461–468
16. Manne S, DuHamel K, Nereo N, Ostroff J, Parsons S, Martini R et al (2002) Predictors of PTSD in mothers of children undergoing bone marrow transplantation: the role of cognitive and social processes. *J Pediatr Psychol* 27(7):607–617
17. Manne S, DuHamel K, Ostroff J, Parsons S, Martini DR, Williams SE et al (2004) Anxiety, depressive, and posttraumatic stress disorders among mothers of pediatric survivors of hematopoietic stem cell transplantation. *Pediatrics* 113(6):1700–1708
18. Manne S, DuHamel K, Ostroff J, Parsons S, Martini DR, Williams SE et al (2003) Coping and the course of mother's depressive symptoms during and after pediatric bone marrow transplantation. *J Am Acad Child Adolesc Psych* 42(9):1055–1068
19. Manne S, DuHamel K, Winkel G, Ostroff J, Parsons S, Martini R et al (2003) Perceived partner critical and avoidant behaviors as predictors of anxious and depressive symptoms among mothers of children undergoing hemopoietic stem cell transplantation. *J Consult Clin Psychol* 71(6):1076–1083
20. Manne S, Nereo N, DuHamel K, Ostroff J, Parsons S, Martini DR et al (2001) Anxiety and depression in mothers of children undergoing bone marrow transplant: symptom prevalence and use of the Beck depression and Beck anxiety inventories as screening instruments. *J Consult Clin Psychol* 69(9):1037–1047
21. Matthes-Martin SLM, Ladenstein R, Emminger W, Felsberger C, Topf R, Gadner H et al (1999) Organ toxicity and quality of life after allogeneic bone marrow transplantation in pediatric patients: a single center retrospective analysis. *Bone Marrow Transplant* 23 (10):1049–1053
22. Mosher CE, Redd W, Rini C, Burkhalter JE, DuHamel KN (2009) Physical, psychological, and social sequelae following hematopoietic stem cell transplantation: a review of the literature. *Psychooncology* 18:113–127
23. Neitzert CS, Ritvo P, Dancey J, Weiser K, Murray C, Avery J (1998) The psychosocial impact of bone marrow transplantation: a review of the literature. *Bone Marrow Transplant* 22(5):409–422
24. Nelson AE, Gleaves L, Nuss S (2003) Mothers' responses during the child's stem cell transplantation: pilot study. *Pediatr Nurs* 29 (3):219–223
25. Nelson AE, Miles MS, Belyea MJ (1997) Coping and support effects on mothers' stress responses to their child's hematopoietic stem cell transplantation. *J Pediatr Oncol Nurs* 14(4):202–212
26. Norberg AL, Boman K (2008) Parent distress in childhood cancer: a comparative evaluation of posttraumatic stress symptoms, depression and anxiety. *Acta Oncol* 47(2):267–274
27. Oppenheim D, Vasselon S, Hartmann O (2002) How do parents perceive high-dose chemotherapy and autologous stem cell transplantation for their children. *Bone Marrow Transplant* 30(1):35–39
28. Phipps S (2006) Psychosocial and behavioral issues in stem cell transplantation. In: Brown RT (ed) *Comprehensive handbook of childhood cancer and sickle cell disease, a biopsychosocial approach*. Oxford University Press, New York
29. Phipps S, Dunavant M, Garvie PA, Lensing S, Rai SN (2002) Acute health-related quality of life in children undergoing stem cell transplant: I. Descriptive outcomes. *Bone Marrow Transplant* 29(5):425–434
30. Phipps S, Dunavant M, Lensing S, Rai SN (2002) Acute health-related quality of life in children undergoing stem cell transplant: II. Medical and demographic determinants. *Bone Marrow Transplant* 29(5):435–442
31. Phipps S, Dunavant M, Lensing S, Rai SN (2004) Patterns of distress in parents of children undergoing stem cell transplantation. *Pediatric Blood & Cancer* 43(3):267–274
32. Phipps S, Dunavant M, Lensing S, Rai SN (2005) Psychosocial predictors of distress in parents of children undergoing stem cell or bone marrow transplantation. *J Pediatr Psychol* 30(2):139–153
33. Quittner AL, Davis MA, Modi AC (2003) Health-related quality of life in pediatric populations. In: Roberts MC (ed) *Handbook of pediatric psychology*, 3rd edn. Guilford, New York, pp 696–709
34. Rini C, Manne S, DuHamel KN, Austin J, Ostroff J, Boulard F et al (2004) Changes in mothers' basic beliefs following a child's bone marrow transplantation: the role of prior trauma and negative life events. *J Trauma Stress* 17(4):325–333
35. Rini C, Manne S, DuHamel KN, Austin J, Ostroff J, Boulard F et al (2004) Mothers' perceptions of benefit following pediatric stem cell transplantation: a longitudinal investigation of the roles of optimism, medical risk, and sociodemographic resources. *Ann Behav Med* 28(2):132–141
36. Streisand R, Rodrigue JR, Houck C, Graham-Pole J, Berlant N (2000) Brief report: parents of children undergoing bone marrow transplantation: documenting stress and piloting a psychological intervention program. *J Pediatr Psychol* 25(5):331–337
37. Taylor SE (1983) Adjustment to threatening events: a theory of cognitive adaptation. *Am Psychol* 38:1161–1173
38. Tsimicalis A, Stinson J, Stevens B (2005) Quality of life of children following bone marrow transplantation: critical review of the research literature. *Eur J Oncol Nurs* 9(3):218–238
39. Vrijmoet-Wiersma CMJ, Kolk AM, Grootenhuis MA, Spek EM, van Klink JMM, Egeler RM et al (2009) Child and parental adaptation to pediatric stem cell transplantation. *Support Care Cancer* 17(6):707–714
40. Vrijmoet-Wiersma CMJ, van Klink JMM, Kolk AM, Koopman HM, Ball LM, Maarten Egeler R (2008) Assessment of parental psychological stress in pediatric cancer: a review. *J Pediatr Psychol* 33(7):694–706