



OPEN

# Pomegranate fruit juice adulteration with apple juice: detection by UV–visible spectroscopy combined with multivariate statistical analysis

Lucia Pappalardo

Pomegranate is rich in high value nutritional substances known to be beneficial against several diseases and its use in medicine is known since ancient times. Due to its properties and delicious taste, pomegranate fresh fruit juices demand has been growing worldwide and its adulteration is becoming a problem. Low-cost, user friendly and fast detection methods are therefore desirable in order to easily and rapidly detect adulteration of short shelf-life fresh fruit juices. For this purpose fresh squeezed arils pomegranate juice samples adulterated with less expensive apple juice concentrate were investigated by UV–visible spectroscopy combined with multivariate statistical analysis. Unsupervised principle component analysis (PCA), supervised projection to latent structure discriminant analysis (PLS-DA) and orthogonal projection to latent structure discriminant analysis (OPLS-DA) were performed on the full spectra. OPLS-DA analysis of UV–visible spectra proved to be a suitable method to detect pomegranate juices adulterated by more than 20% v/v apple juice concentrate.

Pomegranate has been widely cultivated in the Mediterranean since ancient times and nowadays it is widely cultivated also in Middle East and Asia. Pomegranate fresh fruit juice demand is increasing due to its health benefits and it is very popular among consumers concerned about healthy diet and lifestyle. The oldest report about medical applications of pomegranate dates back to 1500 B.C. that mentioned its use in Ancient Egypt to treat several infections<sup>1</sup>. Recent studies have reported pomegranate's antiviral, antimicrobial, antioxidative and anti-inflammatory properties and potential beneficial applications against several diseases<sup>2</sup>. As the demand of fresh fruit juices with high nutritional values is increasing worldwide so their adulteration is becoming a growing problem. Therefore, low-cost, user friendly and fast adulteration detection methods are desirable in order to control easily and rapidly the quality of the short shelf-life fresh fruit juices.

Studies based on mass spectrometry (MS) combined with multivariate statistical analysis for determining the level of adulteration in fruit juices<sup>3–5</sup> and in particular in pomegranate fruit juices<sup>6</sup> have been performed. <sup>1</sup>H nuclear magnetic resonance (NMR) analyses combined with chemometrics have been applied to determine the percentage of pure fruit juices in blends of apple, orange, pineapple and pomegranate juices<sup>7</sup> and adulterations in grape nectars<sup>8</sup>. A method based on Fourier transform infrared (FTIR) spectroscopy combined with unsupervised principal component analysis (PCA) was able to determine adulteration in pomegranate juices concentrate with grape juice concentrate<sup>9</sup>. Furthermore a comprehensive study by UV–Vis spectroscopy and PCA analysis on pomegranate juices mixed with grape and apple juices or simply diluted with water has been published<sup>10</sup>. Boggia et al. concluded that pomegranate adulterated with these fillers can be detected by UV–Vis spectroscopy and unsupervised PCA analysis. However, they determined that this method was not particularly distinctive for pomegranate and apple juices mixtures.

The aim of this work was to expand on the findings of Boggia et al. and investigate how well pomegranate juices adulterated by apple juices can be detected by applying supervised multivariate statistical discriminant

Department of Biology, Chemistry and Environmental Sciences, American University of Sharjah, PO Box 26666, Sharjah, United Arab Emirates. email: [Lpappalardo@aus.edu](mailto:Lpappalardo@aus.edu)

analysis (PLS-DA and OPLS-DA) to UV-Vis spectra of fruit juices and their mixtures. For this purpose, fresh squeezed arils pomegranate juice samples adulterated with less expensive apple juice concentrate were investigated by UV-visible spectroscopy combined with supervised and unsupervised multivariate statistical analysis.

## Material and methods

**Samples.** Fruit juices were purchased at local grocery stores. Fresh squeezed arils pomegranate juices ( $n = 56$ ), and apple juices prepared from concentrate ( $n = 56$ ) were used. The pomegranate and apple juices were purchased, respectively, from two different companies. For each of these juice samples, mixtures in their native fresh conditions at 10%, 20%, 30% and 40% v/v apple/pomegranate juices were prepared by mixing in a vial. Mixtures and pure juices were further diluted 20 times with Milli-Q water (Merck Millipore, Burlington, MA, USA) and 5 ml of each sample were centrifuged at 4000 rpm for 10 min prior to analysis. The dilution and centrifugation were required in order to be able to detect the UV-Vis signal in the proper range of the instrument.

**Instrumentation.** Spectra were acquired on a Shimadzu UV-1800 spectrometer equipped with the software UVProbe ver. 2.43 (Shimadzu, Kyoto, Japan). The temperature of the samples was controlled by a TC1 temperature controller and T-App software ver. 1.30 (Quantum Northwest Inc., Liberty Lake, WA, USA).

**Data acquisition.** UV-Vis absorption spectra were acquired in the 200–800 nm range with 1 nm resolution. Spectra for each sample were recorded at 10 different temperatures in the range 25–80 °C. Two spectra were recorded for each sample at each temperature and their average values were employed for further analysis. Quartz cuvettes of 1 cm path length were used. Spectra showed stability over the temperature range investigated and only data recorded at 25 °C are reported here.

**Multivariate statistical analysis.** The UV-Vis spectra were exported to MS Excel software. The full spectra range from 200 to 800 nm was used for the analysis and the size of each spectrum was 601 data points. Unsupervised principle component analysis (PCA) was performed initially, followed by supervised projection to latent structure discriminant analysis (PLS-DA) and orthogonal projection to latent structure discriminant analysis (OPLS-DA) using the SIMCA software (v.13.5, Umetrics, Umeå, Sweden)<sup>11</sup>. OPLS-DA uses an orthogonal filter to remove all the information in X that are not correlated with Y and is able to augment the classification performance of PLS-DA<sup>12</sup>.


Univariate (UV) scaling was used for all of the variables after mean centering<sup>13</sup>. Pareto scaling and mean centering without scaling were also investigated on the data set under analysis and UV scaling provided the best results. Validation was accomplished using the routine in SIMCA taking 7th-order Monte Carlo cross-validation<sup>11,14</sup>. Goodness of fit of the data to the model [ $R^2$  (cum)] and predictability of the model [ $Q^2$  (cum)] were calculated for the scores plots and variables of importance determined by comparing coefficient plots which resemble the real spectral features<sup>15</sup>.

## Results and discussion

Pomegranate juices have high nutritional values and are sometimes adulterated by counterfeiters by the addition of less expensive juices as for example apple juices. While sophisticated techniques as MS<sup>3–6</sup>, NMR<sup>7,8</sup>, FTIR<sup>9</sup> have proved to be effective in detecting the adulterations in fruit juices, it is desirable to have a faster and less expensive detection method as for example UV-Vis spectroscopy. For this reason, fresh squeezed pomegranate juices, apple juices prepared from concentrate and their mixtures, in their native conditions, were investigated by UV-Vis spectroscopy and full-resolution spectra were analyzed by multivariate statistics in this study.

First goal was to investigate if the method was able to differentiate between apple and pomegranate juices. For this scope the UV-Vis spectra of 100% pomegranate and 100% apple juices samples were recorded and then analyzed by multivariate methods. Unsupervised PCA was calculated and the PCA scores plot showed good clustering of the two classes (Fig. 1a). Then the two classes, apple (A) and pomegranate (P), were specified for the discriminant analysis (PLS-DA and OPLS-DA). The corresponding PLS-DA (Fig. 1b) and OPLS-DA (Fig. 1c) scores plots show a clear separation, quite pronounced in the OPLS results. All observations fell within or close to the 95% Hotelling ellipse, which indicates a coherent dataset (Fig. 1). The analysis also showed excellent cluster correlation [ $R^2$  (cum) = 0.99] and predictability [ $Q^2$  (cum) = 0.99] with good validation results. This first set of experiments proved that UV-spectroscopy combined with multivariate statistical analysis was able to differentiate between apple and pomegranate juices.

In order to determine if pomegranate juice adulterated by apple juice can be detected by this method, the mixtures at 10%, 20%, 30% and 40% v/v apple/pomegranate juices were then analyzed. The multivariate statistical analysis was applied to two groups at the time, where the pure pomegranate juices spectroscopic data were analyzed with respect to each mixture. Figures 2 and 3 show the corresponding PCA, PLS-DA and OPLS-DA scores plots for the analysis of pomegranate juices (P) with respect to mixtures of respectively 10% (A10) (Fig. 2a–c), 20% (A20) (Fig. 2d–f), 30% (A30) (Fig. 3a–c) and 40% (A40) (Fig. 3d–f) v/v apple/pomegranate juices. The  $R^2$  (cum) and  $Q^2$  (cum) values for the overall analyses are summarized in Table 1. The analyses show that the cluster correlation [ $R^2$  (cum)] and predictability [ $Q^2$  (cum)] increase as the percentage of apple juice in the mixtures increases, i.e. the method works better greater the amount of the filler (apple juice). The coefficient plots indicated that the spectral variables of importance responsible for the separation belong to the 225–305 nm region (Fig. 4). The second set of experiments showed that supervised OPLS-DA analysis applied to UV-Vis spectra data was able to detect pomegranate juices adulterated by more than 20% v/v apple juices. While unsupervised PCA analysis alone proved to be unable to properly detect pomegranate juices adulterated with up to 40% v/v apple juice (Figs. 1S–10S).




**Figure 1.** (a) PCA Scores plot; (b) PLS-DA Scores plot; (c) OPLS-DA Scores plot; analysis comparing UV-Vis spectra of apple juice samples (circles) to those of pomegranate juices (squares).


## Conclusions

Detection of adulteration in fresh fruit juices has become of broad interest to consumers, retailers and producers. The possibility to develop inexpensive methods to rapidly detect samples that have been adulterated would represent a major goal in this sector.

In this study UV-visible spectroscopy was combined with supervised and unsupervised multivariate statistical analysis to detect pomegranate juices adulterated with less expensive apple juice concentrate. The analyses show that OPLS-DA applied to UV-Vis spectra can only detected pomegranate fruit juices adulterated by more than 20% v/v apple juices, while unsupervised PCA is unable to properly detect pomegranate juices adulterated with up to 40% v/v apple juice.




**Figure 2.** (a) PCA Scores plot; (b) PLS-DA Scores plot; (c) OPLS-DA Scores plot; analysis comparing UV-Vis spectra of pomegranate juices (squares) to those of 10% (triangles) mixtures. (d) PCA Scores plot; (e) PLS-DA Scores plot; (f) OPLS-DA Scores plot; analysis comparing UV-Vis spectra of pomegranate juices (squares) to those of 20% (pentagons) mixtures.



**Figure 3.** (a) PCA Scores plot; (b) PLS-DA Scores plot; (c) OPLS-DA Scores plot; analysis comparing UV-Vis spectra of pomegranate juices (squares) to those of 30% (inverted triangles) mixtures. (d) PCA Scores plot; (e) PLS-DA Scores plot; (f) OPLS-DA Scores plot; analysis comparing UV-Vis spectra of pomegranate juices (squares) to those of 40% (diamonds) mixtures.

Groups*	PLS-DA		OPLS-DA	
	R <sup>2</sup> (cum)	Q <sup>2</sup> (cum)	R <sup>2</sup> (cum)	Q <sup>2</sup> (cum)
P vs A10	0.277	0.193	0.300	0.222
P vs A20	0.513	0.446	0.846	0.748
P vs A30	0.910	0.769	0.888	0.832
P vs A40	0.962	0.914	0.948	0.922
P vs A	0.991	0.987	0.991	0.989

**Table 1.** Goodness of fit of the data and predictability of the models. \*Pomegranate juices (P), apple juices (A), 10% (A10), 20% (A20), 30% (A30), 40% (A40) v/v apple/pomegranate mixtures.



**Figure 4.** Coefficient plot derived from the OPLS-DA multivariate statistical analysis of UV-Vis spectra of pomegranate juices vs 30% v/v apple/pomegranate juice mixtures. The spectral variables responsible for the separation among the groups are represented in red. These are the regions of the spectra mostly affected by the adulteration.

### Data availability

Data is available upon reasonable request.

Received: 2 August 2021; Accepted: 9 February 2022

Published online: 25 March 2022

### References

- Jayaprakasha, G. K., Negi, P. S. & Jena, B. S. Antimicrobial activities of pomegranate. In *Pomegranates: Ancient Roots to Modern Medicine* (eds Seeram, N. P. *et al.*) 168 (CRC Press, 2006).
- Kandyli, P. & Kokkinomagoulos, E. Food applications and potential health benefits of pomegranate and its derivatives. *Foods* **9**, 122 (2020).
- Jandric, Z. & Cannavan, A. An investigative study on differentiation of citrus fruit/fruit juices by UPLC-QToF MS and chemometrics. *Food Control* **72**, 173–180 (2017).
- Jandric, Z. *et al.* Assessment of fruit juice authenticity using UPLC-QToF MS: A metabolomics approach. *Food Chem.* **148**, 7–17 (2014).
- Vaclavik, L., Schreiber, A., Lacina, O., Cajka, T. & Hajslova, J. Liquid chromatography-mass spectrometry-based metabolomics for authenticity assessment of fruit juices. *Metabolomics* **8**, 793–803 (2011).
- Dasenaki, M. E., Drakopoulou, S. K., Aalizadeh, R. & Thomaidis, N. S. Targeted and untargeted metabolomics as an enhanced tool for the detection of pomegranate juice adulteration. *Foods* **8**, 212 (2019).
- Marchetti, L., Pellati, F., Benvenuti, S. & Bertelli, D. Use of  $^1\text{H}$  NMR to detect the percentage of pure fruit juices in blends. *Molecules* **24**, 2592 (2019).
- Miaw, C. S. W. *et al.* Comparison of different multivariate classification methods for the detection of adulterations in grape nectars by using low-field nuclear magnetic resonance. *Food Anal. Methods* **13**(1), 108–118 (2020).
- Vardin, H., Tay, A., Ozen, B. & Mauer, L. Authentication of pomegranate juice concentrate using FTIR spectroscopy and chemometrics. *Food Chem.* **108**, 742–748 (2008).
- Boggia, R., Casolino, M. C., Hysenaj, V., Oliveri, P. & Zunin, P. A screening method based on UV-visible spectroscopy and multivariate analysis to assess addition of filler juices and water to pomegranate juices. *Food Chem.* **140**, 735–741 (2013).
- Fonville, J. M. *et al.* The evolution of partial least squares models and related chemometric approaches in metabolomics and metabolic phenotyping. *J. Chemom.* **24**, 636–649 (2010).
- Bylesjo, M. *et al.* OPLS discriminant analysis: combining the strengths of PLS-DA and SIMCA classification. *J. Chemom.* **20**, 341–351 (2006).
- Umetrics. *User Guide to SIMCA (v.13)* (MKS Umetrics AB, 2012).
- Trygg, J., Holmes, E. & Lundstedt, T. Chemometrics in metabolomics. *J. Proteome Res.* **6**, 469–479 (2007).
- Galindo-Prieto, B., Eriksson, L. & Trygg, J. Variable influence on projection (VIP) for orthogonal projections to latent structures (OPLS). *J. Chemom.* **28**, 623–632 (2014).

### Acknowledgements

The author expresses her deep gratitude to Dr. István Pelczér for his insightful suggestions and comments on this study. A special thank goes to Mr. Ziad Sara for his assistance with the UV-spectroscopy measurements. This work was supported, in part, by the Open Access Program from the American University of Sharjah.



### Disclaimer

This paper represents the opinions of the author and does not mean to represent the position or opinions of the American University of Sharjah.

### Author contributions

It is a single author paper.

### Funding

AUS FRG13-2-35 by American University of Sharjah.

### Competing interests

The author declares no competing interests.

### Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-07979-7>.

**Correspondence** and requests for materials should be addressed to L.P.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022