





 Cite this: *RSC Adv.*, 2021, 11, 33235

 Received 5th July 2021  
 Accepted 28th September 2021

DOI: 10.1039/d1ra05187c

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

# Consecutive reactions to construct tricarbonyl compounds and synthetic applications thereof†

 Diego Madroño, <sup>a</sup> Cesar A. Mujica-Martinez <sup>b</sup> and Alfredo Vázquez <sup>\*a</sup>

Lithium anions derived from O-carbonate-protected cyanohydrins undergo conjugate addition to cycloalkenones with the concomitant transfer of the alkoxy-carbonyl group to produce tricarbonyl compounds. These products offer numerous possibilities for further elaboration. The synthetic potential of the cascade products was demonstrated by forming bicyclic and tricyclic systems through intramolecular condensation reactions.

## Introduction

Since the pioneering work by Stork *et al.*,<sup>1</sup> O-protected cyanohydrins have become recognized as valuable acyl anion synthons.<sup>2,3</sup> Over the years, this approach has been widely used<sup>4a-k</sup> to achieve the chemical synthesis of organic molecules with diverse architectures (Scheme 1).


Among the different groups used to protect the hydroxyl group in cyanohydrins, carbonates<sup>5a-c</sup> display a moderately electrophilic carbon atom. This reactivity can be exploited to perform subsequent transformations to the anions' reaction with electrophiles (*i.e.*, transfer of an acyl group), thus enabling consecutive reactions. Consecutive reactions, also known as cascade or domino reactions,<sup>6</sup> are a practical strategy to form multiple bonds sequentially, simplifying the construction of organic molecules.

A recent paper<sup>7</sup> demonstrated that anions derived from O-carbonate-protected cyanohydrins undergo conjugate addition to cyclohexenone with concomitant transfer of the alkoxy-carbonyl group to produce  $\beta$ -keto- $\beta'$ -acylcycloalkanecarboxylic acid esters; however, these were isolated as the enol acetate derivatives.


Tricarbonyl compounds have been used for the synthesis of functionalized biphenyls *via* an oxidative aromatization with iodine<sup>8a</sup> and for the formal synthesis of ( $\pm$ )-cochlearol A.<sup>8b</sup> The oxidation of tricarbonyl compounds has been used for the preparation of propellanes, compounds showing a broad spectra of biological and pharmacological activities,<sup>8c</sup> as well as for the synthesis of heterocycles.<sup>8d</sup>

Considering the tremendous synthetic utility of the functionalities present in the putative intermediate products namely  $\beta$ -keto ester,<sup>9a-f</sup> gamma-keto ester,<sup>9g</sup> and 1,4-dione,<sup>10</sup> we attempted to directly obtain these intermediates by modifying Le Lagadec's procedure.<sup>7</sup>

Herein, we present a procedure to construct synthetically valuable tricarbonyl cyclic compounds featuring consecutive Michael–Claisen reactions of lithium anions derived from O-



Scheme 1 Pioneering work of Stork and Maldonado to use protected cyanohydrins as acyl anion equivalents.



Scheme 2 (A and B) Synthetic strategy for the construction of tricarbonyl compounds using consecutive reactions. (B and C) Plausible mechanism for the consecutive reactions illustrated for the formation of 10.

<sup>a</sup>Departamento de Química Orgánica, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad Universitaria, Cd. Mx., 04510, Mexico. E-mail: joseavm@unam.mx

<sup>b</sup>GIFBA, Departamento de Química, Facultad de Ciencias Exactas y Naturales, Centro de Investigación en Materiales CIMA, Universidad de Nariño, San Juan de Pasto, 520002, Colombia

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra05187c



carbonate-protected cyanohydrins with 5, 6, and 7-membered cycloalkenones. The functionalities introduced into the cycloalkenones was exploited by annulation of a five-membered ring onto the olefin of the original cycloalkenone.

## Results and discussion

Our strategy is shown in Scheme 2, consisting of the preparation of cyanocarboxylates **6** from aldehydes **1** according to literature procedures (Scheme 2A). Deprotonation with a suitable base (2 equiv.), followed by conjugate addition of the corresponding anions onto cyclic enones **7** (Scheme 2B) and subsequent reaction of the tricarboxyl product **9** (as an enolate) with electrophiles would afford highly functionalized products **10** (Scheme 2C).


The preparation of some cyanocarboxylates **6** was achieved using the two-phase reaction system procedure reported by Kolis *et al.*<sup>11</sup> (Table 1, Method A). In other cases, higher yields were obtained using ethyl carbonocyanidate (CNCOOEt) in the

presence of DMAP and CH<sub>3</sub>CN as the solvent (Table 1, Method B).<sup>5c</sup> A total of 23 cyanocarboxylates were prepared and satisfactorily characterized.


To evaluate the deprotonation ease of the NC-C-H bond, its  $pK_a$  was determined using a direct approach,<sup>12</sup> in which free energies are calculated directly in THF solution at -78 °C (Table 2). Calculations were carried out using the  $\omega$ B97XD DFT hybrid functional,<sup>13</sup> the 6-311++G(d,p) basis set, and the D2 Grimme dispersion correction<sup>14</sup> as implemented in the Gaussian 16 suite.<sup>15</sup> The solvent was described implicitly using the SMD method.<sup>16</sup> This methodology has been used before to determine  $pK_a$  in several systems.<sup>12,17</sup> Fig. 1 shows that the obtained  $pK_a$  values highly correlate with the Hammett constants<sup>18</sup> ( $\sigma$ ) of the phenyl substituent for compounds **6a**, **6d**, **6e**, and **6f**, for which it was obtained  $pK_a = 37.503 - 11.527\sigma$  and  $R^2 = 0.931$ . Therefore, the deprotonation ease of these compounds increases with the electron-withdrawing properties of the substituents, which results in the stabilization of the corresponding carbanion. Similar correlations are observed with other molecular parameters and also for compounds **7**. The relatively large  $pK_a$  values indicated the use of a strong base would be required to obtain the corresponding carbanion efficiently.

To test our hypothesis and standardize the reaction conditions, we selected cyanocarboxylate **6a** as the carbanion source and commercially available LiHMDS as the base. The base was added dropwise to a solution of **6a** in THF at -78 °C. After 15 min, cyclohex-2-en-1-one was added, and the course of the reaction was monitored by TLC.

Table 1 Preparation of cyanocarboxylates





<sup>a</sup> (Boc)<sub>2</sub>O was used instead of ClCOOMe, <sup>b</sup> The reaction was heated at 40 °C.



<sup>a</sup> (Boc)<sub>2</sub>O was used instead of ClCOOMe. <sup>b</sup> The reaction was heated at 40 °C.

Table 2 Scope of the cascade reactions between cyanocarboxylates **6** and **7** with 2-cyclohexen-1-one







Fig. 1 Correlation of  $pK_a$  with the Hammett constant of the phenyl substituent for compounds **6a**, **6d**, **6e**, **6f**.

To investigate the scope of the cascade process under the optimized conditions described above, the anions of cyanocarboxylates **6** and **7** were added to cyclohexanone. The results of those experiments are summarized in Table 2. A total 15 of tricarbonyl compounds (**8a–o** and **9a–c**) were obtained in moderate yields. All the products were successfully characterized.

Next, we investigated the use of 2-cyclopenten-1-one, 2-cyclohepten-1-one, and 3-methyl-2-cyclohexen-1-one as the enone component for the cascade process to further explore the scope of the methodology. The results are presented in Table 3. For 2-cyclopenten-1-one, the tricarbonyl compound **9a** was obtained in 80% yield, whereas for 2-cyclohepten-1-one the desired product **9b** was obtained in 50% yield. In the case of 3-methyl-2-cyclohexen-1-one, **9c** was obtained in 38% yield. This lower yield was attributed to the steric hindrance caused by the methyl group at C-3 of the cycloalkenone. This assumption is supported by the failure to obtain the cascade product when the more sterically hindered 4,4-dimethyl-2-cyclohexen-1-one was used as the substrate.

Interestingly,  $^1\text{H}$  NMR spectra for all the cascade products obtained from 2-cyclohexen-1-one and 3-methyl-2-cyclohexen-1-one were isolated as a mixture of keto and enol tautomers. In contrast only a single keto tautomer or the keto diastereomers

Table 3 Use of different cycloalkenones to explore the scope of the process



were observed by  $^1\text{H}$  NMR for the adducts derived from 2-cyclopenten-1-one and 2-cyclohepten-1-one (*i.e.*, **9a** and **9b**). Computational results indicate that the keto-tautomer of compounds **9a** and **9b** is 1.365 and 1.366 kcal mol $^{-1}$ , respectively, more stable than the corresponding enol-tautomer. On the contrary, for compound **8a**, the enol-tautomer is only 0.274 kcal mol $^{-1}$  more stable than the corresponding keto-tautomer. This small energy difference could explain the reason to observe this product as a mixture of tautomers. Further investigation on the tautomeric behavior of these systems is currently underway. Clearly, the cascade process can occur for 5, 6, and 7-membered cycloalkenones, even enones showing moderate steric hindrance.

After proving the efficiency of the cascade process, we investigated the possibility of performing annulations *via* intramolecular condensation reactions. Thus, when **14b** was treated with NaH in THF at rt, followed by the addition of triphenylvinylphosphonium bromide (Schweizer's reagent)<sup>19</sup> no reaction was observed. If the reaction mixture was heated under reflux, several spots are observed on TLC. However, when DBU was used as the base (in CH $_3$ CN), 39% of intramolecular Wittig product **10** was obtained after purification, along with unreacted starting material and traces of two unknown compounds. Treatment of **8b** with methylvinyl ketone and cyclopentenone as the Michael acceptors in the presence of DBU (in CH $_3$ CN at room temperature), afforded cyclic products **11** and **12** in 88 and 32% yield, respectively (Table 4). It is noteworthy that decarboxylation occurred during the formation of **11** and **12**.

The preparation of annulated products **10**, **11**, and **12** nicely exemplifies the synthetic potential of  $\beta$ -keto- $\beta'$ -acylcycloalkancarboxylic acid esters **8** as scaffolds to obtain products with increased structural complexity. With some adjustments, we believe that compounds **8** can be used to obtain diverse molecules such as the core of pacifigorgianes<sup>20</sup> **13**, the

Table 4 Further transformations of cascade product **8b**

Temp	Michael acceptor	Product	Yield (%)
rt		<b>10</b>	39
rt		<b>11</b>	88
Reflux		<b>12</b>	32




Fig. 2 Potential use of tricarbonyl compounds **8** to generate structural diversity.

sesquiterpenoid cyperolone<sup>21</sup> **14**, indanones<sup>22</sup> **15**, furans<sup>23</sup> pyrroles and thiophenes **16**, 1,2-azoles **17**, pyrimidines **18** and 1,2-diazines **19** (Fig. 2).

## Experimental

### General information

All experiments involving air and/or moisture-sensitive compounds were conducted in an oven dried round-bottom flask capped with a rubber septum under a positive pressure of nitrogen. Air- and moisture-sensitive liquids were transferred *via* syringe or stainless-steel cannula. Low temperature baths were ice/water (0 °C), CO<sub>2</sub>(s)/acetone (−78 °C). Unless otherwise noted, reaction temperatures refer to that of the bath. Concentration refers to removal of volatiles on a rotary evaporator below 35 °C. Analytical thin-layer chromatography (TLC) was performed using glass plates pre-coated with silica gel (0.25 mm, 60 Å pore size, 230–400 mesh, Merck) impregnated with a fluorescent indicator (254 nm). Materials on TLC plates were visualized under an ultraviolet lamp (254 nm) and/or by submersion of the plate in a solution of phosphomolybdic acid (5%) containing a trace of ceric sulfate in aqueous sulfuric acid (5% v/v) followed by charring on a hot plate. Flash column chromatography (FCC) was performed according to Still *et al.* with silica gel 60 (40–63 μm). All mixed solvent eluents are reported as v/v solutions. Unless otherwise noted, all reported compounds were homogeneous by thin layer chromatography (TLC) and by <sup>1</sup>H NMR.

### Materials and methods

All the reagents were purchased from Sigma-Aldrich and were used as received unless other thing stated. Solvents were distilled prior to use. Anhydrous solvents were distilled under nitrogen atmosphere. THF and diethyl ether were distilled on sodium benzophenone ketyl; MeOH on magnesium activated with 5% iodine. Et<sub>3</sub>N, CH<sub>3</sub>CN dichloroethane and DMF were distilled on CaH<sub>2</sub>.

Proton nuclear magnetic resonance (<sup>1</sup>H NMR) spectra and carbon nuclear magnetic resonance (<sup>13</sup>C NMR) spectra were recorded on Agilent-Inova-300 and Varian VNMRs-400 NMR spectrometers. Proton chemical shifts are expressed in parts per million (ppm, δ scale) and are referenced to tetramethylsilane

(TMS: 0.0). Carbon chemical shifts are expressed in parts per million (ppm, δ scale) and are referenced to tetramethylsilane (TMS: 0.0). For compounds **6e**, **7e**, **7f**, **7h**, **7p**, **8g**, and **8n**, CDCl<sub>3</sub> was used as a standard. This information is included in the ESI.† Data are represented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, dd = doublet of doublets, dt = doublet of triplets, td = triplet of doublets, ddd = doublet of doublets of doublets, ddt = doublet of doublets of triplets, dqd = doublet of quartets of doublets, m = multiplet, br = broad, app = apparent), integration, and coupling constant (*J*) in Hertz (Hz).

IR spectra were recorded on a PerkinElmer Spectrum 400 FT-IR/FIR spectrometer with ATR. Mass spectra were carried out on a JEOL SMX-102a spectrometer.

### Method A

A 2 M aqueous solution of NaCN (4.8 mL) was added dropwise to a mixture of the aldehyde (5 mmol), ClCOOMe (0.43 mL, 5.5 mmol) and (*n*Bu)<sub>4</sub>NBr (160 mg, 0.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6.25 mL) at ambient temperature. The resultant mixture was vigorously stirred overnight, the two phases were separated, and the organic layer was washed with sat. NaCl dried on anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>).

### Method B

CNCOOEt (0.108 mL, 1.1 mmol) was slowly added to a mixture of aldehyde (1 mmol) and DMAP (6 mg, 0.05 mmol) in CH<sub>3</sub>CN (2 mL) under nitrogen atmosphere at ambient temperature. When all the starting material has been consumed (TLC) the reaction mixture is poured on sat NaCl (5 mL) and extracted with EtOAc (3 × 20 mL). The organic extracts were combined and dried on anhydrous Na<sub>2</sub>SO<sub>4</sub>, the solvent evaporated *in vacuo* and the residue purified by column chromatography.

**Cyano(phenyl)methyl methyl carbonate (6a).** Prepared following Method A. The residue was purified by column chromatography (hexanes/EtOAc 95 : 5) to obtain 755 mg (79%) of a translucent yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 3.87 (s, 3H, OCH<sub>3</sub>), 6.27 (s, 1H, CHCN), 7.47–7.48 (m, 3H, *m,p*-C<sub>6</sub>H<sub>5</sub>), 7.53–7.56 (m, 2H, *o*-C<sub>6</sub>H<sub>5</sub>) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 56.0, 66.7, 115.8, 128.0, 129.4, 130.8, 131.3, 154.2 ppm. APCI: *m/z* calculated for C<sub>8</sub>H<sub>8</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 192.0661; found: 192.0703.

**Cyano(furan-2-yl)methyl methyl carbonate (6b).** Prepared following Method A. The residue was purified by column chromatography (hexanes/EtOAc 95 : 5) to obtain 729 mg (80%) of a yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 3.86 (s, 3H, OCH<sub>3</sub>), 6.34 (s, 1H, CHCN), 6.44–6.45 (dd, 1H, *J* = 1.8, 3.2 Hz, 3-C<sub>4</sub>H<sub>3</sub>O), 6.72–6.73 (d, 1H, *J* = 3.2 Hz, 4-C<sub>4</sub>H<sub>3</sub>O), 7.51–7.52 (d, 1H, *J* = 1.8 Hz, 5-C<sub>4</sub>H<sub>3</sub>O). <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 56.1, 59.5, 111.3, 113.3, 113.8, 143.6, 145.4, 154.0 ppm. APCI: *m/z* calculated for C<sub>8</sub>H<sub>8</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 182.0453; found: 182.0502.

**tert-Butyl (cyano(phenyl)methyl)carbonate (6c).** A 2 M aqueous solution of NaCN (4.8 mL) was added dropwise to a mixture of benzaldehyde (5 mmol), Boc<sub>2</sub>O (1.2 g, 5.5 mmol) and (*n*Bu)<sub>4</sub>NBr (16 mg, 0.05 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6.25 mL) at ambient temperature. The resulting mixture was stirred



overnight, the two phases were separated, and the organic layer was washed with sat NaCl, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 845 mg (72%) of a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.44 (s, 9H, *Ot*-Bu), 6.14 (s, 1H, CHCN), 7.34–7.36 (m, 3H, *m,p*-C<sub>6</sub>H<sub>5</sub>), 7.43–7.45 (m, 2H, *o*-C<sub>6</sub>H<sub>5</sub>) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 27.7, 65.8, 85.0, 116.2, 128.0, 129.3, 130.6, 131.7, 151.7 ppm. APCI: *m/z* calculated for C<sub>13</sub>H<sub>16</sub>NO<sub>3</sub> [M + H]<sup>+</sup> = 234.1130; found: 234.1124.

**Cyano(4-methoxyphenyl)methyl methyl carbonate (6d).** Prepared according to Method A with the following modification: the reaction mixture was stirred at 40 °C for 12 h. The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 700 mg (63%) of a pale-yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 3.82 (s, 3H, COOCH<sub>3</sub>), 3.84 (s, 3H, OCH<sub>3</sub>), 6.20 (s, 1H, CHCN), 6.95 (m, 2H *m*-C<sub>6</sub>H<sub>4</sub>OMe), 7.47 (m, 2H, *o*-C<sub>6</sub>H<sub>4</sub>OMe) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 55.5, 55.9, 66.5, 114.7, 116.0, 123.4, 129.9, 154.2, 161.5 ppm. APCI: *m/z* calculated for C<sub>11</sub>H<sub>12</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 222.0766; found: 222.0769.

**(2-Bromophenyl)(cyano)methyl methyl carbonate (6e).** Prepared following Method A. The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 1.23 g (91%) of a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 3.89 (s, 3H, OCH<sub>3</sub>), 6.57 (s, 1H, CHCN), 7.61–7.64 (m, 1H, *m*-C<sub>6</sub>H<sub>4</sub>Br), 7.73–7.76 (m, 1H, *m*-C<sub>6</sub>H<sub>4</sub>Br), 7.87–7.89 (m, 1H, *p*-C<sub>6</sub>H<sub>4</sub>Br), 8.14–7.16 (m, 1H, *o*-C<sub>6</sub>H<sub>4</sub>Br) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 56.4, 63.0, 114.7, 126.0, 126.9, 129.3, 131.6, 134.9, 146.9, 153.6 ppm. ESI<sup>+</sup>: *m/z* calculated for C<sub>10</sub>H<sub>9</sub>BrNO<sub>3</sub> [M + H]<sup>+</sup> = 269.9766; found: 269.9775.

**Cyano(2-nitrophenyl)methyl methyl carbonate (6f).** Prepared following Method A. The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 1.16 g (93%) of a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 3.89 (s, 3H, OCH<sub>3</sub>), 7.01 (s, 1H, CHCN), 7.67–7.71 (m, 1H, *m*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 7.79–7.83 (m, 1H, *m*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 7.94–7.96 (m, 1H, *p*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 8.21–8.24 (m, 1H, *o*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 56.4, 63.0, 114.7, 126.0, 127.0, 129.3, 131.7, 134.9, 146.9, 153.6 ppm. ESI<sup>+</sup>: *m/z* calculated for C<sub>10</sub>H<sub>9</sub>N<sub>2</sub>O<sub>5</sub> [M + H]<sup>+</sup> = 237.0511; found: 237.0510.

**(6-Bromobenzo[*d*][1,3]dioxol-5-yl)(cyano)methyl methyl carbonate (6g).** Prepared according to Method A with the following modification: the reaction mixture was stirred at 40 °C for 12 h. The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 1.32 g (84%) of a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 3.88 (s, 3H, OCH<sub>3</sub>), 6.04 (s, 2H, OCH<sub>2</sub>O), 6.51 (s, 1H, CHCN), 7.04 (s, 1H, *m*-C<sub>6</sub>H<sub>2</sub>Br), 7.16 (s, 1H, *o*-C<sub>6</sub>H<sub>2</sub>Br) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 56.2, 66.2, 102.8, 109.3, 113.3, 114.9, 115.3, 123.8, 148.3, 150.5, 153.9 ppm. ESI<sup>+</sup>: *m/z* calculated for C<sub>11</sub>H<sub>9</sub>BrNO<sub>5</sub> [M + H]<sup>+</sup> = 313.9664; found: 313.9658.

**Cyano(phenyl)methyl ethyl carbonate (7a).** Prepared according to Method B (reaction time 8 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 203 mg (99%) of a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.32 (t, 3H, *J* = 7.2 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 4.21–4.33 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 6.26 (s, 1H, CHCN), 7.44–7.45 (m, 3H, *m,p*-C<sub>6</sub>H<sub>5</sub>),

7.52–7.54 (m, 2H, *o*-C<sub>6</sub>H<sub>5</sub>) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.2, 65.7, 66.5, 115.9, 128.0, 129.4, 130.7, 131.4, 153.5 ppm. APCI: *m/z* calculated for C<sub>11</sub>H<sub>12</sub>NO<sub>3</sub> [M + H]<sup>+</sup> = 206.0817; found: 206.0849.

**Cyano(furan-2-yl)methyl ethyl carbonate (7b).** Prepared according to Method B (reaction time 8 h). The residue was purified by column chromatography (hexanes/EtOAc 95 : 5) to obtain 190 mg (97%) of a translucent yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.32 (t, 3H, *J* = 7.1 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 4.22–4.33 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 6.32 (s, 1H, CHCN), 6.43 (dd, 1H, *J* = 1.8, 3.2 Hz, 3-C<sub>4</sub>H<sub>3</sub>O), 6.71 (d, 1H, *J* = 3.2 Hz, 4-C<sub>4</sub>H<sub>3</sub>O), 7.5 (d, 1H, *J* = 1.8 Hz, 5-C<sub>4</sub>H<sub>3</sub>O) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.2, 59.3, 65.9, 111.3, 113.1, 113.9, 143.8, 145.4, 153.3 ppm. APCI: *m/z* calculated for C<sub>9</sub>H<sub>10</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 196.0610; found: 196.0626.

**Cyano(thiophen-2-yl)methyl ethyl carbonate (7c).** Prepared according to Method B (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 195 mg (92%) of a brown oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.33 (t, 3H, *J* = 7.16 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 4.25–4.32 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 6.48 (s, 1H, CHCN), 7.04–7.05 (m, 1H, 4-C<sub>4</sub>H<sub>3</sub>S), 7.36–7.38 (m, 1H, 3-C<sub>4</sub>H<sub>3</sub>S), 7.46–7.47 (m, 1H, 5-C<sub>4</sub>H<sub>3</sub>S) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.1, 61.5, 65.8, 115.0, 127.3, 129.4, 130.0, 132.7, 153.2 ppm. APCI: *m/z* calculated for C<sub>9</sub>H<sub>10</sub>NO<sub>3</sub>S [M + H]<sup>+</sup> = 212.0381; found: 212.0395.

***tert*-Butyl 2-(cyano((ethoxycarbonyloxy)methyl)-1H-pyrrole-1-carboxylate (7d).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 70 : 30) to obtain 248 mg (84%) of a translucent orange oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.33 (t, 3H, *J* = 7.13 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.58 (s, 9H, *Ot*-Bu), 4.22–4.33 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 6.18–6.20 (t, 1H, *J* = 3.4 Hz, 4-C<sub>4</sub>H<sub>3</sub>N), 6.70–6.71 (m, 1H, 3-C<sub>4</sub>H<sub>3</sub>N), 6.81 (s, 1H, CHCN), 7.31–7.32 (m, 1H, 5-C<sub>4</sub>H<sub>3</sub>N) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.3, 28.0, 60.3, 65.5, 85.7, 110.7, 115.4, 117.6, 123.5, 124.7, 148.4, 153.5 ppm. APCI: *m/z* calculated for C<sub>14</sub>H<sub>19</sub>N<sub>2</sub>O<sub>5</sub> [M + H]<sup>+</sup> = 295.1294; found: 295.1214.

**Cyano(pyridin-2-yl)methyl ethyl carbonate (7e).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 190 mg (92%) of a translucent yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.35 (t, 3H, *J* = 7.15 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 4.27–4.35 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 6.37 (s, 1H, CHCN), 7.37–7.39 (m, 1H, 5-C<sub>5</sub>H<sub>4</sub>N), 7.58–7.60 (m, 1H, 3-C<sub>5</sub>H<sub>4</sub>N), 7.80–7.83 (m, 1H, 4-C<sub>5</sub>H<sub>4</sub>N), 8.66–8.67 (m, 1H, 6-C<sub>5</sub>H<sub>4</sub>N) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.2, 66.0, 67.2, 115.3, 121.9, 125.0, 137.8, 150.3, 150.7, 153.4 ppm. APCI: *m/z* calculated for C<sub>13</sub>H<sub>14</sub>NO<sub>5</sub> [M + H]<sup>+</sup> = 207.0770; found: 207.0797.

**Cyano(pyridin-3-yl)methyl ethyl carbonate (7f).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 197 mg (96%) of a translucent yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.32 (t, 3H, *J* = 7.13 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 4.23–4.33 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 6.30 (s, 1H, CHCN), 7.40–7.42 (m, 1H, 5-C<sub>5</sub>H<sub>4</sub>N), 7.89–7.91 (m, 1H, 4-C<sub>5</sub>H<sub>4</sub>N), 8.70–8.71 (m, 1H, 6-C<sub>5</sub>H<sub>4</sub>N), 8.76–8.77 (m, 1H, 2-C<sub>5</sub>H<sub>4</sub>N) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.2, 64.3, 66.1, 115.1, 124.1, 127.6, 135.6, 149.2, 152.0, 153.3 ppm. APCI: *m/z* calculated for C<sub>13</sub>H<sub>14</sub>NO<sub>5</sub> [M + H]<sup>+</sup> = 207.0770; found: 207.0797.

**Cyano(pyridin-4-yl)methyl ethyl carbonate (7g).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 190 mg (92%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.29 (t, 3H,  $J = 7.14$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 4.22–4.29 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.22 (s, 1H, CHCN), 7.40 (d, 2H,  $J = 6.19$  Hz, 3,5- $\text{C}_5\text{H}_4\text{N}$ ), 8.68 (d, 2H,  $J = 6.19$  Hz, 2,6- $\text{C}_5\text{H}_4\text{N}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.2, 64.8, 66.2, 114.8, 121.6, 139.8, 150.9, 153.3 ppm. APCI:  $m/z$  calculated for  $\text{C}_{13}\text{H}_{14}\text{NO}_5$   $[\text{M} + \text{H}]^+$  = 207.0770, found: 207.0809.

**Cyano(2-nitrophenyl)methyl ethyl carbonate (7h).** Prepared according to Method B (reaction time 2 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 245 mg (98%) of colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.35 (t, 3H,  $J = 7.15$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 4.26–4.34 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 7.00 (s, 1H, CHCN), 7.67–7.70 (m, 1H,  $m\text{-C}_6\text{H}_4\text{NO}_2$ ), 7.79–7.83 (m, 1H,  $m\text{-C}_6\text{H}_4\text{NO}_2$ ), 7.94–7.96 (m, 1H,  $p\text{-C}_6\text{H}_4\text{NO}_2$ ), 8.21–8.23 (m, 1H,  $o\text{-C}_6\text{H}_4\text{NO}_2$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.2, 62.8, 66.2, 114.8, 126.0, 127.1, 129.3, 131.6, 134.9, 146.9, 153.0 ppm. APCI:  $m/z$  calculated for  $\text{C}_{11}\text{H}_{11}\text{N}_2\text{O}_5$   $[\text{M} + \text{H}]^+$  = 251.0668; found: 251.0599.

**Benzofuran-2-yl(cyano)methyl ethyl carbonate (7i).** Prepared according to Method B (reaction time 5 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 216 mg (88%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.36 (t, 3H,  $J = 7.15$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 4.29–4.37 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.49 (s, 1H, CHCN), 7.10 (s, 1H, 3- $\text{C}_8\text{H}_5\text{O}$ ), 7.28–7.32 (m, 1H, 6- $\text{C}_8\text{H}_5\text{O}$ ), 7.38–7.42 (m, 1H, 5- $\text{C}_8\text{H}_5\text{O}$ ), 7.52–7.54 (m, 1H, 7- $\text{C}_8\text{H}_5\text{O}$ ), 7.62–7.64 (m, 1H, 4- $\text{C}_8\text{H}_5\text{O}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.2, 59.9, 66.1, 109.5, 112.0, 113.7, 122.2, 123.9, 126.6, 127.0, 145.8, 153.3, 155.9 ppm. APCI:  $m/z$  calculated for  $\text{C}_{13}\text{H}_{12}\text{NO}_4$   $[\text{M} + \text{H}]^+$  = 246.0766, found: 246.0770.

**Benzo[*b*]thiophen-2-yl(cyano)methyl ethyl carbonate (7j).** Prepared according to Method B (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 226 mg (86%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.36 (t, 3H,  $J = 7.14$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 4.28–4.36 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.58 (s, 1H, CHCN), 7.41–7.44 (m, 2H, 5,6- $\text{C}_8\text{H}_5\text{S}$ ), 7.62–7.63 (m, 1H, 3- $\text{C}_8\text{H}_5\text{S}$ ), 7.81–7.86 (m, 2H, 4,7- $\text{C}_8\text{H}_5\text{S}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.2, 62.4, 66.0, 114.8, 122.7, 124.8, 125.2, 126.2, 126.8, 133.3, 138.4, 140.9, 153.3 ppm. APCI:  $m/z$  calculated for  $\text{C}_{13}\text{H}_{13}\text{NO}_4$   $[\text{M} - \text{OEt}]^+$  = 216.0119, found: 216.0106.

***tert*-Butyl 2-(cyano((ethoxycarbonyl)oxy)methyl)-1*H*-indole-1-carboxylate (7k).** Prepared according to Method B (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 70 : 30) to obtain 275 mg (80%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.36 (t, 3H,  $J = 7.10$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.70 (s, 9H, *tert*-Bu), 4.30–4.35 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.99 (s, 1H, CHCN), 7.08–7.09 (m, 1H, 3- $\text{C}_8\text{H}_5\text{N}$ ), 7.25–7.29 (m, 1H, 5- $\text{C}_8\text{H}_5\text{N}$ ), 7.35–7.39 (m, 1H, 6- $\text{C}_8\text{H}_5\text{N}$ ), 7.57–7.59 (m, 1H, 4- $\text{C}_8\text{H}_5\text{N}$ ), 8.06–8.09 (m, 1H, 7- $\text{C}_8\text{H}_5\text{N}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.3, 28.2, 61.6, 65.7, 86.1, 112.6, 115.3, 116.0, 121.7, 123.6, 126.2, 127.8, 129.3, 137.0, 149.8, 153.4 ppm. APCI:  $m/z$  calculated for  $\text{C}_{18}\text{H}_{21}\text{N}_2\text{O}_5$   $[\text{M} + \text{H}]^+$  = 345.1450, found: 245.1482.

**Methyl 2-(cyano((ethoxycarbonyl)oxy)methyl)benzoate (7l).** Prepared according to Method B (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 232 mg (88%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.33 (t,  $J = 7.11$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ), 3.93 (s, 3H, OMe), 4.25–4.33 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 7.28 (s, 1H, CHCN), 7.51–7.55 (m, 1H,  $m\text{-C}_6\text{H}_4\text{CO}_2\text{Me}$ ), 7.63–7.67 (m, 1H,  $m\text{-C}_6\text{H}_4\text{CO}_2\text{Me}$ ), 7.83–7.86 (m, 1H,  $p\text{-C}_6\text{H}_4\text{CO}_2\text{Me}$ ), 8.05–8.10 (m, 1H,  $o\text{-C}_6\text{H}_4\text{CO}_2\text{Me}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.1, 52.7, 63.6, 65.6, 115.9, 128.1, 128.2, 130.2, 131.5, 132.8, 133.4, 153.2, 166.2 ppm. APCI:  $m/z$  calculated for  $\text{C}_{13}\text{H}_{14}\text{NO}_5$   $[\text{M} + \text{H}]^+$  = 264.0872, found: 264.0841.

**(2-Bromophenyl)(cyano)methyl ethyl carbonate (7m).** Prepared according to Method B (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 270 mg (95%) of translucent oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.34 (t, 3H,  $J = 7.15$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 4.27–4.33 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.57 (s, 1H, CHCN), 7.30–7.34 (m, 1H,  $m\text{-C}_6\text{H}_4\text{Br}$ ), 7.40–7.45 (m, 1H,  $m\text{-C}_6\text{H}_4\text{Br}$ ), 7.61–7.64 (m, 1H,  $p\text{-C}_6\text{H}_4\text{Br}$ ), 7.71–7.74 (m, 1H,  $o\text{-C}_6\text{H}_4\text{Br}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.1, 65.8 ( $\times 2$ ), 115.1, 123.1, 128.3, 129.6, 130.7, 132.0, 133.5, 153.1 ppm. ESI $^+$ :  $m/z$  calculated for  $\text{C}_{11}\text{H}_{11}\text{BrNO}_3$   $[\text{M} + \text{H}]^+$  = 283.9922, found: 283.9927.

**Cyano(2-methoxyphenyl)methyl ethyl carbonate (7n).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 212 mg (90%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.33 (t,  $J = 7.12$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ), 3.88 (s, 3H, OMe), 4.24–4.32 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.58 (s, 1H, CHCN), 6.92–6.95 (m, 1H,  $o\text{-C}_6\text{H}_4\text{OMe}$ ), 7.00–7.05 (m, 1H,  $p\text{-C}_6\text{H}_4\text{OMe}$ ), 7.40–7.44 (m, 1H,  $m\text{-C}_6\text{H}_4\text{OMe}$ ), 7.56–7.58 (m, 1H,  $m\text{-C}_6\text{H}_4\text{OMe}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.1, 55.7, 61.7, 65.4, 111.1, 115.9, 119.5, 120.9, 128.9, 132.0, 153.5, 156.7 ppm. APCI:  $m/z$  calculated for  $\text{C}_{12}\text{H}_{14}\text{NO}_4$   $[\text{M} + \text{H}]^+$  = 236.0923, found: 236.0930.

**Cyano(3-methoxyphenyl)methyl ethyl carbonate (7o).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 222 mg (94%) of a translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.33 (t,  $J = 7.12$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ), 3.83 (s, 3H, OMe), 4.24–4.32 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 6.22 (s, 1H, CHCN), 6.97–7.00 (m, 1H,  $o\text{-C}_6\text{H}_4\text{OMe}$ ), 7.04–7.05 (m, 1H,  $o\text{-C}_6\text{H}_4\text{OMe}$ ), 7.09–7.11 (m, 1H,  $p\text{-C}_6\text{H}_4\text{OMe}$ ), 7.33–7.37 (m, 1H,  $m\text{-C}_6\text{H}_4\text{OMe}$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.1, 55.4, 65.6, 66.2, 113.0, 116.4, 116.5, 120.0, 130.4, 132.5, 153.4, 160.1 ppm. APCI:  $m/z$  calculated for  $\text{C}_{12}\text{H}_{14}\text{NO}_4$   $[\text{M} + \text{H}]^+$  = 236.0923, found: 236.0955.

**Cyano(cyclohexyl)methyl ethyl carbonate (7p).** Prepared according to Method B (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 180 mg (85%) of a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.15–1.29 (m, 5H), 1.33 (t, 3H,  $J = 7.13$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.68–1.93 (m, 6H), 4.21–4.31 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 5.03 (d,  $J = 5.88$  Hz, 1H, CHCN) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.2, 25.3, 25.8, 28.1, 40.3, 65.5, 69.4. APCI:  $m/z$  calculated for  $\text{C}_{11}\text{H}_{18}\text{NO}_3$   $[\text{M} + \text{H}]^+$  = 212.1287, found: 212.1255.

### General procedure for conjugate additions

1 M LiHMDS (THF) (750  $\mu$ L, 0.75 mmol) was slowly added to a solution of cyanocarbonate (0.5 mmol) in THF (2.5 mL) at  $-78$  °C under nitrogen atmosphere, and the resultant mixture stirred at this temperature for 15 min. Then, the enone (0.75 mmol) was added dropwise, and after the addition was completed, the mixture was allowed to reach ambient temperature. When the starting material has been consumed (TLC), the reaction was quenched by the slow addition of sat  $\text{NH}_4\text{Cl}$  (5 mL). The mixture was diluted with water and extracted with EtOAc ( $3 \times 20$  mL); the organics were combined, dried (anh  $\text{Na}_2\text{SO}_4$ ) and concentrated *in vacuo*. The residue was purified by column chromatography ( $\text{SiO}_2$ ).

**Methyl 6-benzoyl-2-hydroxycyclohex-1-ene-1-carboxylate (8a).** Prepared following the general procedure (reaction time 2 h). The residue was purified by column chromatography (hexanes/EtOAc 95 : 5) to obtain 103 mg (79%) of a yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.65–1.95 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.17–2.44 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 3.55 (s, 3H, OMe), 4.48–4.50 (m, 1H, CH), 7.44–7.46 (m, 2H, *m*- $\text{C}_6\text{H}_5$ ), 7.47–7.48 (m, 1H, *p*- $\text{C}_6\text{H}_5$ ), 7.95–7.98 (m, 2H, *o*- $\text{C}_6\text{H}_5$ ), 12.34 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 18.5, 26.1, 28.9, 41.0, 51.5, 97.2, 128.3, 128.7, 132.9, 133.7, 172.0, 174.5, 201.7 ppm. IR ( $\text{cm}^{-1}$ ): 3334, 1738, 1714, 1673, 1333. APCI:  $m/z$  calculated for  $\text{C}_{15}\text{H}_{17}\text{O}_4$  [ $\text{M} + \text{H}$ ] $^+$  = 261.1127, found: 261.1064.

**Ethyl 6-benzoyl-2-hydroxycyclohex-1-ene-1-carboxylate (8b).** Prepared following the general procedure (reaction time 2 h). The residue was purified by column chromatography (hexanes/EtOAc 95 : 5) to obtain 103 mg (75%) of a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 0.97 (t, 3H,  $J = 7.12$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.67–1.94 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.32–2.39 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 4.01–4.06 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 4.50–4.53 (m, 1H, CH), 7.46–7.49 (m, 2H, *m*- $\text{C}_6\text{H}_5$ ), 7.55–7.57 (m, 1H, *p*- $\text{C}_6\text{H}_5$ ), 7.96–8.01 (m, 2H, *o*- $\text{C}_6\text{H}_5$ ), 12.37 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 13.9, 18.8, 26.3, 29.0, 41.0, 60.5, 97.6, 128.4, 128.7, 132.9, 136.6, 171.7, 174.5, 202.1 ppm. IR ( $\text{cm}^{-1}$ ): 3448, 1740, 1682, 1651, 1334. APCI:  $m/z$  calculated for  $\text{C}_{16}\text{H}_{19}\text{O}_4$  [ $\text{M} + \text{H}$ ] $^+$  = 275.1283; found: 275.1242.

***tert*-Butyl 6-benzoyl-2-hydroxycyclohex-1-ene-1-carboxylate (8c).** Prepared following the general procedure (reaction time 5 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 94 mg (62%) of a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.21 (s, 9H, *Ot*-Bu), 1.63–1.74 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.30–2.34 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 4.43–4.47 (m, 1H, CH), 7.43–7.53 (m, 2H, *p*- $\text{C}_6\text{H}_5$ ), 7.49–7.55 (m, 1H, *m*- $\text{C}_6\text{H}_5$ ), 7.94–8.01 (m, 2H, *o*- $\text{C}_6\text{H}_5$ ), 12.55 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 19.0, 26.3, 28.1, 29.1, 41.3, 81.9, 98.8, 128.4, 128.7, 132.9, 133.6, 168.4, 173.8, 202.0 ppm. IR ( $\text{cm}^{-1}$ ): 3061, 1716, 1683, 1648, 1317. APCI:  $m/z$  calculated for  $\text{C}_{14}\text{H}_{13}\text{O}_3$  [ $\text{M} + \text{H}$ ] $^+$  = 303.1596; found: 303.1594.

**Methyl 6-(furan-2-carbonyl)-2-hydroxycyclohex-1-ene-1-carboxylate (8d).** Prepared following the general procedure (reaction time 2 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 102 mg (82%) of a light brown oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.60–1.67 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.83–1.92 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}$ ), 2.27–2.32 (m, 2H,

$\text{COHCH}_2\text{CH}_2$ ), 3.55 (s, 3H, OMe), 4.22–4.23 (m, 1H, CH), 6.51–6.53 (m, 1H, 4- $\text{C}_4\text{H}_3\text{O}$ ), 7.19–7.20 (m, 1H, 3- $\text{C}_4\text{H}_3\text{O}$ ), 7.57–7.59 (m, 1H, 5- $\text{C}_4\text{H}_3\text{O}$ ), 12.33 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 18.6, 26.4, 28.9, 41.8, 51.6, 96.5, 112.3, 117.3, 146.4, 147.1, 172.0, 174.9, 190.8 ppm. IR ( $\text{cm}^{-1}$ ): 3321, 1732, 1715, 1666, 1328. APCI:  $m/z$  calculated for  $\text{C}_{13}\text{H}_{15}\text{O}_5$  [ $\text{M} + \text{H}$ ] $^+$  = 251.0919; found: 251.0824.

***tert*-Butyl 2-(3-hydroxy-2-(methoxycarbonyl)cyclohex-2-ene-1-carbonyl)-1H-pyrrole-1-carboxylate (8e).** Prepared following the general procedure (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 70 : 30) to obtain 128 mg (70%) of a light brown oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.09 (t, 3H,  $J = 7.13$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.56 (s, 9H, *Ot*-Bu), 1.80–1.93 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.32–2.37 (m, 3H,  $\text{COHCH}_2\text{CH}_2$ ), 3.93–4.15 (m, 3H,  $\text{OCH}_2\text{CH}_3$ , CH), 6.16–6.20 (m, 1H, 4- $\text{C}_4\text{H}_3\text{N}$ ), 6.98–6.99 (m, 1H, 3- $\text{C}_4\text{H}_3\text{N}$ ), 7.37–7.38 (m, 1H, 5- $\text{C}_4\text{H}_3\text{N}$ ), 12.42 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.0, 18.8, 26.0, 27.7, 29.1, 50.8, 60.6, 84.9, 97.1, 109.7, 121.4, 128.7, 133.3, 149.2, 169.2, 174.7, 192.0 ppm. IR ( $\text{cm}^{-1}$ ): 3476, 1744, 1717, 1673, 1619, 1310. APCI:  $m/z$  calculated for  $\text{C}_{19}\text{H}_{25}\text{NO}_6$  [ $\text{M} + \text{H}$ ] $^+$  = 364.1760; found: 364.1737.

**Methyl 6-(thiophen-2-carbonyl)-2-hydroxycyclohex-1-ene-1-carboxylate (8f).** Prepared following the general procedure (reaction time 2 h). The residue was purified by column chromatography (hexanes/EtOAc 95 : 5) to obtain 97 mg (69%) of a yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 0.93 (t, 3H,  $J = 7.15$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.84–1.9 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.30–2.34 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 3.96–4.02 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 4.24–4.27 (m, 1H, CH), 7.10–7.12 (m, 1H, 4- $\text{C}_4\text{H}_3\text{S}$ ), 7.59–7.61 (m, 1H, 3- $\text{C}_4\text{H}_3\text{S}$ ), 7.62–7.77 (m, 1H, 5- $\text{C}_4\text{H}_3\text{S}$ ), 12.36 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 13.7, 19.0, 26.9, 29.0, 43.0, 60.5, 97.4, 128.1, 131.7, 133.4, 143.3, 171.6, 174.7, 195.0 ppm. IR ( $\text{cm}^{-1}$ ): 3091, 1737, 1715, 1652, 1333, 1216. APCI:  $m/z$  calculated for  $\text{C}_{14}\text{H}_{17}\text{O}_4\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  = 281.0848; found: 281.0838.

**Methyl 6-(2-bromobenzoyl)-2-hydroxycyclohex-1-ene-1-carboxylate (8g).** Prepared following the general procedure (reaction time 1 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 143 mg (84%) of a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.68–1.77 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.31–2.45 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 3.70 (s, 3H, OMe), 4.22–4.24 (m, 1H, CH), 7.26–7.31 (m, 1H, *m*- $\text{C}_6\text{H}_4\text{Br}$ ), 7.36–7.40 (m, 1H, *m*- $\text{C}_6\text{H}_4\text{Br}$ ), 7.52–7.54 (m, 1H, *p*- $\text{C}_6\text{H}_4\text{Br}$ ), 7.60–7.64 (m, 1H, *o*- $\text{C}_6\text{H}_4\text{Br}$ ), 12.36 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 18.5, 24.4, 29.0, 45.3, 51.7, 96.6, 119.9, 127.3, 128.5, 131.6, 134.1, 141.2, 172.1, 174.9, 203.6 ppm. IR ( $\text{cm}^{-1}$ ): 3440, 1742, 1702, 1655, 1333, 1087. APCI:  $m/z$  calculated for  $\text{C}_{15}\text{H}_{16}\text{BrO}_4$  [ $\text{M} + \text{H}$ ] $^+$  = 339.0232; found: 239.0232.

**Ethyl 6-(2-bromobenzoyl)-2-hydroxycyclohex-1-ene-1-carboxylate (8h).** Prepared following the general procedure (reaction time 5 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 153 mg (87%) of a colorless oil (mp = 92–94 °C, EtOAc).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.20 (t, 3H,  $J = 7.17$  Hz  $\text{OCH}_2\text{CH}_3$ ), 1.63–1.81 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.32–2.44 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 4.13–4.19 (m, 2H, 7.38) (m, 1H, *m*- $\text{C}_6\text{H}_4\text{Br}$ ), 7.54–7.56 (m, 1H, *p*- $\text{C}_6\text{H}_4\text{Br}$ ), 7.59–7.76 (m, 1H, *o*- $\text{C}_6\text{H}_4\text{Br}$ ), 12.46 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.2, 18.3, 24.1, 28.9, 45.2, 60.6, 96.5, 119.8,



127.1, 128.3, 131.5, 133.9, 141.1, 169.4, 174.8, 203.4 ppm. IR (cm<sup>-1</sup>): 3064, 1703, 1651, 1617, 1304, 1086. APCI: *m/z* calculated for C<sub>16</sub>H<sub>18</sub>BrO<sub>3</sub> [M + H]<sup>+</sup> = 353.0388; found: 353.0352.

**Methyl 2-hydroxy-6-(2-nitrobenzoyl)cyclohex-1-ene-1-carboxylate (8i).** Prepared following the general procedure (reaction time 30 min). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 132 mg (86%) of a light-yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.61–1.71 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.26–2.30 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.63 (s, 3H, OMe), 4.16–4.17 (m, 1H, CH), 7.20–7.24 (m, 1H, *m*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 7.30–7.33 (m, 1H *m*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 7.46–7.48 (m, 1H, *p*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 7.55–7.57 (m, 1H, *o*-C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>), 12.30 (bs, 1H, OH) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 18.5, 24.3, 28.9, 45.3, 51.6, 96.6, 119.8, 127.3, 128.4, 131.6, 134.0, 141.1, 172.0, 174.9, 203.6 ppm. IR (cm<sup>-1</sup>): 3001, 1742, 1702, 1657, 1440. APCI: *m/z* calculated for C<sub>15</sub>H<sub>16</sub>NO<sub>6</sub> [M – CO<sub>2</sub>Me]<sup>+</sup> = 247.0845; found: 247.0862.

**Methyl 2-hydroxy-6-(3-methoxybenzoyl)cyclohex-1-ene-1-carboxylate (8j).** Prepared following the general procedure (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 77 mg (50%) of a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 0.98 (t, 3H, *J* = 7.13 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.62–1.79 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.31–2.37 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.85 (s, 3H, OMe), 4.01–4.02 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 4.46–4.48 (m, 1H, CH), 7.08–7.11 (m, 1H, *o*-C<sub>6</sub>H<sub>4</sub>), 7.35–7.39 (m, 1H, *o*-C<sub>6</sub>H<sub>4</sub>), 7.49–7.50 (m, 1H, *m*-C<sub>6</sub>H<sub>4</sub>), 7.56–7.58 (m, 1H, *p*-C<sub>6</sub>H<sub>4</sub>), 12.42 (bs, 1H, OH) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 13.8, 18.6, 26.1, 28.8, 41.0, 55.4, 60.4, 97.5, 112.9, 119.0, 120.7, 129.5, 137.8, 159.8, 171.6, 174.3, 201.8 ppm. IR (cm<sup>-1</sup>): 3074, 1738, 1716, 1682, 1329. APCI: *m/z* calculated for C<sub>17</sub>H<sub>21</sub>O<sub>5</sub> [M + H]<sup>+</sup> = 305.1389; found: 305.1392.

**Ethyl 2-hydroxy-6-picolinoylcyclohex-1-ene-1-carboxylate (8k).** Prepared following the general procedure (reaction time 10 h). The residue was purified by column chromatography (hexanes/EtOAc 70 : 30) to obtain 84 mg (61%) of a brown oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 0.88 (t, 3H, *J* = 7.17 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.68–1.77 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.33–2.36 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.96–4.01 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 5.12–5.15 (m, 1H, CH), 7.47–7.49 (m, 1H, 5-C<sub>5</sub>H<sub>4</sub>N), 7.82–7.84 (m, 1H, 3-C<sub>5</sub>H<sub>4</sub>N), 8.05–8.06 (m, 1H, 4-C<sub>5</sub>H<sub>4</sub>N), 8.69–8.71 (m, 1H, 6-C<sub>5</sub>H<sub>4</sub>N), 12.35 (bs, 1H, OH) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 13.6, 19.0, 25.9, 29.0, 39.2, 60.2, 97.7, 122.3, 126.9, 136.9, 148.8, 152.7, 169.2, 174.2, 202.8 ppm. IR (cm<sup>-1</sup>): 3054.9, 1739, 1695, 1652, 1583, 1347. APCI: *m/z* calculated for C<sub>15</sub>H<sub>18</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 276.1236, found: 276.1200.

**Ethyl 2-hydroxy-6-nicotinoylcyclohex-1-ene-1-carboxylate (8l).** Prepared following the general procedure (reaction time 1.5 h). The residue was purified by column chromatography (hexanes/EtOAc 70 : 30) to obtain 105 mg (76%) of a light-yellow oil. <sup>1</sup>H NMR (400 MHz, DMSO) δ: 0.91 (t, 3H, *J* = 7.12 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.55–1.72 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.25–2.43 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.97–4.13 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 4.41–4.43 (m, 1H, CH), 7.22–7.39 (m, 1H, 5-C<sub>5</sub>H<sub>4</sub>N), 8.19–8.21 (m, 1H, 4-C<sub>5</sub>H<sub>4</sub>N), 8.72–8.73 (m, 1H, 6-C<sub>5</sub>H<sub>4</sub>N), 9.12–9.16 (m, 1H, 2-C<sub>5</sub>H<sub>4</sub>N), 12.29 (bs, 1H, OH) ppm. <sup>13</sup>C NMR (400 MHz, DMSO) δ: 14.0, 18.7, 25.9, 28.8, 48.4, 60.7, 97.6, 124.5, 131.5, 136.2, 149.7, 153.9, 171.5, 174.5, 201.7 ppm. IR (cm<sup>-1</sup>): 3048, 1737, 1716,

1784, 1584, 1331. APCI: *m/z* calculated for C<sub>15</sub>H<sub>18</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 276.1236; found: 276.1204.

**Methyl 2-hydroxy-6-(4-methoxybenzoyl)cyclohex-1-ene-1-carboxylate (8m).** Prepared following the general procedure (reaction time 12 h). The residue was purified by column chromatography (hexanes/EtOAc 90 : 10) to obtain 77 mg (53%) of a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.61–1.93 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.28–2.40 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.57 (s, 3H, CO<sub>2</sub>Me), 3.88 (s, 3H, OMe), 4.45–4.48 (m, 1H, CH), 6.93–6.98 (m, 2H, *m*-C<sub>6</sub>H<sub>4</sub>OMe), 7.94–8.00 (m, 2H, *o*-C<sub>6</sub>H<sub>4</sub>OMe), 12.35 (bs, 1H, OH) ppm. RMN <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 18.7, 26.6, 29.0, 40.7, 48.3, 58.7, 97.5, 113.9, 130.8, 163.5, 170.0, 174.6, 198.3 ppm. IR (cm<sup>-1</sup>): 3328, 1742, 1714, 1662, 1334. APCI: *m/z* calculated for C<sub>16</sub>H<sub>19</sub>NO<sub>5</sub> [M + H]<sup>+</sup> = 291.1232; found: 291.1251.

**Methyl 6-(6-bromobenzo[d][1,3]dioxole-5-carbonyl)-2-hydroxycyclohex-1-ene-1-carboxylate (8n).** Prepared following the general procedure (reaction time 3 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 150 mg (78%) of a white solid (mp = 92–94 °C AcOEt) <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.66–1.76 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.02–2.35 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.69 (s, 3H, OMe), 4.15–4.17 (m, 1H, CH), 6.03 (s, 2H, OCH<sub>2</sub>O), 7.03 (s, 1H, *m*-C<sub>6</sub>H<sub>2</sub>Br), 7.05 (s, 1H, *o*-C<sub>6</sub>H<sub>2</sub>Br), 12.32 (bs, 1H, OH) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 18.5, 24.6, 29.0, 45.0, 51.7, 96.7, 102.5, 108.5, 112.0, 114.2, 134.0, 147.3, 150.0, 172.0, 174.9, 202.5 ppm. IR (cm<sup>-1</sup>): 3384, 1716, 1699, 1662, 1328, 1037. APCI: *m/z* calculated for C<sub>16</sub>H<sub>16</sub>BrO<sub>6</sub> [M + H]<sup>+</sup> = 383.0130; found: 383.0155.

**tert-Butyl 2-(3-hydroxy-2-(methoxycarbonyl)cyclohex-2-ene-1-carbonyl)-1H-indole-1-carboxylate (8o).** Prepared following the general procedure (reaction time 6 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 123 mg (60%) of a brown oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.05 (t, 3H, *J* = 7.13 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.60 (s, 9H, *tert*-Bu), 1.89–1.99 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH), 2.34–2.37 (m, 2H, COHCH<sub>2</sub>CH<sub>2</sub>), 3.99–4.14 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 4.28–4.30 (m, 1H, CH), 7.23–7.27 (m, 1H, 5-C<sub>8</sub>H<sub>5</sub>N), 7.30 (s, 1H, 3-C<sub>8</sub>H<sub>5</sub>N), 7.37–7.44 (m, 1H, 6-C<sub>8</sub>H<sub>5</sub>N), 7.62–7.65 (m, 1H, 4-C<sub>8</sub>H<sub>5</sub>N), 7.98–8.01 (m, 1H, 7-C<sub>8</sub>H<sub>5</sub>N), 12.45 (bs, 1H, OH) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 14.0, 18.7, 25.8, 27.9, 29.0, 43.4, 60.5, 84.5, 96.7, 114.5, 115.7, 117.0, 122.5, 123.1, 127.3, 137.5, 138.9, 149.6, 169.2, 174.8, 193.5 ppm. IR (cm<sup>-1</sup>): 3053, 1736, 1679, 1652, 1613, 1321. APCI: *m/z* calculated for C<sub>15</sub>H<sub>18</sub>NO<sub>4</sub> [M + H]<sup>+</sup> = 414.1917; found: 414.1865.

**Methyl 2-benzoyl-5-oxocyclopentane-1-carboxylate (9a).** Prepared following the general procedure (reaction time 5 h). The residue was purified by column chromatography (hexanes/EtOAc 70 : 30) to obtain 99 mg (80%) of a translucent yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ: 1.94–1.99 (m, 1H, CH<sub>2</sub>CH<sub>2</sub>CH), 2.44–2.58 (m, 3H, COCH<sub>2</sub>CH<sub>2</sub>CH), 3.75 (s, 3H, OMe), 3.89–3.92 (d, 1H, *J* = 9.39 Hz, COCHCO), 4.50–5.57 (m, 1H, CH<sub>2</sub>CHCO), 7.49–7.60 (m, 2H, *m*-C<sub>6</sub>H<sub>5</sub>), 7.61–7.62 (m, 1H, *p*-C<sub>6</sub>H<sub>5</sub>), 8.02–8.05 (m, 2H, *o*-C<sub>6</sub>H<sub>5</sub>) ppm. <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ: 25.9, 37.9, 47.2, 53.0, 57.1, 128.8, 129.1, 134.0, 135.4, 168.6, 199.0, 209.1 ppm. IR (cm<sup>-1</sup>): 3053, 1748, 1724, 1678, 1321. APCI: *m/z* calculated for C<sub>13</sub>H<sub>11</sub>O<sub>3</sub> [M + H]<sup>+</sup> = 247.0970; found: 247.1008.

**Methyl 2-benzoyl-7-oxocycloheptane-1-carboxylate (9b).** Prepared following the general procedure (reaction time 5 h). The residue was purified by column chromatography (hexanes/



EtOAc 80 : 20) to obtain 68 mg (50%) of a pale-yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.49–1.62 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 1.95–2.70 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.63–2.73 (m, 2H,  $\text{COCH}_2\text{-CH}_2$ ), 3.64 (s, 3H, OMe), 4.01–4.03 (m, 1H,  $\text{COCHCO}$ ), 4.39–4.41 (d, 1H,  $J = 10.18$  Hz,  $\text{CH}_2\text{CHCO}$ ), 7.45–7.49 (m, 2H,  $m\text{-C}_6\text{H}_5$ ), 7.56–7.57 (m, 1H,  $p\text{-C}_6\text{H}_5$ ), 7.95–7.97 (m, 2H,  $o\text{-C}_6\text{H}_5$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 24.1, 26.9, 31.7, 43.2, 45.5, 52.6, 59.9, 128.6, 128.9, 133.5, 135.5, 170.0, 201.1, 207.6 ppm. IR ( $\text{cm}^{-1}$ ): 3061, 1742, 1702, 1679, 1322. APCI:  $m/z$  calculated for  $\text{C}_{16}\text{H}_{19}\text{O}_4$   $[\text{M} + \text{H}]^+ = 275.1283$ ; found: 275.1294.

**Methyl 6-benzoyl-2-hydroxy-6-methylcyclohex-1-ene-1-carboxylate (9c).** Prepared following the general procedure (reaction time 24 h). The residue was purified by column chromatography (hexanes/EtOAc 80 : 20) to obtain 52 mg (38%) of a yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.51 (s, 3H,  $\text{OCH}_3$ ), 1.70–1.74 (m, 1H,  $\text{CH}_2\text{CH}_2\text{C}$ ), 1.90–1.94 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.12–2.18 (m, 1H,  $\text{CH}_2\text{CH}_2\text{C}$ ), 2.48–2.54 (m, 2H,  $\text{COHCH}_2\text{CH}_2$ ), 3.39 (s, 3H, OMe), 7.32–7.36 (m, 2H,  $m\text{-C}_6\text{H}_5$ ), 7.40–7.45 (m, 1H,  $p\text{-C}_6\text{H}_5$ ), 7.81–7.83 (m, 2H,  $o\text{-C}_6\text{H}_5$ ), 12.61 (bs, 1H, OH) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 18.1, 24.9, 29.6, 34.8, 48.2, 51.1, 104.7, 128.3 ( $\times 2$ ), 131.7, 136.7, 172.1, 172.8, 203.9 ppm. 3056, 1717, 1678, 1650, 1308. APCI:  $m/z$  calculated for  $\text{C}_{15}\text{H}_{15}\text{O}_3$   $[\text{M} + \text{H}]^+ = 275.1283$ ; found: 275.1240.

**Ethyl 4-oxo-1-phenyl-3,4,5,6,7,7a-hexahydro-3aH-indene-3a-carboxylate (10).** DBU (109  $\mu\text{L}$ , 0.73 mmol) was added to a solution of **30** (100 mg, 0.37 mmol) in acetonitrile (3.7 mL) at 0  $^\circ\text{C}$  under nitrogen atmosphere. After stirring the mixture at this temperature for 15 min, triphenylvinylphosphonium bromide (202 mg, 546  $\mu\text{mol}$ ) was added portion wise. After the addition was completed, the mixture was stirring for 12 h at room temperature. The reaction was quenched by the addition of  $\text{H}_2\text{O}$  (10 mL) and extracted with EtOAc ( $3 \times 20$  mL); the organics were combined, dried (anh  $\text{Na}_2\text{SO}_4$ ) and concentrated *in vacuo*. The residue was purified by column chromatography ( $\text{SiO}_2$ ) (hexanes/EtOAc 70 : 30) to obtain 40 mg (39%) of translucent yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.25 (t, 3H,  $J = 3.17$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.56–1.78 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$ ), 2.11–2.14 (m, 1H,  $\text{COHCH}_2\text{CH}_2$ ), 2.46–2.50 (m, 1H,  $\text{COHCH}_2\text{CH}_2$ ), 2.52–2.54 (m, 1H,  $\text{CCH}_2\text{CH}$ ), 3.06–3.07 (m, 1H,  $\text{CCH}_2\text{CH}$ ), 3.984.00 (m, 1H, CH), 4.02–4.22 (m, 2H,  $\text{OCH}_2\text{CH}_3$ ), 5.93–5.95 (m, 1H,  $\text{CCHCH}_2$ ), 7.31–7.35 (m, 5H,  $\text{C}_6\text{H}_5$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.1, 21.5, 26.3, 38.3, 39.3, 51.5, 61.7, 65.8, 125.0, 126.3, 127.4, 128.4, 135.3, 144.4, 172.1, 208.7. IR ( $\text{cm}^{-1}$ ): 2945, 1709, 1676, 1315. APCI:  $m/z$  calculated for  $\text{C}_{18}\text{H}_{21}\text{O}_3$   $[\text{M} + \text{H}]^+ = 285.1491$ ; found: 285.1467.

**2-Acetyl-1-phenyl-3,4,5,6,7,7a-hexahydro-4H-inden-4-one (11).** DBU (82  $\mu\text{L}$ , 0.55 mmol) was added to a solution of **30** (100 mg, 0.37 mmol) in acetonitrile (1.2 mL) at 0  $^\circ\text{C}$  under nitrogen atmosphere. After stirring the mixture at this temperature for 15 min, methyl vinyl ketone (35  $\mu\text{L}$ , 401  $\mu\text{mol}$ ) was added dropwise. After the addition was completed, the mixture was stirring for 12 h at 80  $^\circ\text{C}$ . The reaction was quenched by the addition of  $\text{H}_2\text{O}$  (5 mL). The mixture was diluted with water and extracted with EtOAc ( $3 \times 20$  mL); the organics were combined, dried (anh  $\text{Na}_2\text{SO}_4$ ) and concentrated *in vacuo*. The residue was purified by column chromatography ( $\text{SiO}_2$ ) (hexanes/EtOAc 80 : 20) to obtain 82 mg (88%) of

translucent oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.40–1.44 (m, 1H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.62–1.74 (m, 1H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.75–1.77 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}$ ), 1.81 (s, 3H,  $\text{COCH}_3$ ), 2.36–2.39 (m, 2H,  $\text{COCH}_2\text{CH}_2$ ), 2.84–2.85 (m, 1H,  $\text{CHCH}_2\text{C}$ ), 2.97–2.99 (m, 1H,  $\text{CHCH}_2\text{C}$ ), 3.12–3.15 (m, 1H, CH), 3.48–3.50 (m, 1H, CH), 7.12–7.15 (m, 2H,  $m\text{-C}_6\text{H}_5$ ), 7.32–7.37 (m, 3H,  $o,p\text{-C}_6\text{H}_5$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 22.4, 26.2, 29.8, 35.3, 39.4, 48.7, 53.7, 127.8, 128.4, 128.6, 136.2, 138.4, 155.1, 198.8, 212.4 ppm. IR ( $\text{cm}^{-1}$ ): 1348, 1682, 1704, 1720, 2939. APCI:  $m/z$  calculated for  $\text{C}_{17}\text{H}_{19}\text{O}_2$   $[\text{M} + \text{H}]^+ = 255.1385$ ; found: 255.1379.

**8-Phenyl-2,3,3a,3b,5,6,7,7a-octahydrocyclopenta[a]indene-1,4-dione (12).** DBU (82  $\mu\text{L}$ , 0.55 mmol) was added to a solution of **30** (100 mg, 0.37 mmol) in acetonitrile (1.2 mL) at 0  $^\circ\text{C}$  under nitrogen atmosphere. After stirring the mixture at this temperature for 15 min, 2-cyclopenten-1-one (34  $\mu\text{L}$ , 401  $\mu\text{mol}$ ) was added dropwise. After the addition was completed, the mixture was stirring for 24 h at room temperature. The reaction was quenched by the addition of  $\text{H}_2\text{O}$  (5 mL). The mixture was diluted with water and extracted with EtOAc ( $3 \times 20$  mL); the organics were combined, dried (anh  $\text{Na}_2\text{SO}_4$ ) and concentrated *in vacuo*. The residue was purified by column chromatography ( $\text{SiO}_2$ ) (hexanes/EtOAc 80 : 20) to obtain 31 mg (32%) of translucent oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 1.51–1.61 (m, 1H), 1.71–1.80 (m, 2H), 2.02–2.10 (m, 2H), 2.24–2.37 (m, 1H), 2.47–2.50 (m, 2H), 2.61–2.65 (m, 2H), 2.84–2.89 (m, 1H), 3.63–3.70 (m, 1H, CH), 3.88–3.94 (m, 1H, CH), 7.39–7.43 (m, 2H,  $m\text{-C}_6\text{H}_5$ ), 7.99–8.02 (m, 3H,  $o,p\text{-C}_6\text{H}_5$ ) ppm.  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 24.0, 27.8, 28.9, 39.1, 44.7, 53.3, 54.9, 59.3, 128.5, 129.5, 130.5, 132.1, 137.6, 151.2, 200.5, 212.3 ppm. IR ( $\text{cm}^{-1}$ ): 1335, 1590, 1686, 2951. APCI:  $m/z$  calculated for  $\text{C}_{17}\text{H}_{19}\text{O}_2$   $[\text{M} + \text{H}]^+ = 267.1385$ ; found: 267.1349.

## Conclusions

We utilized a cascade process to obtain tricarbonyl compounds by adding anions of cyanocarboxylates derived from aromatic aldehydes onto 5, 6, and 7-membered cycloalkenones. During the process, we exploited the dual role of cyanocarboxylates: as “latent” acylcarbanions and as acylating reagents. The synthetic potential of the cascade products obtained was successfully demonstrated by forming bicyclic and tricyclic systems through intramolecular condensation reactions. These results suggest that tricarbonyl compounds can be used as scaffolds to generate structural diversity. The investigation of further transformations is currently underway and will be published in due course.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

The authors thank Facultad de Química, UNAM for financial support (Grant PAIP 50009062). We thank Professor D. E. Ward for revising the manuscript, and for helpful suggestions. We also thank Rosa Isela del Villar, Nayeli López, Georgina Duarte,

and Marisela Gutierrez for recording NMR, MS, and IR spectra. DM thanks CONACYT for Graduate Scholarship.

## Notes and references

- 1 G. Stork and L. A. Maldonado, *J. Am. Chem. Soc.*, 1971, **93**, 5286–5287.
- 2 R. J. H. Gregory, *Chem. Rev.*, 1999, **99**, 3649–3682.
- 3 J. D. Albright, *Tetrahedron*, 1983, **39**, 3207–3233.
- 4 (a) G. Stork and L. Maldonado, *J. Am. Chem. Soc.*, 1974, **96**, 5273–5274; (b) G. Stork, J. C. Depezay and J. D'Angelo, *Tetrahedron Lett.*, 1975, 389–392; (c) J. Tsuji, *J. Am. Chem. Soc.*, 1981, **103**, 5259–5266; (d) G. Stork and T. Takahashi, *J. Am. Chem. Soc.*, 1977, **99**, 1275–1276; (e) G. Stork, T. Takahashi, I. Kawamoto and T. Suzuki, *J. Am. Chem. Soc.*, 1978, **100**, 8272–8273; (f) T. Takahashi, K. Kitamura, H. Nemoto, J. Tsuji and I. Miura, *Tetrahedron Lett.*, 1983, **24**, 3489–3492; (g) T. Takahashi, H. Iwamoto, K. Nagashima, T. Okabe and T. Doi, *Angew. Chem., Int. Ed.*, 1997, **36**, 1319–1321; (h) O. S. Park, H. J. Hwang and W. Y. Lee, *Arch. Pharmacol. Res.*, 1993, **16**, 205–208; (i) A. Kende, K. Liu, I. Kaldor, G. Dorey and K. Koch, *J. Am. Chem. Soc.*, 1995, **117**, 8258–8270; (j) I. Kadota, Y. Hu, G. K. Packard and S. D. Rychnovsky, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**(33), 11992–11995; (k) E. H. Granados-Covarrubias and L. A. Maldonado, *J. Org. Chem.*, 2009, **74**, 5097–5099.
- 5 (a) A. T. Au, *Synth. Commun.*, 1984, **14**, 743–748, For recent methodologies, see: (b) H. M. Torres-Domínguez, L. A. Maldonado and R. Le Lagadec, *Tetrahedron Lett.*, 2020, **61**, 151414–151417; (c) S. Aoki, S. Kotani, M. Sugiura and M. Nakajima, *Tetrahedron Lett.*, 2010, **51**, 3547–3549.
- 6 (a) L. F. Tietze and A. Modi, *Med. Res. Rev.*, 2000, **20**(4), 304–322; (b) K. C. Nicolaou, D. J. Edmonds and P. G. Bulger, *Angew. Chem., Int. Ed.*, 2006, **45**, 7134–7186; (c) K. C. Nicolaou and J. S. Chen, *Chem. Soc. Rev.*, 2009, **38**, 2993–3009; (d) J. Poulin, C. M. Grisé-Bard and L. Barriault, *Chem. Soc. Rev.*, 2009, **38**, 3092–3101; (e) C. Grondal, M. Jeanty and D. Enders, *Nat. Chem.*, 2010, **2**, 167–178; (f) H. Pellissier, *Chem. Rev.*, 2013, **113**(1), 442–524; (g) S. Zhi, S. Sanjun, X. Ma and W. Zhang, *Org. Biomol. Chem.*, 2019, **17**, 7632–7650.
- 7 H. M. Torres-Domínguez, L. A. Maldonado and R. Le Lagadec, *Synth. Commun.*, 2017, **47**, 1250–1255.
- 8 (a) A. Sharma, J. Pandey and R. P. Tripathi, *Tetrahedron Lett.*, 2009, **50**, 1812–1816; (b) T. Venkatesh, P. S. Mainkar and S. Chandrasekhar, *J. Org. Chem.*, 2021, **86**, 5412–5416; (c) K. Asahi and H. Nishino, *Tetrahedron*, 2008, **64**, 1620–1634; (d) S. R. Lima and F. Coelho, *ACS Omega*, 2020, **5**, 8032–8045, and references therein.
- 9 (a) F. W. Sum and L. Weiler, *Tetrahedron*, 1981, **37**, 303–317; (b) B. Lygo, N. Oconnor and P. R. Wilson, *Tetrahedron*, 1988, **44**, 688–6888; (c) S. Benetti, R. Romagnoli, C. De Risi, G. Spalluto and V. Zanirato, *Chem. Rev.*, 1995, **95**, 1065–1114; (d) C. Simon, C. Thierry and J. Jean Rodriguez, *Eur. J. Org. Chem.*, 2004, 4957–4980; (e) D. Bonne, Y. Coquerel, T. Constantieux and J. Rodriguez, *Tetrahedron: Asymmetry*, 2010, **21**, 1085–1109; (f) T. Govender, P. I. Arvidsson, G. E. M. Maguire, H. G. Kruger and T. Naicker, *Chem. Rev.*, 2016, **116**, 9375–9437; (g) Y. S. Rao, *Chem. Rev.*, 1976, **76**, 625–694.
- 10 (a) V. Bhardwaj, D. Gumber, V. Abbot, S. Dhiman and P. Sharma, *RSC Adv.*, 2015, **5**, 15233–15266; (b) S. Khaghaninejad and M. M. Heravi, *Adv. Heterocycl. Chem.*, 2014, **111**, 95–146; (c) T. J. Donohoe and R. D. C. Pullin, *Chem. Commun.*, 2012, **48**, 11924–11938; (d) C. Schmuck and D. Rupprecht, *Synthesis*, 2007, 3095–3110.
- 11 S. P. Kolis, M. T. Clayton, J. L. Grutsch and M. M. Faul, *Tetrahedron Lett.*, 2003, **44**, 5707–5710.
- 12 B. Thapa and H. Schlegel, *J. Phys. Chem. A*, 2016, **120**, 5726–5735.
- 13 J. D. Chai and M. Head-Gordon, *Phys. Chem. Chem. Phys.*, 2008, **10**, 6615–6620.
- 14 S. J. Grimme, *Comput. Chem.*, 2006, **27**, 1787–1799.
- 15 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr, J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16*, Gaussian, Inc., Wallingford CT, 2016.
- 16 A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2009, **113**, 6378–6396.
- 17 (a) S. Dhers, A. Mondal, D. Aguilà, J. Ramírez, S. Vela, P. Dechambenoit, M. Rouzières, J. R. Nitschke, R. Clérac and J. M. Lehn, *J. Am. Chem. Soc.*, 2018, **140**, 8218–8227; (b) M. A. Rosero-Mafla, J. I. Castro, N. E. Sánchez, C. A. Mujica-Martinez and M. N. Chaur, *ChemistrySelect*, 2020, **5**, 7685–7694.
- 18 (a) C. D. Johnson, *The Hammett Equation*, Cambridge University Press, Cambridge, 1973; (b) C. Hansch, A. Leo and R. Taft, *Chem. Rev.*, 1991, **91**, 165–195.
- 19 A. Kuźnik, R. Mazurkiewicz and B. Fryczkowska, *Beilstein J. Org. Chem.*, 2017, **13**, 2710–2738.
- 20 A. Srikrishn and D. H. Dethe, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 2012, **51**(2), 345–355.
- 21 P. Klahn, A. Duschek, C. Liébert and S. F. Kirsch, *Org. Lett.*, 2012, **14**, 1250–1253.
- 22 A. E. Cotman, B. Modéc and B. Mohar, *Org. Lett.*, 2018, **20**, 2921–2924.
- 23 C. Zhang, H. Jiang and S. Zhu, *Chem. Commun.*, 2017, **53**, 2677–2680.