

RESEARCH ARTICLE

Characterization of recombinant β -fructofuranosidase from *Bifidobacterium adolescentis* G1

Toshima Omori, Keiji Ueno, Kei Muramatsu, Masanori Kikuchi, Shuichi Onodera and Norio Shiomi*

Abstract

Background: We have previously reported on purification and characterization of β -fructofuranosidase (β -FFase) from *Bifidobacterium adolescentis* G1. This enzyme showed high activity of hydrolysis on fructo-oligosaccharides with a low degree of polymerization. Recently, genome sequences of *B. longum* NCC2705 and *B. adolescentis* ATCC 15703 were determined, and *cscA* gene in the both genome sequences encoding β -FFase was predicted. Here, cloning of *cscA* gene encoding putative β -FFase from *B. adolescentis* G1, its expression in *E. coli* and properties of the recombinant protein are described.

Results: Using the information of *cscA* gene from *Bifidobacterium adolescentis* ATCC 15703, *cscA* gene from *B. adolescentis* G1 was cloned and sequenced. The N-terminal amino acid sequence of purified β -FFase from *B. adolescentis* G1 was identical to the deduced amino acid sequences of *cscA* gene from *B. adolescentis* G1. To confirm the translated product of the *cscA* gene, the recombinant protein was expressed in *Escherichia coli*. Molecular mass of the purified recombinant enzyme was estimated to be about 66,000 by SDS-PAGE and 60,300 by MALDI TOF-MS. The optimum pH of the enzyme was 5.7 and the enzyme was stable at pH 5.0-8.6. The thermostability of the enzyme was up to 50°C. The K_m (mM), V_{max} ($\mu\text{mol}/\text{mg}$ of protein/min), k_0 (sec^{-1}) and k_0/K_m ($\text{mM}^{-1} \text{sec}^{-1}$) for 1-kestose, neokestose, nystose, fructosylnystose, sucrose and inulin were 1.7, 107, 107.5, 63.2, and 1.7, 142, 142.7, 83.9, and 3.9, 152, 152.8, 39.2, and 2.2, 75, 75.4, 34.3, and 38, 79, 79.4, 2.1, and 25.9, 77, 77.4, 3.0, respectively. The hydrolytic activity was strongly inhibited by AgNO_3 , SDS, and HgCl_2 .

Conclusion: The recombinant enzyme had similar specificity to the native enzyme, high affinity for 1-kestose, and low affinity for sucrose and inulin, although properties of the recombinant enzyme showed slight difference from those of the native one previously described.

Background

Bifidobacteria are saccharolytic anaerobes generally present in human intestine. Growth of bifidobacteria is selectively promoted by prebiotics [1].

Fructo-oligosaccharides, such as 1-kestose, nystose and fructosylnystose, consist of β -2,1-linked fructose to sucrose, and they are naturally contained in artichoke tubers [2], chicory roots [3] and burdock roots [4,5].

These saccharides have been produced and commercially manufactured from sucrose with bacterial fructosyltransferase [6] and β -fructofuranosidases (β -FFases) [7-9], and have been on the market as prebiotics. Fructo-oligosaccharides are not hydrolyzed by digestive enzymes of mammalian origin, so they are able to reach large intestine, and to be selectively degraded by the resident microbes, such as bifidobacteria.

We have already reported that *B. adolescentis* G1 were isolated from feces of human adults, and produce the unique β -FFase which has high affinity toward 1-kestose, nystose and fructosylnystose [10-12]. Recently, genome sequences of *B. longum* NCC2705 (accession no.

* Correspondence: n-shiomi@rakuno.ac.jp

¹ Department of Food and Nutrition Sciences, Graduate School of Dairy Science Research, Rakuno Gakuen University, 582 Bunkyo-dai Midorimachi, Ebetsu, 069-8501, Japan

Full list of author information is available at the end of the article

AE014295) and *B. adolescentis* ATCC 15703 (AP009256) were determined, and *cscA* gene in the both genome sequences encoding β -FFase was predicted [13]. The recombinant β -FFase from *B. infantis* [14], *B. lactis* [15] and *B. longum* [16] have been studied using mixtures of fructo-oligosaccharides (Actilight, Raftilose and Raftiline) as a substrate, although detailed substrate specificity of the enzyme to sole fructo-oligosaccharide remains unclear. In our previous study purification and the substrate specificity of β -FFase from *B. adolescentis* G1 was demonstrated [10,11]. However, we have not revealed the gene encoding β -FFase from *B. adolescentis* G1 yet. This study is aimed at cloning of *cscA* gene from *B. adolescentis* G1 and characterizing the recombinant protein of *cscA* gene expressed in *Escherichia coli*.

Results and discussion

Deduced amino acid sequence of *cscA* gene from *B. adolescentis* G1

The open reading frame (ORF) of *cscA* gene from *B. adolescentis* G1 consisted of 1,557 nucleotides which encoded a 518 amino acid polypeptide. The molecular mass and pI of the deduced polypeptide were predicted to be 58,119 and 4.89, respectively. N-terminal sequence of purified enzyme was found in the deduced polypeptide. The deduced polypeptide of *cscA* gene from *B. adolescentis* G1 shows high identity that from bifidobacteria (*B. adolescentis* ATCC15703 [accession no. AP009256, protein ID BAF39931]: 98%, *B. longum* NCC2705 [accession no. AE014295, protein ID AAN23970]: 85%, *B. breve* UCC2003 [accession no. AY549965, protein ID AAT28190]: 84%, *B. animalis* subsp. *lactis* DSM10140 [accession no. AY509036, protein ID AAS87041]: 72%) (Fig. 1). Deduced polypeptides of *cscA* gene from *B. adolescentis* G1 belong to GH family 32, and confirmed existence of NDPNG motif and EC motif involved in catalysis. Two acidic residues have been identified as being related in the catalytic mechanism of yeast invertase: Asp-23 as a nucleophile and Glu-204 as an acid/base catalyst [17,18]. Asp within NDPNG motif and Glu within EC motif were significant part of β -FFase from *B. adolescentis* G1 (position of NDPNG: Asn-53-Gly-57, EC: Glu-235-Cys-236). Additionally, Asp-309 within RDP motif was investigated on *Acetobacter diazotrophicus* levansucrase, the motif was found at a conserved position, not only in levanses but also in invertase [19]. The Asp (D) in RDP motif in levansucrase from *Bacillus subtilis* was identified as a transition state stabilizer [20]. The existence of the motif was likely to keep a common functional role for β -FFase. It seems that the motif was also important portion for β -FFase from *B. adolescentis* G1 (position of RDP: Arg-180-Pro-182). The motifs are in highly conserved position in β -fructofuranosidase of GH32 [21].

Heterologous expression and purification of recombinant protein

To investigate translated product of *cscA* gene from *B. adolescentis* G1 in detail, recombinant protein was produced by *E. coli* Rosetta2 (DE3) with pET-G1cscA. The recombinant protein was confirmed in cell extract by SDS-PAGE and enzyme assay. The recombinant protein was purified from cell extract (total protein 163 mg, total enzyme activity 1,480 U, specific activity 9.1 U/mg) (Fig. 2(a)) by TALON CellThru Resin, DEAE Sepharose CL-6B, Toyopearl HW-55S. Purification procedures of recombinant protein were summarized in Table 1. Specific activity of the final purified enzyme was 86.0 U/mg. SDS-PAGE analysis of the purified enzyme showed a single band with an apparent molecular mass of about 66,000 (Fig. 2(b)). MALDI-TOF mass spectrum of purified recombinant protein shows a mass-peak at near 60,300, in agreement with the calculated mass of recombinant protein including extra sequences from pET32b (+) vector (data not shown). The reason of difference between calculated mass and actual measurement mass by SDS-PAGE is not clear. However, Shirai *et al* reported that acidic proteins (those for which isoelectric points are less than 6.0) exhibited mobility smaller than calculated, this may be due to negative charge repulsion with SDS [22]. In case of our study, the smaller mobility of recombinant protein (Fig. 2) than the mobility from calculated mass might be thought by the same reason.

Effects of pH and temperature


The effects of pH on hydrolytic activities toward 1-kestose were examined. The optimum pH of recombinant protein was observed to be 5.7 (Fig. 3). The characterization is similar to other β -FFase from bifidobacteria [14] that enzyme reaction proceeds strongly at acidulant condition. The recombinant protein was stable in the pH range from 5.0 to 8.6 (residual activity more than 90%) (Fig. 4). Although this extend of pH-stability is slightly different from the native enzyme [10], it is not poles apart. The enzyme was stable up to 50°C (residual activity more than 90%), and inactivated at 60°C (Fig. 5). This property is also similar to other bifidobacterium β -FFase [14].

Effects of various metal salts and chemical reagents

The hydrolase activity of the enzyme was not inhibited by MgCl₂, FeCl₃, CoCl₂, MnCl₂, SnCl₂ and CaCl₂, deoxycholic acid sodium salt, TritonX-100, Tween80 and ZnCl₂ had enhanced the enzyme activity. The enzyme activity was strongly inhibited by AgNO₃, SDS, HgCl₂, and was partially inhibited by CuSO₄. The result was summarized in Table 2. They formed mercaptides with free SH groups, so inhibited the hydrolysis of 1-kestose, as described previously [10].


| | |
|----------------------|--|
| B. adoG1_CscA | -----MTGFTPDAPVLHEIKNHGEE ^T KAEGVAAFAAKRNNRWYPKFHIAS |
| B. ado_CscA | -----MTGFTPDAPVLHEIKNHSEEL ^T KAEGVAAFAAKRNNRWYPKFHIAS |
| B. lon_CscA | -----MTDFTPETPVLTP ^I HDHAAELAKAEAGVAEMAANRNNRWYPKYHIAS |
| B. bre_CscA | -----MTDFTPETPVLTP ^I RDHAAELAKAEAGVAEMAANRNNRWYPKYHIAS |
| B. lac_CscA | MATLPTNIPANGILTPD ^L DPVLT ^P ISDHAEQLSLAEAGVSALETRNRNRWYPKFHIAS :* .. *** *: *: *: *: *: *: *: *: *: *: *: * |
| B. adoG1_CscA | NGGWINDPNGLCFYKGRWHV ^F YQLHPYGTQWGP ^M HGHVSSTDMVNWKREPIMFAPSLEE |
| B. ado_CscA | NGGWINDPNGLCFYKGRWHV ^F YQLHPYGTQWGP ^M HGHVSSTADMVNWKREPIMFAPSLEE |
| B. lon_CscA | NGGWINDPNGLCFYKGRWHV ^F YQLHPYGTQWGP ^M HGHVSSTDMLNWKREPIMFAPSLEQ |
| B. bre_CscA | NGGWINDPNGLCFYKGRWHV ^F YQLHPYGTQWGP ^M HGHVSSTDMLNWKREPIMFAPSLEQ |
| B. lac_CscA | NGGWINDPNGLCRYNGRWHV ^F YQLHPHG ^T QWGP ^M HGHVSSDNMVDWHREP ^I AFAPSLEQ ***** * : ***** : ***** : : * : * : * * * : * * * : * |
| B. adoG1_CscA | EKDG ^V FSGSAVIGDDGK ^L RFY ^T GHRWANGK ^D NTGGDW ^V QMLAEPD ^N DELTSAT ^K RG ^M V |
| B. ado_CscA | EKDG ^V FSGSAVIGDDGK ^L RFY ^T GHRWANGK ^D NTGGDW ^V QMLAEPD ^N DELTSAT ^K RG ^M V |
| B. lon_CscA | EKDG ^V FSGSAVIDDNGD ^L RFY ^T GHRWANGH ^D NTGGDW ^V QMTAL ^P DNDDELTSAT ^K QGM ^I |
| B. bre_CscA | EKDG ^V FSGSAVIDDNGD ^L RFY ^T GHRWANGH ^D NTGGDW ^V QMTAL ^P DNDDELTSAT ^K QGM ^I |
| B. lac_CscA | ERHG ^V FSGSAVIGDDGK ^P WI ^F YTGHRWANGK ^D NTGGDW ^V QMLAKPNDD ^L K ^T F ^T KEGM ^I * : ***** * : * : : ***** : ***** * * : * : * : * : * * : * |
| B. adoG1_CscA | IDCPTDKVNH ^H YRDPK ^V WKTGD ^K WYMTFGV ^S SAEK ^R GQMWLFSSDDM ^V KW ^T YEQVL ^F FEHP |
| B. ado_CscA | IDCPTDKVNH ^H YRDPK ^V WKTGD ^K WYMTFGV ^S SAEK ^R GQMWLFSSDDM ^V KW ^T YEQVL ^F FEHP |
| B. lon_CscA | IDCPTDKVDH ^H YRDPK ^V WKTGD ^T WYMTFGV ^S SADK ^R GQMWLFSSKDM ^V RWE ^Y ERVLF ^Q HP |
| B. bre_CscA | IDCPTDKVDH ^H YRDPK ^V WKTGD ^T WYMTFGV ^S SEDK ^R GQMWLFSSKDM ^V RWE ^Y ERVLF ^Q HP |
| B. lac_CscA | IDCPTDEVDH ^H FRDPK ^V WKTGD ^T WYMTFGV ^S SKEH ^R GQMWLYTSSDM ^V HWSFDR ^V LF ^F HP ***** * : * : * : ***** * : ***** : : ***** : * : * : * * : : * : * : * |
| B. adoG1_CscA | DPDV ^F MLECDP ^F FPIK ^D VGNEK ^W IGFSAMG ^A QPSGFMN ^R NV ^N NAGYMI ^G TWT ^P GEQ ^F K |
| B. ado_CscA | DSNV ^F MLECDP ^F FPIK ^D VEGNEK ^W IGFSAMG ^A KPSGFMN ^R NV ^N NAGYMI ^G TWT ^P GEQ ^F K |
| B. lon_CscA | DPDV ^F MLECDP ^F FPIK ^D KGNEK ^W IGFSAMG ^S KPSGFMN ^R VS ^N NAGYMI ^G TWE ^P GGEF ^K |
| B. bre_CscA | DPDV ^F MLECDP ^F FPIK ^D KGNEK ^W IGFSAMG ^S KPSGFMN ^R NV ^N NAGYMI ^G TWE ^P GGEF ^K |
| B. lac_CscA | DPNV ^F MLECDP ^F FPIR ^D ARGNEK ^W IGFSAMG ^A KPN ^G FMN ^R NV ^N NAGY ^M VGT ^W K ^P GES ^F K * : * : ***** * : ***** * : * : ***** * : * : * : * : * : * : * |
| B. adoG1_CscA | PETE ^F R ^L WDCGH ^N YAP ^Q SFN ^D GK ^R QI ^V YG ^W MSPF ^V EPI ^M QD ^D GWCG ^N L ^T L ^P REI ^T LGA |
| B. ado_CscA | PETE ^F R ^L WDCGH ^N YAP ^Q SFN ^D GK ^R QI ^V YG ^W MSPF ^V EPI ^M QD ^D GWCG ^N L ^T L ^P REI ^T LGA |
| B. lon_CscA | PETE ^F R ^L WDCGH ^N YAP ^Q SFN ^V DGR ^Q I ^V YG ^W MSPF ^V Q ^P I ^P ME ^D DGW ^C G ^Q L ^T L ^P REI ^T LDG |
| B. bre_CscA | PETE ^F R ^L WDCGH ^N YAP ^Q SFN ^V DGR ^Q I ^V YG ^W MSPF ^V Q ^P I ^P ME ^D DGW ^C G ^Q L ^T L ^P REI ^T LDD |
| B. lac_CscA | PETE ^F R ^L WDEGH ^N FYAP ^Q SFN ^T EG ^R Q ^I M ^Y GW ^M S ^P F ^V A ^P I ^P ME ^E DGW ^C G ^N L ^T L ^P REI ^T LDG ***** * : * : ***** * : ***** * : * : ***** * : * : * : * : * : * |
| B. adoG1_CscA | DGDL ^H TAPVAEMEGLRE ^D TVD ^F GAIDL ^D VS ^G EKT ^I VDD-AEAVEIEM ^T IDLAN ^S TAERAG |
| B. ado_CscA | DGDL ^H TAPVAEMEGLRE ^D TVD ^F GAIDL ^D VS ^G EKT ^I VDD-AEAVEIEM ^T IDLAN ^S TAERAG |
| B. lon_CscA | DGDV ^V TAPVAEMEGLRE ^D TLD ^H GSIT ^L DM ^D GE ^Q VIADD-AEAVEIEM ^T IDLAA ^S TAERAG |
| B. bre_CscA | DGDV ^V TAPVAEMEGLRE ^D TLD ^H GSIT ^L DM ^D GE ^Q VIADD-AEAVEIEM ^T IDLAA ^S TADRAG |
| B. lac_CscA | DGDL ^V TAPT ^I EMEGLRE ^N TIG ^F DSL ^D LGT ^N QT ^S TIL ^D DDGGALEIEM ^R LD ^L NK ^T TAERAG * : * : * : ***** : * : * : * : * : * : * : * : * : * : * : * |
| B. adoG1_CscA | LCV ^H ATEDGAYTSVAYD ^D QIGR ^V V ^V DRQANA ^Q GDRGYRTAP ^L SEAE ^L LAGEL ^K LRV ^Y VDR |
| B. ado_CscA | LRV ^H ATEDGAYTSVAYD ^D QIGR ^V V ^V DRQANA ^Q GDRGYRTAP ^L SEAE ^L LAGEL ^K LRV ^Y VDR |
| B. lon_CscA | LKI ^H ATEDGAYTYVAYD ^D QIGR ^V V ^V DRQAMANG ^D RGYRAAP ^L TDAELASG ^L DL ^R V ^F VDR |
| B. bre_CscA | LKI ^H ATEDGAYTYVAYD ^D QIGR ^V V ^V DRQAMANG ^D HGYRAAP ^L TDAELASG ^L DL ^R V ^F VDR |
| B. lac_CscA | LHV ^H ATSDGHYTAIV ^F DAQIG ^G V ^I DR ^Q NVANG ^D KGYR ^V AKLS ^D TE ^L AAD ^L DL ^R V ^F IDR * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * |
| B. adoG1_CscA | GCVE ^V YVNDGR ^Q V ^L SS ^S YASEG ^P RAIK ^L VAES ^G TLK ^V SLV ^L LH ^H M ^K SIGLE |
| B. ado_CscA | GCVE ^V YVNDGR ^Q V ^L SS ^S YASEG ^P RAIK ^L VAES ^G TLK ^V SLV ^L LH ^H M ^K SIGLE |
| B. lon_CscA | GSVE ^V YVNGGH ^Q V ^L SS ^S YASEG ^P RAIK ^L VAES ^G SLK ^V DSL ^K LH ^H M ^K SIGLE |
| B. bre_CscA | GSVE ^V YVNGGH ^Q V ^L SS ^S YASEG ^P RAIK ^L VAEF ^G N ^L K ^V ES ^L KLH ^H M ^K SIGLE |
| B. lac_CscA | GCVE ^V YV ^D G ^G KHAM ^S YSF ^P DGARAV ^E L ^V SE ^G T ^H I ^D LT ^M H ^S L ^K SIGLE * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * : * |

Figure 1 Deduced amino acid sequence of *cscA* from *Bifidobacterium*. B.adoG1_CscA, deduced amino acid sequence of *cscA* gene from *B. adolescentis* G1; B.ado_CscA, *B. adolescentis* ATCC 15703 (A1A2J8); B.lon_CscA, *B. longum* NCC2705 (Q8CY57); B.bre_CscA, *B. breve* UCC 2003 (3115295E); B.lac_CscA, *B. lactis* (Q564C7) are aligned. The three carboxylic acids in NDPNG, RDP and EC motifs are crucial for enzyme activity are shaded in gray. Underlined sequences indicate the N-terminus of purified β -FFase from *B. adolescentis* G1. Consensus line: asterisks (*) indicated identical residues; colons (:) indicated conserved substitutions; and periods (.) indicated semi-conserved substitutions.



Substrate specificity and kinetic parameters

To examine the substrate specificity of the recombinant β -FFase, the enzyme was incubated in 10 mM fructo-oligosaccharides such as 1-kestose, neokestose, nystose or fructosylnystose at 37°C for 0, 10 min, 1 and 4 h. Reaction products were analyzed by HPAEC (Fig. 6). This enzyme catalyzed hydrolytic reaction on the fructo-oligosaccha-



rides. In the reaction of 1-kestose and neokestose, fructose and sucrose were produced (Fig. 6(a) and 6(b)). These results indicate that the enzyme hydrolyzed not only terminal β -2,1 fructoside bond of 1-kestose, but also β -2,6 fructoside bond (fru-2,6-glc) of neokestose. When the enzyme was incubated with nystose, fructose and 1-kestose were produced (Fig. 6(c)), with fructosylnystose, fructose and nystose were produced (Fig. 6(d)), with sucrose, glucose and fructose were produced (Fig. 6(e)), with inulin, fructose was produced (Fig. 6(f)). By the prolonged incubation of nystose or fructosylnystose, sucrose and 1-kestose were also produced. This result showed that the enzyme is capable of degrading each substrate via an exo-type of cleavage, releasing terminal fructosyl residues as well as other β -FFase reported from bifidobacteria [14,15]. To estimate the rate parameters of hydrolysis of several substrates, Michaelis constants (K_m) and maximum velocities (V_{max}) for each substrate were measured.

Table 1: Summary of purification procedure recombinant β -FFase.

| Procedure | Total protein (mg) | Total activity (units) | Specific activity (units/mg) | Purification (-fold) | Recovery (%) |
|-----------------------|--------------------|------------------------|------------------------------|----------------------|--------------|
| Crude extract | 163 | 1,480 | 9.08 | 1.00 | 100 |
| Metal affinity resins | 30.0 | 600 | 20.0 | 2.20 | 40.5 |
| DEAE-Sepharose CL-6B | 11.3 | 900 | 79.7 | 8.78 | 60.8 |
| Toyopearl HW-55S | 3.65 | 314 | 86.0 | 9.47 | 21.2 |




Figure 4 pH Stability curve of recombinant β -FFase. After preincubation of enzyme solutions containing 0.1% BSA in Britton-Robinson buffer with different pHs (3.95, 4.25, 4.53, 4.97, 5.67, 6.30, 6.58, 6.89, 7.23, 7.60, 7.92, 8.28, 8.64, 9.13 or 9.53) for 20 h at 4°C, the mixture was adjusted to pH 5.7, and then, residual activities were measured. Residual activities indicated closed squares.

K_m , V_{max} , k_0 , k_0/K_m are listed in Table 3. These parameters indicated that the recombinant enzyme had high affinity and high rate for hydrolyzing FOS, especially for 1-kestose, neokestose and nystose. The hydrolyzing activity against sucrose and inulin was lower than formers, so the enzyme was thought mainly as relating FOS degradation.

The recombinant protein indicated a higher relative efficiency for fructo-oligosaccharides such as 1-kestose,





Figure 5 Temperature stability curve of recombinant β -FFase. Enzyme solution containing 0.1% BSA was incubated at different temperature (4, 30, 40, 45, 50, 55, 60 or 65°C) for 15 min, cooled to 0°C, and then, residual activities were measured. Residual activities indicated closed squares.

Table 2: Effects of various metal salts or chemical reagents on recombinant β -FFase.

| Reagent | Relative activity (%) |
|------------------------------|-----------------------|
| None | 100 |
| AgNO ₃ | 0 |
| CuSO ₄ | 51.8 |
| MgCl ₂ | 99.1 |
| FeCl ₃ | 110.6 |
| CoCl ₂ | 105.1 |
| MnCl ₂ | 100.2 |
| SnCl ₂ | 99.2 |
| ZnCl ₂ | 121.1 |
| CaCl ₂ | 102.1 |
| HgCl ₂ | 0 |
| SDS | 0 |
| Deoxycholic acid sodium salt | 136.9 |
| TritonX-100 | 137.8 |
| Tween 80 | 137.7 |

A reaction mixture containing 25 μ l of enzyme solution, 25 μ l of 0.2 M sodium phosphate buffer (pH 5.7), 10 μ l of each chemical (10 mM), 15 μ l of Milli-Q water and 25 μ l of 40 mM 1-kestose was incubated at 37°C for 10 min.

neokestose and nystose than sucrose and inulin. The result is similar to the kinetic parameters from native enzyme [11]. Janer *et al* reported that recombinant β -FFase from *B. lactis* (72% identity with amino acid sequence from *B. adolescentis* G1 β -FFase) showed high affinity to low molecular fructans (Raftilose, DP < 5) compared to low or high polymerized inulin (Raftiline LS or HP) or sucrose [15]. These observations were very similar to our result. Some properties (pH-optima, pH-stability, temperature-stability, effects of various metal salts or chemical reagents) are different between the recombinant and native enzymes, but the characterization are similar each other. We investigated substrate specificity for neokestose in this study, although it was not used in the previous research. Neokestose (β -D-fructofuranosyl (2 \rightarrow 6) - α -D-glucopyranosyl-(1 \rightarrow 2)- β -D-fructofuranoside) and its related fructo-oligosaccharides are contained in onion bulbs and asparagus roots. Neokestose and the related saccharides were reported to be synthesized by fructan:fructan 6^G-fructosyltransferase from the vegetables [23,24]. Recombinant β -FFase from *B. adolescentis* G1 preferentially hydrolyzed neokestose rather than 1-kestose. The enzyme hydrolyzes not only terminal β -2,1 fructoside bond of 1-kestose or nystose, but also β -2,6 fructoside bond (fru-2-6-glc) of neokestose. The result indicated that neokestose had potentiality to



Substrate
 (a) 1-Kestose
 (b) Neokestose
 (c) Nystose
 (d) Fructosylnystose
 (e) Sucrose
 (f) Inulin

Glc: glucose
 Fru: fructose
 Suc: sucrose
 1-K: 1-kestose
 N-k: neokestose
 Nys: nystose
 F-nys: fructosylnystose
 Inu: inulin

Figure 6 High performance anion-exchange chromatograms of the reaction products formed from fructo-oligosaccharide by recombinant β -FFase. Recombinant β -FFase was incubated with 10 mM 1-kestose (a), neokestose (b), nystose (c), fructosylnystose (d), sucrose (e) or inulin (f) for up to 4 h at 37°C. The reaction mixtures were analyzed by high performance anion-exchange chromatography (HPAEC). Glc, Glucose; Fru, Fructose; Suc, Sucrose; 1-K, 1-Kestose; N-k, Neokestose; Nys, Nystose; F-nys, Fructosylnystose; Inu, Inulin.

Table 3: Rate parameters of recombinant β -FFase.

| Substrate | K _m (a) | V _{max} (b) | k ₀ (c) | k ₀ /K _m (d) |
|------------------|--------------------|----------------------|--------------------|------------------------------------|
| 1-Kestose | 1.7 | 107 | 107.5 | 63.2 |
| Neokestose | 1.7 | 142 | 142.7 | 83.9 |
| Nystose | 3.9 | 152 | 152.8 | 39.2 |
| Fructosylnystose | 2.2 | 75 | 75.4 | 34.3 |
| Sucrose | 38 | 79 | 79.4 | 2.1 |
| Inulin | 25.9 | 77 | 77.4 | 3.0 |

a), mM; b), μ mol/mg of protein/min; c), sec⁻¹; d), mM⁻¹ · sec⁻¹

be a good efficient prebiotics, which can promote the human health due to growth of bifidobacteria in the gut.

Our result exhibited that the recombinant β -FFase from *B. adolescentis* G1 had unique properties to hydrolyze preferably low DP fructo-oligosaccharides as well as native one. We suppose that research about β -FFase efficiently to hydrolyze fructo-oligosaccharides is to be of much help for developing more effective prebiotics, probiotics and synbiotics.

Conclusion

In this article, cloning the *cscA* gene from *Bifidobacterium adolescentis* G1 and characterization of the recombinant protein of *cscA* gene expressed in *Escherichia coli* were described. Molecular mass of the purified recombinant enzyme was estimated to be about 66,000 by SDS-PAGE and 60,300 by MALDI TOF-MS. The optimum pH of the enzyme was 5.7 and the enzyme was stable at pH 5.0-8.6. The thermostability of the enzyme was up to 50°C. The K_m (mM), V_{max} (μ mol/mg of protein/min/), k_0 (sec⁻¹) and k_0/K_m (mM⁻¹ sec⁻¹) for 1-kestose, neokestose, nystose, fructosylnystose, sucrose and inulin were 1.7, 107, 107.5, 63.2, and 1.7, 142, 142.7, 83.9, and 3.9, 152, 152.8, 39.2, and 2.2, 75, 75.4, 34.3, and 38, 79, 79.4, 2.1, and 25.9, 77, 77.4, 3.0, respectively. The hydrolytic activity was strongly inhibited by AgNO₃, SDS, HgCl₂. The recombinant enzyme had similar specificity to the native enzyme, high affinity for 1-kestose, and low affinity for sucrose and inulin, although properties of the recombinant enzyme showed slight difference from those of the native one previously described.

Experimental

Bacterial strains, plasmids and culture condition

B. adolescentis G1 was cultured in GAM broth (Nissui Pharmaceutical Co., Ltd, Tokyo, Japan), and then, it was used for the extraction of genomic DNA. *Escherichia coli* DH5 α was used as host cell with the plasmid pGEM-T vector (Promega, Madison, WI, USA) for cloning and sequencing. *Escherichia coli* Rosetta2 (DE3) (Novagen, Madison, WI, USA) was used as host cell with the plas-

mid pET-32b (+) vector (Novagen) for protein expression. *E. coli* strains were grown in Luria Bertani (LB) medium supplemented with 100 μ g/ml carbenicillin.

Analysis of N-terminal amino acid Sequences

N-terminal sequences of β -FFase purified from *Bifidobacterium adolescentis* G1 was determined by ABI 477A protein sequencer/120A PTH analyzer system.

Gene cloning

After grown *B. adolescentis* G1 in GAM broth, the cells were collected by centrifugation (1,700 \times g, 10 min, 4°C) and the supernatant was removed, the precipitate was resuspended by adding 20 mM Tris-HCl buffer (pH 8.0), and centrifugated (1,700 \times g, 10 min, 4°C). Genomic DNA was extracted from the precipitate by DNeasy Tissue Kit (Qiagen, Courtaboueuuf, France).

Based on *cscA* gene encoding β -FFase from *Bifidobacterium adolescentis* ATCC 15703 [accession no. AP009256, protein ID BAF39931], a 1,659 bp of DNA including *cscA* gene from *B. adolescentis* G1 was amplified by PCR using *B. adolescentis* G1 genomic DNA as a template and using G1FFase1-for (5'-CCCAACAATTCATAACCCAG-3') and G1FFase2-rev (5'-TTCCCATATACCCCTTGCTA-3') as primers. PCR condition was: initial step of denaturation by 94°C for 2 min, followed by 30 cycles of 94°C for 15 sec, 57°C for 30 sec and 68°C for 90 sec, and then a final step at 68°C for 10 min using of KOD-plus- (Toyobo, Osaka, Japan). After adenine was attached to the PCR products by A addition kit (Qiagen), these products were ligated into pGEM-T vector using T4 DNA ligase (Promega) and *E. coli* DH5 α was transformed by the resulting vector. The transformants were grown and harvested, and the plasmids were isolated by Sigma GenElute Plasmid Mini-Prep Kit (Sigma-Aldrich, St. Louis, MO, USA), and its insert DNA was sequenced. The plasmid was named pGEM-G1cscA.

Expression of a recombinant protein in E. coli

For construction of expression vector, *cscA* gene was amplified by PCR using pGEM-G1cscA as a template. Primers used were ffaseNtEcoRI-for (5'-TCCGAATTC-

GATGACTGGCTTTACTCCGGA-3') and *ffase*CtXhoI-rev (5'-TTGCTCGAGTTCCAGTCCGATGGACTTCAT-3'). These primers had recognition sequence of *Eco*RI and *Xho*I, respectively. PCR condition was: initial step of denaturation by 94°C for 2 min, followed by 30 cycles of 94°C for 15 sec, 57°C for 30 sec and 68°C for 60 sec, and then a final step at 68°C for 7 min using of KOD-plus-. The PCR product was digested with *Eco*RI and *Xho*I, followed by ligation into pET-32b (+) vector cleaved with the same restriction enzyme using Quick T4 DNA ligase (New England Biolabs, Inc., Ipswich, MA). Finally, *E. coli* Rosetta2 (DE3) was transformed by the ligated pET vector. Resulting plasmid was named pET-G1cscA. The nucleotide sequence of the plasmid was analyzed by ABI 3730 × 1 sequencer (Applied Biosystems, Foster City, CA, USA), and confirmed it had no error. The transformants were selected on LB agar plates containing 100 µg/ml carbenicillin. A single colony of transformant was inoculated into 26 ml LB broth containing 100 µg/ml carbenicillin and grown in a shaking incubator at 37°C until the cell density at 600 nm reached 0.4. By adding 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG), and then, incubated for an additional 3 h, recombinant protein was induced. The cells were harvested by centrifugation (10,000 × g, 5 min, 4°C) and washed twice with 20 mM Tris-HCl. Finally, the cells were harvested by centrifugation (5,000 × g, 5 min, 4°C), and stored at -80°C until the preparation of the crude enzyme.

Purification of recombinant protein

All operations were done at 4°C. The cells were suspended in 8 ml wash buffer (50 mM NaH₂PO₄, 0.3 M NaCl, pH 7.5) containing protease inhibitor cocktail EDTA-free (Roche Molecular Biochemicals, Mannheim Germany), and the suspension was disrupted by ultrasonication. Supernatant after centrifugation (12,000 × g, 5 min) was used as the crude extract. The recombinant protein was purified from the crude extract by His-tag affinity resin (TALON CellThru Resin, Takara-Bio, Kyoto, Japan) according to the manufacturer's instructions. The active fraction was dialyzed for 1 day against the 10 mM sodium phosphate buffer (pH 7.5). To remove the His-tag peptides from recombinant protein in the purified fraction, recombinant enterokinase (rEK, Novagen) was used according to the manufacturer's instructions. The treated fraction was dialyzed for 1 day against the 10 mM sodium phosphate buffer (pH 7.5). This dialyzed was applied to a column (φ 1.7 cm × 20.2 cm) of DEAE-Sepharose CL-6B (GE Healthcare, Milwaukee, WI, USA) equilibrated with the same buffer. The adsorbed proteins were eluted with a linear gradient of sodium chloride from 0 to 1.0 M in the same buffer at the flow rate of 30 ml/h. The active fraction was dialyzed against 50 mM sodium phosphate buffer containing 0.1 M NaCl

(pH 7.5) for 1 day, and the dialyzed was concentrated by ultrafiltration with Vivaspin 20 (30,000 MWCO, Sartorius, Germany). The ultrafiltrate was filtered on a column (φ 2.4 cm × 67 cm) of Toyopearl HW-55S (Tosoh Co. Ltd, Tokyo, Japan) equilibrated with 50 mM sodium phosphate buffer containing 0.1 M NaCl (pH 7.5) at the flow rate of 30 ml/h. The active fraction was used as the purified enzyme solution.

Measurement of molecular mass

SDS-PAGE was conducted on a 12.5% (v/w) polyacrylamide gel by the method of Laemmli [25]. Proteins in the gel were stained with Coomassie Brilliant Blue R-250. LMW Marker (GE Healthcare) was used as molecular mass marker. The concentration of protein was spectrophotometrically measured at 280 nm. MALDI TOF-MS spectra were measured using a Shimadzu-Kratos mass spectrometer (KOMPACT Probe) in positive ion mode with 10 mg sinapic acid dissolved in 600 µl trifluoroacetic acid and 400 µl acetonitrile as a matrix. Ions were formed by a pulsed UV laser beam (nitrogen laser, 337 nm). Calibration was done using bovine serum albumin (BSA) as an external standard.


Enzyme assays

For the measurement of β-FFase activity, 50 µl of 20 mM 1-kestose in distilled water was mixed with 25 µl of 0.2 M sodium phosphate buffer (pH 5.7) and 25 µl of purified enzyme solution and incubated at 37°C for 10 min. The reaction was stopped by boiling for 5 min. One unit of β-FFase activity was defined as the amount of enzyme which produced 1 µmol of fructose per min under the above reaction conditions. For quantification of fructose, high performance anion exchange chromatography (HPAEC) was done on a DX300 chromatograph (Dionex Corp., Sunnyvale, USA) with a CarboPac PA-1 anion exchange column (Dionex Corp.) and a pulsed amperometric detector (PAD) as described previously [26].

For the determination of optimum pH, McIlvaine buffer with pH range 3.0-8.5 were used. The reaction was stopped by adding 900 µl of 150 mM NaOH.

To investigate the pH stability of enzyme, the mixture of 25 µl of Britton- Robinson buffer with pH range 3.0-10.0 and 25 µl of purified enzyme solution containing 0.1% BSA was kept at 4°C for 20 h, then the mixture was adjusted to pH 5.7, and incubated with 10 mM 1-kestose at 37°C for 10 min. The reaction was stopped by heating the samples at 100°C for 5 min.

For temperature stability profiles, 25 µl of 0.2 M sodium phosphate buffer (pH 5.7) and 25 µl of purified enzyme solution containing 0.1% BSA were mixed, and they were incubated at 4, 30, 40, 45, 50, 55, 60 and 65°C for 15 min, respectively, and then, each solution was cooled to 0°C. The mixtures were incubated with 50 µl of 20 mM 1-kes-



tose at 37°C for 10 min. The reaction was stopped by heating the samples at 100°C for 5 min.

The effects of metal salts and chemical reagents on the enzyme activity were investigated. The reaction mixture, 25 μ l of 40 mM 1-kestose, 25 μ l of 0.2 M sodium phosphate buffer (pH 5.7), 10 μ l of each metal salt or chemical reagents, 15 μ l of distilled water and 25 μ l purified enzyme was incubated at 37°C for 10 min. The reaction was stopped by boiling for 5 min.

To measure rate parameters of hydrolysis against different substrates, a reaction mixture containing 50 μ l of each substrate at various concentrations, 25 μ l of 0.2 M sodium phosphate buffer (pH 5.7) and 25 μ l of enzyme solution was incubated at 37°C for 10 min. Molecular activities (k_0) were calculated by using maximum velocities (V_{max}) and relative molecular mass of the enzyme.

Substrates

Saccharides as substrates were prepared as follows. Crystalline 1-kestose [1^F - β -D-fructofuranosylsucrose, 1-kesto-

triose] and nystose [$1^F(1-\beta$ -D-fructofuranosyl) $_2$ sucrose, 1, 1-kestotetraose] were prepared from sucrose using *Scopulariopsis brevicaulis* β -FFase [27]. Fructosylnystose [$1^F(1-\beta$ -D-fructofuranosyl) $_3$ sucrose, 1, 1, 1-kestopentaose] was prepared from Jerusalem artichoke tubers in our laboratory. Sucrose and inulin were purchased from Wako (Wako Pure Chemicals Industries, Osaka, Japan). Neokestose was prepared from asparagus roots as described previously [28]. The structures of the substrates were shown (Fig. 7).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

TO and KU collected data of gene sequences and recombinant protein, and contributed to drafting the manuscript. KM and MK collected data of native enzyme from bifidobacteria. NS and SO generated study design and drafted manuscript. All authors read and approved the final manuscript.

Author Details

Department of Food and Nutrition Sciences, Graduate School of Dairy Science Research, Rakuno Gakuen University, 582 Bunkyo-dai Midorimachi, Ebetsu, 069-8501, Japan

Received: 17 January 2010 Accepted: 12 April 2010

Published: 12 April 2010

References

- Gibson GR, Roberfroid MB: Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutr* 1995, **125**:1401-1412.
- Bacon JSD, Edelman J: The carbohydrates of the Jerusalem artichoke and other compositae. *Biochem J* 1951, **48**:114-126.
- Ende W Van den, Mintiens A, Speleers H, Onuoha AA, Van Laere A: The metabolism of fructans in roots of *Cichorium intybus* during growth, storage and forcing. *New Phytol* 1996, **132**:555-563.
- Ishiguro Y, Ueno K, Abe M, Onodera S, Fukushi E, Benkeblia N, Shiomi N: Isolation and structural determination of reducing fructooligosaccharides newly produced in stored edible burdock. *J Appl Glycosci* 2009, **56**:159-164.
- Abe M, Ueno K, Ishiguro Y, Omori T, Onodera S, Shiomi N: Purification, cloning and functional characterization of fructan: fructan 1-fructosyltransferase from edible burdock (*Arctium lappa* L.). *J Appl Glycosci* 2009, **56**:239-246.
- van Balken JAM, van Dooren ThJGM, Tweel WJJ van den, Kamphuis J, Meijer EM: Production of 1-kestose with intact mycelium of *Aspergillus phoenicis* containing sucrose-1 F -fructosyltransferase. *Appl Microbiol Biotechnol* 1991, **35**:216-221.
- Takeda H, Sato K, Kinoshita S, Sasaki H: Production of 1-kestose by *Scopulariopsis brevicaulis*. *J Ferment Bioeng* 1994, **77**:386-389.
- Hirayama M, Sumi N, Hidaka H: Purification and properties of a fructooligosaccharides-producing β -fructofuranosidase from *Aspergillus niger* ATCC 20611. *Agric Biol Chem* 1989, **53**:667-673.
- Hidaka H, Adachi T, Tokunaga T, Nakajima Y, Kono T: The road of fructooligosaccharide research and business development. In *Recent Advances in Fructooligosaccharides Research* Edited by: Shiomi N, Benkeblia N, Onodera S. Kerala: Research Signpost Publisher; 2007:375-395.
- Muramatsu K, Onodera S, Kikuchi M, Shiomi N: Purification and some properties of β -fructofuranosidase from *Bifidobacterium adolescentis* G1. *Biosci Biotechnol Biochem* 1993, **57**:1681-1685.
- Muramatsu K, Onodera S, Kikuchi M, Shiomi N: Substrate specificity and subsite affinities of β -fructofuranosidase from *Bifidobacterium adolescentis* G1. *Biosci Biotechnol Biochem* 1994, **58**:1642-1645.
- Muramatsu K, Onodera S, Kikuchi M, Shiomi N: The production of β -fructofuranosidase from *Bifidobacterium* spp. *Biosci Biotechnol Biochem* 1992, **56**:1451-1454.

13. Schell MA, Karmirantzou M, Snel B, Vilanova D, Berger B, Pessi G, Zwahlen MC, Desiere F, Bork P, Delley M, Pridmore RD, Arigoni F: **The genome sequence of *Bifidobacterium longum* reflects its adaptation to the human gastrointestinal tract.** *Proc Natl Acad Sci USA* 2002, **99**:14422-14427.
14. Warchol M, Perrin S, Grill JP, Schneider F: **Characterization of a purified β -fructofuranosidase from *Bifidobacterium infantis* ATCC 15697.** *Lett Appl Microbiol* 2002, **35**:462-467.
15. Janer C, Rohr LM, Pelaez C, Laloi M, Cleusix V, Requena T, Meile L: **Hydrolysis of oligofructoses by the recombinant β -fructofuranosidase from *Bifidobacterium lactis*.** *Syst Appl Microbiol* 2004, **27**:279-285.
16. Kullin B, Abratt VR, Reid SJ: **A functional analysis of the *Bifidobacterium longum* *cscA* and *scrP* genes in sucrose utilization.** *Appl Microbiol Biotechnol* 2006, **72**:975-81.
17. Reddy VA, Maley F: **Identification of an active-site residue in yeast invertase by affinity labeling and site-directed mutagenesis.** *J Biol Chem* 1990, **265**:10817-10820.
18. Reddy A, Maley F: **Studies on identifying the catalytic role of Glu-204 in the active site of yeast invertase.** *J Biol Chem* 1996, **271**:13953-13958.
19. Batista FR, Hernández L, Fernández JR, Arrieta J, Menéndez C, Gómez R, Tambara Y, Pons T: **Substitution of Asp-309 by Asn in the Arg-Asp-Pro (RDP) motif of *Acetobacter diazotrophicus* levansucrase affects sucrose hydrolysis, but not enzyme specificity.** *Biochem J* 1999, **377**:503-506.
20. Meng G, Fütterer K: **Structural framework of fructosyl transfer in *Bacillus subtilis* levansucrase.** *Nat Struct Biol* 2003, **10**:935-941.
21. **Carbohydrate-Active enzymes Database** [<http://www.cazy.org/>]
22. Shirai A, Matsuyama A, Yashiroda Y, Hashimoto A, Kawamura Y, Arai R, Komatsu Y, Horinouchi S, Yoshida M: **Global analysis of gel mobility of proteins and its use in target identification.** *J Biol Chem* 2008, **283**:10745-10752.
23. Ueno K, Onodera S, Kawakami A, Yoshida M, Shiomi N: **Molecular characterization and expression of a cDNA encoding fructan:fructan 6⁶-fructosyltransferase from asparagus (*Asparagus officinalis*).** *New phytologist* 2005, **165**:813-824.
24. Fujishima M, Sakai H, Ueno K, Takahashi N, Onodera S, Benkeblia N, Shiomi N: **Purification and characterization of a fructosyltransferase from onion bulbs and its key role in the synthesis of fructo-oligosaccharides *in vivo*.** *New phytologist* 2005, **165**:513-524.
25. Laemmli UK: **Cleavage of structural proteins during the assembly of the head of bacteriophage T4.** *Nature* 1970, **227**:680-685.
26. Shiomi N, Onodera S, Chatterton NJ, Harrison PA: **Separation of fructooligosaccharide isomers by anion-exchange chromatography.** *Agric Biol Chem* 1991, **55**:427-428.
27. Takeda H, Sato K, Kinoshita S, Sasaki H: **Production of 1-kestose by *Scopulariopsis brevicaulis*.** *J Ferment Bioeng* 1994, **77**:386-389.
28. Shiomi N, Yamada J, Izawa M: **Isolation and identification of fructo-oligosaccharides in roots of asparagus (*Asparagus officinalis* L.).** *Agr Biol Chem* 1976, **40**:567-575.

doi: 10.1186/1752-153X-4-9

Cite this article as: Omori *et al.*, Characterization of recombinant β -fructofuranosidase from *Bifidobacterium adolescentis* G1 *Chemistry Central Journal* 2010, **4**:9

Publish with **ChemistryCentral** and every scientist can read your work free of charge

“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>


ChemistryCentral