

SCIENTIFIC REPORTS

OPEN

Erratum: Predictive analytics of environmental adaptability in multi-omic network models

Claudio Angione & Pietro Lió

Scientific Reports 5:15147; doi: 10.1038/srep15147; published online 20 October 2015; updated on 20 May 2016

This Article contains typographical errors.

In the Results section under subheading ‘METRADE: a novel method to integrate and optimize gene expression and codon usage in FBA’,

“Through the function $h(\cdot)$, the gene expression arrays have a continuous effect on the FBA model, rather than only an on/off effect on reactions as in the Boolean approaches.”

should read:

“Through the function $h(\cdot)$, the gene expression arrays have a continuous effect on the FBA model, rather than only an on/off effect on reactions as in the Boolean approaches.”

In the Results section under subheading ‘Mapping genotype-phenotype associations to multidimensional objective spaces’,

“Then, we solve the bilevel problem (4) replacing the function h with the function $k(y_i) = [1 + |\log(y_i)|]^{\text{sgn}(y_i-1)}$, where σ_i^2 is the variance of the gene set responsible for the i th reaction, and γ is a weight for the variance.”

should read:

“Then, we solve the bilevel problem (4) replacing the function h with the function $k(y_i) = [1 + \frac{\gamma}{\sigma_i^2} |\log(y_i)|]^{\text{sgn}(y_i-1)}$, where σ_i^2 is the variance of the gene set responsible for the i th reaction, and γ is a weight for the variance.”



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>