RESEARCH ARTICLE

Open Access

The influence of amaranth (Amaranthus hypochondriacus) dietary nitrates on the aerobic capacity of physically active young persons



Tomas Liubertas^{1*}, Ramutis Kairaitis¹, Loreta Stasiule², Sandrija Capkauskiene², Arvydas Stasiulis², Pranas Viskelis³, Jonas Viškelis³ and Dalia Urbonaviciene³

Abstract

Background: Recent evidence indicates that elevating plasma nitrites through dietary nitrates (NO_3^-) supplementation is associated with enhanced muscle efficiency, fatigue resistance and performance. Beetroot (in various forms) is the dominant source of dietary NO_3^- primarily due to its vast availability and the simple form of preparation suitable for final consumption. After a few years of research and experimentation, our scientific team identified alternative source rich with dietary NO_3^- as possible nitric oxide precursor, amaranth (*Amaranthus hypochondriacus*) with a standardized concentration 9–11% of NO_3^- . This study aimed to evaluate the effect of single-dose ($\pm 400 \text{ mg}$ of dietary NO_3^-) and long-term (6 days) supplementation of amaranth concentrate derived dietary NO_3^- on aerobic capacity in physically active young people.

Methods: We conducted a randomized, double-blind, placebo-controlled human study. Thirteen healthy and physically active young male participants were randomized into experimental and placebo groups. The aerobic capacity was tested during increasing cycling exercise (ICE) with pulmonary gas exchange recording and analysis.

Results: The peak power of the ICE, the maximum oxygen consumption and the first ventilatory threshold were significantly increased after long-term consumption of dietary amaranth (from 4.44 ± 0.50 to 4.55 ± 0.43 W/kg; from 37.7 ± 2.7 to 41.2 ± 5.4 mL/kg/min and from 178.6 ± 30.3 to 188.6 ± 35.2 W, p < 0.05; respectively) in experimental group.

Conclusions: Long-term (6 days) use of dietary NO_3^- from amaranth may improve the aerobic capacity during ICE in young physically active male persons. It can be recommended as the nutritional supplement during last week of preparation for competition in endurance events.

Keywords: Aerobic capacity, Dietary nitrates, Amaranth, Cycling, Young persons

Full list of author information is available at the end of the article



^{*} Correspondence: tomas.liubertas@stud.lsu.lt

¹Department of Coaching Science, Lithuanian Sports University, 44221 Kaunas. Lithuania

Introduction

Green leafy vegetables and roots are the main source of dietary nitrates (NO₃⁻) [1–3]. NO₃⁻ is a naturally occurring compound as well as an approved food additive [1, 2]. A number of studies have already confirmed the benefits of dietary nitrates to human health [4]: their consumption reduces blood pressure, suppresses platelet aggregation, protects against ischemic diseases, and improves endothelial function [1]. Nitric oxide and nitrites, both NO₃⁻ products, affect vasodilatation by increasing blood flow [5], thus increasing the oxygen uptake and oxidative processes in the working muscles [6]. Additionally, nitrates show to increase the bioavailability of blood plasma, which is important for the exogenous pathway of nitrates-nitrite-NO and acts as a regulator of hypoxic signals and NO-induced vasodilatation [7].

The effects of nitrate/nitrite/NO on the muscle circulatory system and mitochondrial and contractile efficacy [8, 9] may increase muscle blood flow circulation and improve the metabolic response to physical activity [10]. The evidence supports that even the concentration of plasma nitrites is an independent factor of physical performance [5, 11]. Nevertheless, studies on the effects of nitrates on work capacity indicators are highly controversial so far.

Studies have shown nitrates to have a positive effect on work efficiency and oxygen expenditure [10, 12–17], but other studies have not found visible and conclusive changes in given performance [14, 18–22].

A large number of researchers found that 300-500 mg of beetroot nitrates have a single and long-lasting positive effect on the aerobic performance of physically active individuals [10, 12, 16–19, 23]. Recently became popular and actively researched beetroot (in various forms) is the dominant source of dietary NO₃⁻, primarily due to its vast availability and the simple form of preparation suitable for final consumption. Remarkably, limited studies have evaluated NO₃ rich leafy vegetables and, more specifically, amaranth on exercise performance. Importantly, amaranth is not only rich in NO₃, potassium (> 10% by weight) and antioxidant polyphenols (e.g. amaranthine), but also devoid of sugar and oxalates. It has recently been reported that red spinach extract as a nutritional supplement can elicit an ergogenic response by delaying the ventilatory threshold during graded treadmill exercise testing [24]. After a few years of research and experimentation, our scientific team identified alternative source rich in dietary NO₃⁻ as possible and alternative nitric oxide precursor, amaranth (Amaranthus hypochondriacus) with a standardized concentration 9–11% of NO₃⁻. Since NO₃⁻ supplementation increases plasma NO2-, this intervention may therefore have the potential to improve muscle blood flow [5], thus increasing the oxygen uptake and oxidative processes in the working muscles [6] and exercise tolerance. Thus, based on research data available we formulated the hypothesis that $400\,\mathrm{mg}$ of dietary $\mathrm{NO_3}^-$ from amaranth (dietary amaranth) will increase the aerobic capacity of physically active young people.

In this study we aimed to evaluate the effect of single and long-term doses of dietary amaranth on the aerobic capacity of physically active young persons.

Materials and methods

Participants

The study recruited 13 volunteering graduate students (all males) from Lithuanian Sports University. Every participant was informed about the research objectives and methods and signed an Informed Consent form for participation. The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Kaunas Regional Ethics Committee, Nr. BE-2-11, 21 March 2017. The anthropometric data and age of the participants are presented in Table 1.

Measurements

Anthropometry

Electronic weighing scales (Body Composition Analyzer TBF-300, Tanita, Japan). were used to measure the weight and relative fat mass of the participants. Height was measured using a stadiometer (Leicester height measure, UK).

Increasing cycling exercise (ICE)

For the assessment of aerobic capacity, an ICE was performed on a cycle ergometer (Ergoline-800, Denmark). The seat and handlebar positions on the cycle ergometer were adjusted for each subject prior to the initial exercise test and maintained in that position for the subsequent exercise tests. Prior to the ICE, a 5-min warm-up was performed. The ICE consisted of 3 min of cycling at 40 W, then the ramp protocol was applied, and the workload was continuously increased by 30 W per min. The cadence was 70 rpm. The participants were encouraged to exercise until voluntary fatigue, and the test was stopped when the participant was not able to maintain a cadence above 65 rpm.

Table 1 Characteristics of study participants

Variable	Placebo group ¹	Experimental group ²	<i>p</i> -Value
Age (years)	21.9 ± 1.9	21.3 ± 0.8	0.940
Height (cm)	182.1 ± 5.9	180.5 ± 8.3	0.829
Body weight (kg)	88.1 ± 12.5	84.3 ± 23.7	0.668
Body mass index (BMI)	26.5 ± 2.8	25.6 ± 5.8	0.568
Relative fat mass (%)	13.1 ± 3.8	18.9 ± 5.6	0.063

Values are reported as the mean \pm standard deviation (SD). ¹ n = 6; ² n = 7

Pulmonary gas exchange recording and analysis

The subjects breathed through a face mask, and pulmonary gas exchange parameters were measured breath-by-breath using a wireless, portable spirometric system "Oxycon Mobile" (Viasys Healthcare; California, USA). Prior to each exercise session, the spirometric system was calibrated according to the recommendations of manufacturers. The maximum value of oxygen uptake (VO₂) over the 20 s of cycling was referred to as VO2 max, and the first and second ventilatory thresholds (VT1 and VT2) were determined from the data of the incremental cycling exercise. The determination was based on the analysis of the relationship between ventilatory equivalents of oxygen or carbon dioxide and cycling power. The VT1 was identified as the first point at which the ventilatory equivalent for O2 increased without a concurrent increase in the ventilatory equivalent for CO₂. The VT2 was identified as the point of constant increase of ventilatory equivalent of CO₂. A least squares method was used to fit two lines representing ventilatory equivalents versus load plots. The intersection point of the two regression lines was assigned to VT1 and VT2. Heart rate (HR) was recorded continuously with a wireless Polar monitoring system (Polar, Finland).

Biochemical analysis of blood

Blood samples for the measurement of blood lactate concentration [La] (Lactate Pro2, Japan) were taken from fingertips at the end of the 5th min of recovery after the ICE.

Protocol

A randomized double-blinded design was used in this study. During the first visit, the participants had their anthropometric measurements taken and performed the ICE (1 T) (Fig. 1).

After 2 days of rest, participants repeated the ICE (2 T) 1 h after consumption of supplement with (experimental group) or without (placebo group) *Amaranthus hypochondriacus*. Then, participants in both groups consumed supplement for 4 days during breakfast and, on the next day, performed a third ICE (3 T) after consuming supplements 1 h before the test.

The experimental group consumed a hand-made oat bar supplement (OB) 60 g total weight - made of oats, honey, vanilla, containing 4 g standardized $Amaranthus\ hypochondriacus$ concentrate (9–11% equivalent to \pm 400 mg of active ingredient (NO $_3$ -). The placebo group consumed visually and flavory identical oat bar (OB) - 60 g, containing oats, honey and vanilla (excluding active ingredient - ie. $Amaranthus\ hypochondriacus$ concentrate). Participants were asked not to change their nutritional habits during the period of study.

Statistical analysis

The statistical analysis was carried out with SPSS (Statistical Package for Social Sciences, version 19.0) and Microsoft Office Excel 2007. The normal distribution of variables was checked using the Kolmogorov-Smirnov test. Non-parametric data analysis methods were used to assess the effect of dietary amaranth on aerobic capacity. The significance of the difference between the independent samples was evaluated using the Mann-Whitney test.




Fig. 1 Study protocol. Note: ICE – increasing cycling exercise test; OB - handmade oat bar; 1–9 –days of study. (1 T) - test Nr. 1 executed at day 1 of the study; (2 T) - test Nr. 2 executed at day 4 of the study; (3 T) - test Nr. 3 executed at day 9 of the study




Fig. 2 Peak power of increasing cycling exercise (ICE) following single-dose (2 T) and long-term (3 T) doses of supplements in the experimental and placebo groups. * p < 0.05 - statistically significant difference compared to the 1 T




Fig. 3 Maximal oxygen uptake (VO2max) following single-dose (2 T) and long-term (3 T) doses of supplements in the experimental and placebo groups. * p < 0.05 - statistically significant difference compared to the 1 T




Fig. 4 First ventilatory threshold (VT1) and second ventilatory threshold (VT2) following single-dose (2 T) and long-term (3 T) doses of supplements in the experimental and placebo groups. *p < 0.05- statistically significant difference between the 2 T and 3 T

The difference between dependent samples was assessed by the Wilcoxon test. Statistical significance was accepted when p < 0.05. All data are reported as the mean \pm standard deviation (SD).

Results and discussion

Results

No significant changes of the parameters of aerobic capacity were observed in Placebo group (Figs. 2, 3, 4). Peak power of ICE has increased significantly in the experimental group, from 4.42 ± 0.50 W/kg during the first testing to 4.55 ± 0.43 W/kg during the third testing (P=0.043; Fig. 2). The single dose of supplements did not have any significant effect on the VO₂max in the experimental group. After long-term use of dietary amaranth, absolute and relative values of VO₂max demonstrated a significant increase in the experimental group (from 3.282 ± 0.51 l/min and 37.7 ± 2.7 mL/kg/min during the first test to 3.599 ± 0.51 l/min (p=0.028) and 41.2 ± 5.4 mL/kg/min (p=0.043) during the third test, respectively (Fig. 3).

VT1 did not change after a single dose of dietary amaranth, but a significant difference was observed in the experimental group after long-term supplementation (third testing). After a single dose dietary amaranth VT1 was 2.02 ± 0.13 W/kg and after long-term use of dietary amaranth the value of VT1 increased significantly to

 2.14 ± 0.16 W/kg (p = 0.028; Fig. 4a). The VT2 did not change significantly in any of the groups (p > 0.05, Fig. 4b).

Other physiological variables measured during ICE did not change significantly in any group of participants (Table 2).

Table 2 Maximal physiological responses during increasing cycling exercise following single-dose (2 T) and long-term (3 T) doses of supplements in the experimental and placebo groups (HR - heart rate; RER - respiratory exchange ratio; [La] - blood lactate concentration)

Indexes	Experimental group		
	1 T	2 T	3 T
HR, beats/min	186.1 ± 8.2	183.8 ± 9.1	184.2 ± 7.4
RER	1.18 ± 0.07	1.16 ± 0.04	1.24 ± 0.06
[La] at 5th min after ICE, mmol/l	14.3 ± 2.9	15.2 ± 4.8	12.4 ± 1.6
Indexes	Placebo group		
	1 T	2 T	3 T
HR, beats/min	189.2 ± 5.7	188.2 ± 8.1	187.0 ± 7.6
RER	1.20 ± 0.07	1.16 ± 0.04	1.22 ± 0.07
[La] at 5th min after ICE, mmol/l	13.0 ± 2.1	13.6 ± 4.6	13.9 ± 4.2

Key: Values are expressed as mean \pm SD

Table 3 Effects of single and long-term use of nitrates on the indicators of work capacity of healthy and physically active persons

References	Participants	Supplemen-tation	Concentration of nitrates mmol/day	Duration of consumption	Protocol	Findings
[12]	Healthy, physically active people, $n = 8$; placebo (P) and experimental groups (E)	500 ml beetroot juice /day	~ 340 mg or 11 mmol	6 days	Time trial at 70% between the ventilation threshold and VO ₂ max	The increase in pulmonary O_2 uptake following the onset of moderate exercise was reduced by 19% in the beetroot juice condition. During severe exercise, the O_2 uptake slow component was reduced, and the timeto-exhaustion was extended. Trial duration: E: 675 ± 203 s P: 583 ± 145 s ($p < 0.05$)
[19]	Healthy people, $n = 9$;	NaNO ₃	0.033mmol $\text{NaNO}_3 \text{kg/ body}$ WT three times daily	Randomized and double-blinded; the washout period between the two trials was at least 7 days	Maximal combined arm and leg exercise tests.	Dietary nitrate reduced VO ₂ max from $3.72+/-0.33$ to $3.62+/-0.31$ L/min, $p < 0.05$. Despite the reduction in VO ₂ max the time-to-exhaustion trended towards an increase after nitrate supplementation (524+/- 31 vs $563+/-30$ s, $p = 0.13$).
[10]	Healthy, physically active people; n=7	500 ml beetroot juice /day	300 mg or ~ 5,1 mmol	6 days	Leg extension until fatigue at 30% of MVC	20% reduced O_2 cost of exercise following dietary NO_3 - supplementation appears to be due to a reduced ATP cost of muscle force production. Trial duration: E: $734 \pm 109 \text{ s}$ P: $586 \pm 80 \text{ s}$ ($p < 0,05$)
[18]	Healthy, physically active people; n = 9	500 ml beetroot juice /day	300 mg or ~ 5,1 mmol	A. 2,5 h before B. 5 days C. 15 days	Increasing exercise test	A. E: 325 ± 71 Wmax P: 322 ± 68 Wmax B. E:328 ± 68 Wmax P: 323 ± 67 Wmax C. E:331 ± 68 Wmax P: 323 ± 68 Wmax After 15 days maximal power was increased by 4%
[16]	Healthy, physically active people; n = 9	500 ml of beetroot juice (BR)	380 mg or ~ 6,1 mmol	6 days	Subjects completed treadmill exercise tests and knee-extension exercise tests for estimation of Q _{max} .	The O_2 cost of walking, moderate-intensity running, and severe-intensity running was reduced by BR; time-to-exhaustion during severe-intensity running was increased by 15%. Trial time: E: 8,7 \pm 1,8 min P: 7,6 \pm 1,5 min (p < 0,05)
[17]	Healthy, physically active men; n = 15	Beetroot juice	4,4 mg or 0.07 mmol nitrate/kg body WT/day	6 days	Increased exercise test simulating a 5000-m altitude	Short-term dietary nitrate supplementation improves arterial and muscle oxygenation status but not cerebral oxygenation status during exercise in severe hypoxia. Trial duration: E: 597 ± 22 s P: 568 ± 23 s (p < 0,05)
[11]	Healthy, recreationally active men; n = 10	70, 140, or 280 ml of beetroot juice or placebo (PL) 70, 140, or 280 ml	~ 4.2, ~ 8.4, or ~ 16.8 mmol nitrates or placebo containing ~ 0.04, ~ 0.08, or ~ 0.12 mmol nitrates	4–5 weeks	Two, 5-min bouts of moderate-intensity exercise and one bout of severe-intensity exercise that was continued until task failure as a measure of exercise tolerance, 2.5 h post-ingestion	8.4 and 16.8 mmol of nitrates significantly improved the time-to-task failure by 14 and 12%, re- spectively, during severe- intensity exercise. End- exercise VO ₂ during

Table 3 Effects of single and long-term use of nitrates on the indicators of work capacity of healthy and physically active persons (Continued)

References	Participants	Supplemen-tation	Concentration of nitrates mmol/day	Duration of consumption	Protocol	Findings
					of beetroot juice or placebo.	moderate-intensity exercise was reduced significantly following the ingestion of 280 ml of BR ($p < 0.05$).
[23]	Healthy, physically active men; n = 16	500 ml beetroot juice	5 mmol	1.5 h before physical load	A continuous cycle exercise test involving 20-min stages at 50 and 70% VO ₂ max and a final stage at 90% VO ₂ max until volitional exhaustion.	Dietary nitrate reduced VO_2 max by 15,63% and increased the time-to-exhaustion by 16%. Trial duration: E: 185 ± 122 s P: 160 ± 109 s $(p < 0,05)$
[25]	Healthy, recreationally active participants n = 15 (males = 8, females = 7)	1000 mg red spinach extract powder (RSE) or placebo (PBO) (maltodextrin) in gelatin capsules	~ 90 mg	One single dose	At one occasion 65–75-min post-RSE/PBO ingestion, venipuncture was performed (PRE), after which graded exercise test - the Bruce protocol - was performed.	Significantly increased plasma NO_3 A large effect on the ventilatory threshold compared to the placebo (\pm 0.12 \pm 0.14 L/min) with the ventilatory threshold occurred at a significantly higher relative VO_2 (\pm 3.6 \pm 5.2%, p < 0.05).

Discussion

In the present study we analyzed the effect of single dose (equivalent to ± 400 mg) and long term consumption of dietary nitrates obtained from amaranth (*Amaranthus hypochondriacus*) on aerobic capacity of healthy physically active young people. Our principal findings are that the VO₂max, VT1 and peak power measured during gradually increasing cycling exercise increased significantly only after 6 days of supplementation, but demonstrated no significant effect after a single dose.

Our data coincides with other studies of a similar nature carried out by other authors (Table 3). The researchers cited used beetroot juice with varying amounts of NO₃. The most popular use of the fixed duration of the dietary NO₃ is 4-6 days, and the amount varies from 5 mmol up to 9 mmol. Positive and significant effect for VO₂max was found by Bailey [10, 12], Lansley [16], Vonhatalo [18], and Wylie [11]. Their studies involved healthy, physically active people and tested physical performance. Vanhatalo [18] explored the long-term effects of nitrates with 5 and 15 days of 500 ml of 5.2 mmol nitrate from beetroot juice and found that the 15-day period significantly improved the maximum aerobic power (Table 3). Most previous studies have used beetroot juice as a source of dietary NO3-, but only a few have investigated the effect of other sources. In one study, the immediate ergogenic effect (delaying the ventilatory threshold) of red spinach extract was demonstrated [24]. The authors reported that after acute ingestion of 1000 mg of red spinach extract, VO2 at the ventilatory threshold was significantly higher, although no significant changes were seen in the time-toexhaustion or maximal aerobic power. By contrast, in our study, significant changes were observed after 6 days of dietary amaranth consumption. Our dose of NO3–was lower (400 mg) than that in the report by More et al. [25], and this could be why a single dose had no effect.

The mechanisms by which NO_3 supplementation might influence oxygen uptake and utilization could include a reduction of NO_3 to NO, directly influencing mitochondrial efficiency, vascular tone and/or tissue oxygenation. However, the evidence for these mechanisms depends on the measurement of NO_2 in the plasma [9].

Amaranth is a leafy vegetable, and its leaves and grains contain large amounts of NO₃ as well as other nutrients [26]. The NO₃ content of amaranth may range from 965 to 4259 mg/kg [1] or from 1800 to 9200 mg/kg [27]. There is scientific evidence suggesting that nitrate-rich spinach can augment nitric oxide levels, enhance endothelial function, and lower blood pressure acutely [13]. Moore and co-authors [25] reported that acute ingestion of 1000 mg of an amaranth extract substantially increased only the plasma NO₃ level and not the NO₂ level. Subramanian and Gupta [28] reported that an acute 2000 mg dose of amaranth extract, delivering ~ 180 mg (~ 2.9 mmol) of NO₃, increased plasma NO₃ and NO₂. This increase is similar to, or exceeds that observed with acute ingestion of relatively higher NO₃ doses from beetroot juice [11, 18].

There were some limitations to our study. We did not measure the plasma contents of NO₃ or NO₂; thus, based on the results of other studies, we can only assume that it increased after 6 days of amaranth consumption. Further studies are needed to measure acute

and long-term changes in plasma NO3 and NO2 after the consumption of given amaranth supplement. Secondly, the number of study participants was rather small, so the statistical power was moderate (49.9% for the absolute VO2 max in the experimental group). We did not utilize a crossover type of experiment design in this particular study which is a particularly useful option in evaluating the safe and effective use of tested substances when participants switch from one substance to another. It has both advantages and disadvantages as compared to parallel study. Usually the order effect and carry-over effects are discussed among limitations. In our case the study involved repeated graded exercise tests so often testing might have some training effect. The washout period of ingesting dietary nitrates is unknown, so the experiment could be rather long with increased influences of different covariates. In this case a randomized double-blinded was used as a way of carrying out an experiment in an attempt to minimize subjective biases on the part of the experimenter and on the part of the participant.

Conclusion

Long-term (6 days) supplementation with dietary NO_3 from amaranth may improve aerobic capacity during ICE in young physically active male persons. It can be recommended as the nutritional supplement during last week of preparation for competition in endurance events.

Abbreviations

ICE: Increased cycling exercise; OB: Oat bar; VT1: First ventilatory threshold; VT2: Second ventilatory threshold; 1 T: Test Nr. 1 executed at day 1 of the study; 2 T: Test Nr. 2 executed at day 4 of the study; 3 T: Test Nr. 3 executed at day 9 of the study; HR: Heart rate; RER: Respiratory exchange ratio; La: Blood lactate concentration; SD: Standard deviation; NO $_3$ ⁻: Nitrate; NO: Nitric oxide; W: Watts; VO $_2$ max: The maximum rate of oxygen consumption measured during incremental exercise

Acknowledgements

We would like to show our gratitude to all the participants of this study and thank them for sharing their experience during the course of this research. Especially we would like to express our gratitude and we thank 3 "anonymous" reviewers for their so-called insights and for their comments on an earlier version of this manuscript.

Authors' contributions

The study was designed by T.L., R.K. and A.S.; the active components were extracted, fractionated and prepared for supplementation by P.V. and D.U.; the subjects were recruited and hosted the informative session by R.K.; L.S., A.S., S.C., checked that subjects followed the diet guidelines and the timing of supplement ingestion; data were collected and analyzed by A.S., L.S., S.C., T.L., P.V.; L.S., S.C., T.L. and J.V. conducted the statistical analysis; and data interpretation and manuscript preparation were undertaken by T.L., R.K., L.S., S.C., A.S., P.V. J.V. and D.U. The author(s) read and approved the final manuscript.

Funding

Authors received no specific funding for this work.

Availability of data and materials

The data that support the findings of this study are available on request from the corresponding Tomas Liubertas, T.L. The data are not publicly

available due to General Data Protection Regulation (GRDP) of European Union (restrictions containing information that could compromise the privacy of research participants);

Ethics approval and consent to participate

The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Kaunas Regional Ethics Committee, Nr. 8F-2-11, 21 March 2017.

Consent for publication

All authors approved the final version of the paper.

Competing interests

The authors declare no conflict of interest.

Author details

¹Department of Coaching Science, Lithuanian Sports University, 44221 Kaunas, Lithuania. ²Department of Applied Biology and Rehabilitation, Lithuanian Sports University, 44221 Kaunas, Lithuania. ³Institute of Horticulture, Lithuanian Research Centre for Agriculture and Forestry, 54333 Babtai, Lithuania.

Received: 27 January 2020 Accepted: 30 June 2020 Published online: 13 July 2020

References

- Lidder S, Webb AJ. Vascular effects of dietary nitrate (as found in green leafy vegetables and beetroot) via the nitrate-nitrite-nitric oxide pathway. Br J Clin Pharmacol. 2013;75(3):677–96. https://doi.org/10.1111/j.1365-2125.2012. 04420.x PMID: 22882425; PMCID: PMC3575935.
- Santamaria P. Nitrate in vegetables: toxicity, content, intake and EC regulation. J Sci Food Agric. 2006;86:10–7.
- Hord NG, Tang Y, Bryan NS. Food sources of nitrates and nitrites: the physiologic context for potential health benefits. Am J Clin Nutr. 2009;90(1): 1–10. https://doi.org/10.3945/ajcn.2008.27131 Epub 2009 May 13.
- Stanaway L, Rutherfurd-Markwick K, Page R, Ali A. Performance and health benefits of dietary nitrate supplementation in older adults: a systematic review. Nutrients. 2017;9(11):1171.
- Totzeck M, Hendgen-Cotta UB, Luedike P, Berenbrink M, Klare JP, Steinhoff H-J, et al. Nitrite regulates hypoxic vasodilation via myoglobin-dependent nitric oxide generation. Circulation. 2012;126(3):325-+.
- Dreißigacker U, Wendt M, Wittke T, Tsikas D, Maassen N. Positive correlation between plasma nitrite and performance during high-intensive exercise but not oxidative stress in healthy men. Nitric Oxide. 2010;23(2):128–35.
- Lundberg JO, Carlstrom M, Larsen FJ, Weitzberg E. Roles of dietary inorganic nitrate in cardiovascular health and disease. Cardiovasc Res. 2011;89(3):525–32.
- Larsen FJ, Schiffer TA, Borniquel S, Kent Sahlin K, Ekblom B, Lundberg JO, Weitzberg E. Dietary inorganic nitrate improves mitochondrial efficiency in humans. Cell Metab. 2011;13(2):149–59.
- Larsen FJ, Schiffer TA, Weitzberg E, Lundberg JO. Regulation of mitochondrial function and energetics by reactive nitrogen oxides. Free Radic Biol Med. 2012;53(10):1919–28.
- Bailey SJ, Fulford J, Vanhatalo A, Winyard PG, Blackwell JR, DiMenna FJ, Wilkerson DP, Benjami N, Jones AM. Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. J Appl Physiol. 2010;109:135–48.
- Wylie LJ, Kelly J, Bailey SJ, Blackwell JR, Skiba PF, Winyard PG, Jeukendrup AE, Vanhatalo A, Jones AM. Beetroot juice and exercise: pharmacodynamic and dose-response relationships. J Appl Physiol. 2013;115(3):325–36.
- Bailey SJ, Winyard P, Vanhatalo A, Blackwell JR, DiMenna FJ, Wilkerson DP, Tarr J, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the O₂ cost of low-intensity exercise and enhances tolerance to high-intensity exercise in humans. J Appl Physiol. 2009;107(4):1144–55.
- Bond H, Morton L, Braakhuis AJ. Dietary nitrate supplementation improves rowing performance in well-trained rowers. Int J Sport Nutr Exerc Metab. 2012;22:251–6.
- Cermak NM, Gibala MJ, van Loon LJ. Nitrate supplementation's improvement of 10-km time-trial performance in trained cyclists. Int J Sport Nutr Exerc Metab. 2012;22(1):64–71.

- Lansley KE, Winyard PG, Bailey SJ, Vanhatalo A, Wilkerson DP, Blackwell JR, Gilchrist M, Benjamin N. Acute dietary nitrate supplementation improves cycling time trial performance. Med Sci Sports Exerc. 2011a;43(6):1125–31.
- Lansley KE, Winyard PG, Fulford J, Vanhatalo A, Bailey SJ, Blackwel JR, DiMenna FJ, Gilchrist M, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the O₂ cost of walking and running: a placebocontrolled study. J Appl Physiol. 2011b;110(3):591–600.
- Masschelein E, Van Thienen R, Wang X, Van Schepdael A, Thomis M, Hespel P. Dietary nitrate improves muscle but not cerebral oxygenation status during exercise in hypoxia. J Appl Physiol. 2012;113(5):736–45.
- Vanhatalo A, Bailey SJ, Blackwell J, DiMenna FJ, Pavey TG, Wilkerson DP, Benjamin N, Winyard PG, Jones AM. Acute and chronic effects of dietary nitrate supplementation on blood pressure and the physiological responses to moderate-intensity and incremental exercise. [randomized controlled trial]. Am J Phys Regul Integr Comp Phys. 2010;299(4):R1121–31.
- Larsen FJ, Weitzberg E, Lundberg JO, Ekblom B. Dietary nitrate reduces maximal oxygen consumption while maintaining work performance in maximal exercise. Free Radic Biol Med. 2010;48(2):342–7.
- Murphy M, Eliot K, Heuertz RM, Weiss E. Whole beetroot consumption acutely improves running performance. J Acad Nutr Diet. 2012;112(4):548–52.
- Peacock O, Tjonna AE, James P, Wisløff U, Welde B, Böhlke N, Smith A, Stokes K, Cook C, Sandbakk O. Dietary nitrate does not enhance running performance in elite cross-country skiers. Med Sci Sports Exerc. 2012;44(11): 2213–9.
- Wilkerson DP, Hayward GM, Bailey SJ, Vanhatalo A, Blackwell JR, Jones AM. Influence of acute dietary nitrate supplementation on 50 mile time trial performance in well-trained cyclists. Eur J Appl Physiol. 2012;112(12):4127–34.
- Thompson KG, Turner L, Prichard J, et al. Influence of dietary nitrate supplementation on physiological and cognitive responses to incremental cycle exercise. Respir Physiol Neurobiol. 2014;193:11–20.
- Wasserman K, Hansen JE, Sue DY, Whipp BJ, Casaburi R. Principles of exercise testing and interpretation. 3rd ed. Philadelphia: Lea & Deliger; 1999. p. 17–61, 153–154.
- Moore AN, Haun CT, Kephart WC, Holland AM, Mobley CB, Pascoe DD, Roberts MD, Martin JS. Red spinach extract increases ventilatory threshold during graded exercise testing. Sports. 2017;5(4):80.
- Fasuyi AO, Akindahunsi AO. Nutritional evaluation of Amaranthus cruentus leaf meal based broiler diets supplemented with cellulase/glucanase/ xylanase enzymes. Am J Food Technol. 2009;4:108–18.
- 27. Prakash D, Pal M. Nutritional and antinutritional composition of vegetable and grain amaranth leaves. J Sci Food Agric. 1991;57(4):573–83.
- 28. Subramanian D, Gupta S. Pharmacokinetic study of amaranth extract in healthy humans: a randomized trial. Nutrition. 2016;32(7–8):748–53.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

