



# Crystal structure, solvothermal synthesis, thermo-gravimetric studies and DFT calculations of a five-coordinate cobalt(II) compound based on the *N,N*-bis(2-hydroxyethyl)glycine anion

Yanling Zhou,<sup>a,b\*</sup> Xianrong Liu,<sup>a</sup> Qijun Wang,<sup>a</sup> Lisheng Wang<sup>a</sup> and Baoling Song<sup>a</sup>

Received 4 August 2016

Accepted 14 September 2016

Edited by A. J. Lough, University of Toronto, Canada

**Keywords:** Five-coordinate transition metal; crystal structure; solvothermal synthesis; mono-nuclear; ESI-MS; DFT.

**CCDC reference:** 1504322

**Supporting information:** this article has supporting information at journals.iucr.org/e


<sup>a</sup>School of Chemistry and Chemical Engineering, Guangxi University, Nanning, 530004, People's Republic of China, and <sup>b</sup>Guangxi Experiment Centre of Science and Technology, Guangxi University, Nanning, 530004, People's Republic of China. \*Correspondence e-mail: zyl8289@126.com

The reaction of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ , *N,N*-bis(2-hydroxyethyl)glycine and triethylamine ( $\text{Et}_3\text{N}$ ) in ethanol solution under solvothermal conditions produced crystals of [*N,N*-bis(2-hydroxyethyl)glycinato]chloridocobalt(II),  $[\text{Co}(\text{C}_6\text{H}_{12}\text{NO}_4)\text{Cl}]$ . The  $\text{Co}^{\text{II}}$  ion is coordinated in a slightly distorted trigonal-bipyramidal environment which is defined by three O atoms occupying the equatorial plane and the N and Cl atoms in the apical sites. In the crystal, two types of  $\text{O}-\text{H} \cdots \text{O}$  hydrogen bonds connect the molecules, forming a two-dimensional network parallel to (001). The molecular structure of the title compound confirms the findings of FTIR, elemental analysis, ESI-MS analysis and TG analysis. By using the density functional theory (DFT) (B3LYP) method with 6-31G(d) basis set, the molecular structure has been calculated and optimized.

## 1. Chemical context


In recent years, coordination compounds have attracted a great deal of interest for their structural aesthetics and potential functional applications (Fujita *et al.*, 2004). The design of molecular structures is highly influenced by factors such as the coordination nature of the metal ion, the coordinating ability and functionality of the organic ligands and the reaction conditions (Zhang *et al.*, 2015; Yin *et al.*, 2015). Hence, the prediction of crystal structure is largely considered to be serendipitous except for simple compounds such as mononuclear molecules. The  $3d^7$  configuration of  $\text{Co}^{\text{II}}$  is particularly suited for the construction of metal-organic compounds (Kurmo, 2009). One of the interesting structural aspects of studying cobalt compared to nickel, iron or manganese is the range of coordination geometries – octahedral, tetrahedral, square-pyramidal, trigonal-bipyramidal and square-planar – which are all stable (Kurmo, 2009). There are several coordination modes for the cobalt ion. The common mode is six-coordinate (Bryant *et al.*, 2015; Artetxe *et al.*, 2015), and only relatively few four-coordinate (Gupta *et al.*, 2015) and five-coordinate (Lee *et al.*, 2015) cobalt complexes have been recorded. Generally, five-coordinate compounds have two classical configurations, trigonal-bipyramidal and square-pyramidal, and the extent of each geometry each can be determined by the  $\tau$  value (Addison *et al.*, 1984).

The carboxylate unit is widely used in the synthesis of coordination compounds and is part of commonly used ligands. It is a good bridging group, favouring the formation of




OPEN ACCESS

products (Zhou *et al.*, 2009). Very recently, we have been investigating  $\text{Co}^{\text{II}}$  compounds constructed from ligands containing carboxylate and hydroxyl groups, which usually form multinuclear and/or polymeric structures and show interesting magnetic behavior (Zhou *et al.*, 2009; Zeng *et al.*, 2010). Similarly, herein, we chose *N,N*-bis(2-hydroxyethyl)glycine ( $\text{bicH}_3$ ) containing two hydroxyl oxygen atoms, one carboxylate oxygen atom and one nitrogen atom, which can potentially coordinate to a metal ion as a tetradentate ligand (He *et al.*, 1999).  $\text{bicH}_3$  contains the properties of both amino acid and amino alcohol as a result of the *N*-substituted amino, carboxyl, and two hydroxyl groups in the molecule. To the best of our knowledge, the crystal structures of metal–organic compounds with the  $\text{bicH}_3$  ligand have not been very well explored to date. Potential coordination modes for  $\text{bicH}_2^-$ ,  $\text{bicH}^{2-}$  and  $\text{bic}^{3-}$  are shown in Fig. 1. In the course of our ongoing studies on  $\text{Co}^{\text{II}}$  compounds containing ligands with carboxylate moieties, we have directly assembled the title compound  $[\text{Co}(\text{bicH}_2\text{Cl})]$ , **1**, using the flexible tetradentate ligand  $\text{bicH}_3$  and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  under solvothermal conditions.



## 2. Structural commentary

The molecular structure of the title compound is shown in Fig. 2. The  $\text{Co}^{\text{II}}$  ion is five-coordinated by four atoms from the  $\text{bicH}_2^-$  ligand (one carboxylate oxygen atom, two hydroxyl



**Figure 1**  
Coordination modes for  $\text{bicH}_2^-$  (a–e),  $\text{bicH}^{2-}$  (f–h) and  $\text{bic}^{3-}$  (i, j).

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).


$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O2}-\text{H2} \cdots \text{O4}^{\text{i}}$	0.85 (1)	1.79 (1)	2.6271 (19)	165 (2)
$\text{O1}-\text{H1} \cdots \text{O4}^{\text{ii}}$	0.79 (3)	1.89 (3)	2.6567 (19)	165 (3)

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{3}{2}, -z$ ; (ii)  $-x + 1, -y + 1, -z$ .


oxygen atoms, one nitrogen atom) and one terminal chlorine atom in a slightly distorted trigonal–bipyramidal environment ( $\tau = 0.94$ ,  $\tau = |\alpha - \beta|/60$ ,  $\alpha$  and  $\beta$  being the two largest angles around the central atom; values for  $\tau$  in perfect coordination geometries are 1.0 for trigonal–bipyramid and 0.0 for square–pyramidal). In a similar reported compound which was formed by  $\text{bicH}_2^{2-}$  and a  $\text{Cu}^{\text{II}}$  ion, a five-coordinate mode was observed (He *et al.*, 1999); the difference is that one nitrogen atom of benzimidazole or isoquinoline has replaced the terminal chloride ion in compound **1**. In **1**, the bond lengths around the  $\text{Co}^{\text{II}}$  ion are  $\text{Co1}-\text{N1} = 2.1626$  (15),  $\text{Co1}-\text{O1} = 2.0482$  (13),  $\text{Co1}-\text{O2} = 2.0463$  (14),  $\text{Co1}-\text{O3} = 2.0095$  (14) and  $\text{Co1}-\text{Cl1} = 2.2701$  (6)  $\text{\AA}$ . The length of the  $\text{Co}-\text{O}$  (carboxylate) bond is shorter than that of  $\text{Co}-\text{O}$  (hydroxyl), which may be due to the difference between the electron density of carboxylate oxygen atoms and that of hydroxyl oxygen atoms (He *et al.*, 1999). According to the total valence–charge balance and the bond lengths, we can conclude that cobalt is in oxidation state +II.

## 3. Supramolecular features

In the crystal, two types of  $\text{O}-\text{H} \cdots \text{O}$  hydrogen bonds (Table 1) connect the molecules, forming a two-dimensional network parallel to (001) (Fig. 3). The  $\text{O}-\text{H}$  groups behave as donors to the non-coordinating carboxylate oxygen atom of



**Figure 2**  
The molecular structure of the title compound, showing the atom labeling. Displacement ellipsoids are drawn at the 30% probability level.



**Figure 3**  
Part of the crystal structure showing the two different O—H...O hydrogen bonds as distinct colors, blue for O1—H...O4<sup>ii</sup> bonds and green for O2—H2...O4<sup>i</sup> bonds (symmetry codes as in Table 1).


symmetry-related molecules. The hydroxy group containing O1 acts as a bifurcated O—H...O(O,O) donor while carboxylate atom O4 is a bifurcated (O—H,O—H)...O acceptor.

#### 4. Database survey

A search of the Cambridge Structural Database (CSD; Groom *et al.*, 2016) was carried out for structures containing the bicH<sub>3</sub> ligand. This revealed bicH<sub>3</sub> coordinating to a lanthanide metal (Inomata *et al.*, 2001), Cd and Na (Katsoulakou *et al.*, 2011), Cu, Ni and Zn (Thakuria & Das, 2007; Liu *et al.*, 2013; Lo & Ng, 2010), Re, Mn and Fe (Kirillov *et al.*, 2005; Sun *et al.*, 1997; Graham *et al.*, 2009). A related structure with copper and bromide (Yamaguchi *et al.*, 1991) shows a very similar mononuclear crystal structure to the title compound. There are only a small number of reports for the ligand coordinating to Co (Funes *et al.*, 2015; Zhao & Liu, 2010; Liu *et al.*, 2015).



**Figure 4**  
The optical microscope image of single crystals of compound **1**.



**Figure 5**  
ESI mass spectrum of compound **1**.

#### 5. Synthesis and crystallization

The ligand bicH<sub>3</sub> (0.5 mmol) in a ethanol solution (2 mL) was added to a ethanol solution (5 mL) of CoCl<sub>2</sub>·6H<sub>2</sub>O (1 mmol). 0.02 mL of triethylamine was added dropwise to the mixed solution and stirred for 15 min at room temperature. The reactants were sealed in a 12 mL Teflon-lined autoclave, heated at 413 K for three days and then cooled to room temperature at a rate of 10 K h<sup>-1</sup>. Purple single crystals (Fig. 4) were obtained along with purple powder. The crystals were picked out, washed with distilled water, and dried in air (yield *ca* 50.3% based on Co<sup>II</sup>). Analysis calculated (%) for C<sub>6</sub>H<sub>12</sub>ClCoNO<sub>4</sub>: C 28.30, H 3.93, N 5.50; Found C 28.31, H 3.95, N 5.54%. FTIR data for **1** (KBr, cm<sup>-1</sup>): 3383(*m*), 2964(*w*), 1593(*s*), 1434(*m*), 1407(*m*), 1309(*w*), 1058(*w*), 890(*w*).

#### 6. ESI–MS spectroscopic analysis

The ESI mass spectra were recorded using an LCQ–FLEET mass spectrometer (Thermo). To give further evidence for the inner structure of compound **1**, characterization of the molecule in solution was accomplished by ESI–MS experiments. For the methanol solution of **1**, the ESI mass spectrum (Fig. 5) exhibits the main ion peak observed at an *m/z* of 254.93, which can be assigned as [Co(C<sub>6</sub>H<sub>11</sub>NO<sub>4</sub>)Cl]<sup>+</sup> (fit: 254.970860). The observed *m/z* clearly matches the assigned formula as well as the simulated spectra. This suggests that compound **1** produced in solution was stable during the ionization process. ESI–MS can also be used to examine a series of inner-bridge replacement reactions for multinuclear Co compounds (Zhou *et al.*, 2010; Hu *et al.*, 2013), which is an important complement to ligand exchange, ion exchange, template exchange and supramolecular transformations (Chakrabarty *et al.*, 2011; Miras *et al.*, 2009). The study of the chemistry of coordination compounds by mass spectroscopy is an excellent tool to demonstrate the stability and existence of multinuclear molecules in solution.




Figure 6  
The TG curve of compound **1**.

## 7. TG analysis

The TG analysis was performed on Pyris Diamond TG/DTA. The appearance of the flexible polydentate ligands inspired us to investigate the thermal stability of the network. The crushed single-crystal samples were heated to 1073 K in an  $N_2$  atmosphere at a heating rate of  $5\text{ K min}^{-1}$  (Fig. 6). The TGA curve for **1** shows that the framework begins to decompose at 413 K, and the 21.6% remaining weight is assuming to the mass loss percentage of cobalt (cal. 22.9%). One similar compound,  $[CuCd(bicH_2(NO_3)Cl_2(H_2O))] \cdot H_2O$ , constructed with  $bicH_2$  is quite unstable and begins to lose lattice water at 393 K (Liu *et al.*, 2013). Other compounds synthesized using the same ligand also show mass loss below 413 K (Inomata *et al.*, 2001), owing to the loss of coordinating water. However, another reported complex with five-coordinate cobalt begins to disintegrate at 669 K (Lee *et al.*, 2015). In summary, the crystals synthesized using  $bicH_2^{2-}$  tend to decompose at a relatively low temperature.

## 8. DFT calculations

All the calculations were performed by using the *GAUSSIAN09* program package (Frisch *et al.*, 2009). The molecular structure of the title compound in the ground state was optimized *in vacuo* without considering the solvent, and a basis set (b1) consisting of a standard LanL2DZ basis set (Dunning & Hay, 1976; Wadt & Hay, 1985; Hay & Wadt, 1985) for Co, while the other atoms, C, H, N, O were described by a standard 6-31G(d) set. To investigate the energy differences between the high-spin and low-spin states of the title compound, the  $\Delta E$  of these two energy states was evaluated using the B3LYP/b1 method (Carabineiro *et al.*, 2008; Saraçoğlu & Cukurovali, 2016). Vibrational frequencies were calculated for all stationary points to verify that each was a minimum (NIMAG = 0) on the potential energy surface.

To gain an insight of the electronic structures, bonding properties and relative stability of the two different spin

ground states ( $S = 3/2$  and  $1/2$ ) for compound **1**, the calculations in the DFT method were investigated. The optimized geometries calculated for compound **1** are presented in Fig. 7. The structure of **1** presented an almost perfect trigonal-bipyramidal geometry by means of the X-ray diffraction. The chlorine and nitrogen atoms occupy the axial positions, while the equatorial plane is occupied by three oxygen atoms. According to the energies for the two calculated structures (see Supporting information), the X-ray structure determined for complex **1** should correspond to the complex with high-spin ground state. The result can also be indicated by the mean ( $\delta$ ) and maximum ( $\Delta$ ) absolute deviations obtained for the coordination distances (Co–X). For the high-spin form of complex **1**, values of  $\delta = 0.104\text{ \AA}$  and  $\Delta = 0.148\text{ \AA}$  indicate a reasonable agreement. As for the high-spin ground state and the experimental value, the biggest difference for the bond lengths is found to be  $0.148\text{ \AA}$  for Co–N, similar to what has been observed in related  $Co^{II}$  compounds (Carabineiro *et al.*, 2008). The results of the schematic representation of both ground states supported its coordination behavior and the value of  $\Delta E$  is  $13.4\text{ kcal mol}^{-1}$ , which shows that compound **1** can well exist stably. The cartesian coordinates for the two calculated structures are given in the Supporting information.

## 9. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms bonded to C atoms were placed in calculated positions with  $C-H = 0.97\text{ \AA}$  and  $U_{iso}(H) = 1.2U_{eq}(C)$ . Hydroxyl hydrogen atoms H1 and H2 were




Figure 7  
Optimized geometries (B3LYP) for the low-spin ( $S = 1/2$ , bottom) and the high-spin ( $S = 3/2$ , top) ground states and the relative energy of the two ground states ( $\text{kcal mol}^{-1}$ ).



**Table 2**  
Experimental details.

Crystal data	
Chemical formula	[Co(C <sub>6</sub> H <sub>12</sub> NO <sub>4</sub> )Cl]
<i>M<sub>r</sub></i>	256.55
Crystal system, space group	Orthorhombic, <i>Pbca</i>
Temperature (K)	296
<i>a</i> , <i>b</i> , <i>c</i> (Å)	8.3925 (9), 14.0939 (15), 15.8448 (17)
<i>V</i> (Å <sup>3</sup> )	1874.2 (3)
<i>Z</i>	8
Radiation type	Mo <i>K</i> α
<i>μ</i> (mm <sup>-1</sup> )	2.10
Crystal size (mm)	0.84 × 0.27 × 0.24
Data collection	
Diffractionmeter	Bruker APEXII CCD
Absorption correction	Multi-scan ( <i>SADABS</i> ; Bruker, 2004)
<i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>	0.512, 0.604
No. of measured, independent and observed [ <i>I</i> > 2σ( <i>I</i> )] reflections	15130, 1933, 1854
<i>R<sub>int</sub></i>	0.021
(sin θ/λ) <sub>max</sub> (Å <sup>-1</sup> )	0.625
Refinement	
<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.025, 0.066, 1.14
No. of reflections	1933
No. of parameters	125
No. of restraints	3
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
Δ <i>ρ</i> <sub>max</sub> , Δ <i>ρ</i> <sub>min</sub> (e Å <sup>-3</sup> )	0.51, -0.43

Computer programs: *APEX2* and *SAINT* (Bruker, 2004), *SIR2004* (Burla *et al.*, 2007), *SHELXL2014* (Sheldrick, 2015) and *OLEX2* (Dolomanov *et al.*, 2009).

refined independently, H1 with a refined isotropic displacement parameter and H2 with *U*<sub>iso</sub>(H) = 1.5*U*<sub>eq</sub>(O).

## Acknowledgements

This work was supported by the NSFC (No. 21401030), GXNSF (No. 2014GXNSFBA118048), as well as the Guangxi Experiment Centre of Science and Technology (YXKT2014008). The computational resources are partly provided by Multifunction Computer Center of Guangxi University.

## References

Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.  
 Artetxe, B., Reinoso, S., San Felices, L., Vitoria, P., Pache, A., Martín-Caballero, J. & Gutiérrez-Zorrilla, J. M. (2015). *Inorg. Chem.* **54**, 241–252.  
 Bruker (2004). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bryant, M. R., Burrows, A. D., Fitchett, C. M., Hawes, C. S., Hunter, S. O., Keenan, L. L., Kelly, D. J., Kruger, P. E., Mahon, M. F. & Richardson, C. (2015). *Dalton Trans.* **44**, 9269–9280.  
 Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Casciarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G., Siliqi, D. & Spagna, R. (2007). *J. Appl. Cryst.* **40**, 609–613.

Carabineiro, S. A., Bellabarba, R. M., Gomes, P. T., Pascu, S. I., Veiros, L. F., Freire, C., Pereira, L. C. J., Henriques, R. T., Oliveira, M. C. & Warren, J. E. (2008). *Inorg. Chem.* **47**, 8896–8911.  
 Chakrabarty, R., Mukherjee, P. S. & Stang, P. J. (2011). *Chem. Rev.* **111**, 6810–6918.  
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.  
 Dunning, T. H. Jr & Hay, P. J. (1976). *Modern Theoretical Chemistry*, Vol. 3, edited by H. F. Schaefer III, p. 1. New York: Plenum.  
 Frisch, M. J., *et al.* (2009). *GAUSSIAN09*. Gaussian, Inc., Wallingford, CT.  
 Fujita, M., Powell, A. & Creutz, C. E. (2004). *Creutz Comprehensive Coordination Chemistry II: From Biology to Nanotechnology*, Vol. 7, pp. 1–56 Oxford: Elsevier.  
 Funes, A. V., Carrella, L., Sorace, L., Rentschler, E. & Alborés, P. (2015). *Dalton Trans.* **44**, 2390–2400.  
 Graham, K., Darwish, A., Ferguson, A., Parsons, S. & Murrie, M. (2009). *Polyhedron*, **28**, 1830–1833.  
 Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst. B72*, 171–179.  
 Gupta, S. K., Kuppuswamy, S., Walsh, J. P., McInnes, E. J. & Murugavel, R. (2015). *Dalton Trans.* **44**, 5587–5601.  
 Hay, P. J. & Wadt, W. R. (1985). *J. Chem. Phys.* **82**, 270–283, 299–310.  
 He, X. F., Long, L. S., Le, X. Y., Chen, X. M., Ji, L. N. & Zhou, Z. Y. (1999). *Inorg. Chim. Acta*, **285**, 326–331.  
 Hu, Y. Q., Zeng, M. H., Zhang, K., Hu, S., Zhou, F. F. & Kurmoo, M. (2013). *J. Am. Chem. Soc.* **135**, 7901–7908.  
 Inomata, Y., Takei, T. & Howell, F. S. (2001). *Inorg. Chim. Acta*, **318**, 201–206.  
 Katsoulakou, E., Konidaris, K. F., Terzis, A., Raptopoulou, C. P., Perlepes, S. P., Manessi-Zoupa, E. & Kostakis, G. E. (2011). *Polyhedron*, **30**, 397–404.  
 Kirillov, A. M., Haukka, M., Kirillova, M. V. & Pombeiro, A. L. (2005). *Adv. Synth. Catal.* **347**, 1435–1446.  
 Kurmoo, M. (2009). *Chem. Soc. Rev.* **38**, 1353–1379.  
 Lee, J. Y., Lee, J. Y. & Lee, H. M. (2015). *Inorg. Chem. Commun.* **52**, 16–19.  
 Liu, Y., Kuang, D. Z., Feng, Y. L. & Fu, W. W. (2013). *Transition Met. Chem.* **38**, 849–853.  
 Liu, Y., Zhou, D., Liu, H.-H. & He, C.-C. (2015). *Acta Cryst. E71*, m199–m200.  
 Lo, K. M. & Ng, S. W. (2010). *Acta Cryst. E66*, m1485.  
 Miras, H. N., Wilson, E. F. & Cronin, L. (2009). *Chem. Commun.* pp. 1297–1311.  
 Saraçoğlu, H. & Cukurovali, A. (2016). *Mol. Cryst. Liq. Cryst.* **625**, 173–185.  
 Sheldrick, G. M. (2015). *Acta Cryst. C71*, 3–8.  
 Sun, Z., Gantzel, P. K. & Hendrickson, D. N. (1997). *Polyhedron*, **16**, 3267–3271.  
 Thakuria, H. & Das, G. (2007). *Polyhedron*, **26**, 149–153.  
 Wadt, W. R. & Hay, P. J. (1985). *J. Chem. Phys.* **82**, 284–298.  
 Yamaguchi, H., Inomata, Y. & Takeuchi, T. (1991). *Inorg. Chim. Acta*, **181**, 31–36.  
 Yin, Z., Zhou, Y. L., Zeng, M. H. & Kurmoo, M. (2015). *Dalton Trans.* **44**, 5258–5275.  
 Zeng, M. H., Zhou, Y. L., Wu, M. C., Sun, H. L. & Du, M. (2010). *Inorg. Chem.* **49**, 6436–6442.  
 Zhang, W. X., Liao, P. Q., Lin, R. B., Wei, Y. S., Zeng, M. H. & Chen, X. M. (2015). *Coord. Chem. Rev.* **293–294**, 263–278.  
 Zhao, J.-P. & Liu, F.-C. (2010). *Acta Cryst. E66*, m848.  
 Zhou, Y. L., Meng, F. Y., Zhang, J., Zeng, M. H. & Liang, H. (2009). *Cryst. Growth Des.* **9**, 1402–1410.  
 Zhou, Y. L., Zeng, M. H., Wei, L. Q., Li, B. W. & Kurmoo, M. (2010). *Chem. Mater.* **22**, 4295–4303.

## supporting information

*Acta Cryst.* (2016). E72, 1463-1467 [doi:10.1107/S2056989016014596]

## Crystal structure, solvothermal synthesis, thermogravimetric studies and DFT calculations of a five-coordinate cobalt(II) compound based on the *N,N*-bis(2-hydroxyethyl)glycine anion

Yanling Zhou, Xianrong Liu, Qijun Wang, Lisheng Wang and Baoling Song

### Computing details

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINTE* (Bruker, 2004); data reduction: *SAINTE* (Bruker, 2004); program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2007); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).

### [*N,N*-Bis(2-hydroxyethyl)glycinato]chloridocobalt(II)

#### Crystal data

[Co(C<sub>6</sub>H<sub>12</sub>NO<sub>4</sub>)Cl]  
 $M_r = 256.55$   
 Orthorhombic, *Pbca*  
 $a = 8.3925$  (9) Å  
 $b = 14.0939$  (15) Å  
 $c = 15.8448$  (17) Å  
 $V = 1874.2$  (3) Å<sup>3</sup>  
 $Z = 8$   
 $F(000) = 1048$

$D_x = 1.818$  Mg m<sup>-3</sup>  
 Mo *K*α radiation,  $\lambda = 0.71073$  Å  
 Cell parameters from 9897 reflections  
 $\theta = 2.6$ – $26.4^\circ$   
 $\mu = 2.10$  mm<sup>-1</sup>  
 $T = 296$  K  
 Prism, purple  
 $0.84 \times 0.27 \times 0.24$  mm

#### Data collection

Bruker APEXII CCD  
 diffractometer  
 Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2004)  
 $T_{\min} = 0.512$ ,  $T_{\max} = 0.604$   
 15130 measured reflections

1933 independent reflections  
 1854 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$   
 $\theta_{\max} = 26.4^\circ$ ,  $\theta_{\min} = 2.6^\circ$   
 $h = -10 \rightarrow 10$   
 $k = -17 \rightarrow 17$   
 $l = -19 \rightarrow 19$

#### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.025$   
 $wR(F^2) = 0.066$   
 $S = 1.14$   
 1933 reflections  
 125 parameters  
 3 restraints

Primary atom site location: structure-invariant  
 direct methods  
 Hydrogen site location: mixed  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.030P)^2 + 1.540P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.51 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.41813 (3)	0.62528 (2)	0.09419 (2)	0.02152 (10)
Cl1	0.15183 (6)	0.60446 (4)	0.07726 (4)	0.04084 (15)
O1	0.46124 (15)	0.51884 (10)	0.17919 (8)	0.0254 (3)
O2	0.42621 (16)	0.75436 (10)	0.15315 (9)	0.0315 (3)
H2	0.3575 (17)	0.7980 (10)	0.1454 (15)	0.047*
O3	0.48651 (16)	0.61778 (10)	-0.02717 (8)	0.0320 (3)
O4	0.68239 (18)	0.63508 (9)	-0.11936 (9)	0.0305 (3)
N1	0.67257 (18)	0.64315 (11)	0.10881 (9)	0.0226 (3)
C1	0.6259 (2)	0.50535 (15)	0.20204 (13)	0.0319 (4)
H1A	0.6481	0.4381	0.2078	0.038*
H1B	0.6469	0.5356	0.2559	0.038*
C2	0.7323 (2)	0.54745 (14)	0.13522 (13)	0.0308 (4)
H2A	0.8398	0.5533	0.1571	0.037*
H2B	0.7353	0.5056	0.0866	0.037*
C3	0.5816 (2)	0.79707 (15)	0.15957 (15)	0.0367 (5)
H3A	0.5852	0.8407	0.2069	0.044*
H3B	0.6066	0.8318	0.1084	0.044*
C4	0.6996 (2)	0.71678 (15)	0.17250 (13)	0.0336 (4)
H4A	0.8075	0.7408	0.1677	0.040*
H4B	0.6865	0.6901	0.2285	0.040*
C5	0.6287 (2)	0.63779 (12)	-0.04522 (12)	0.0240 (4)
C6	0.7392 (2)	0.66736 (15)	0.02554 (11)	0.0301 (4)
H6A	0.8412	0.6359	0.0186	0.036*
H6B	0.7571	0.7353	0.0226	0.036*
H1	0.432 (3)	0.468 (2)	0.1653 (18)	0.054 (9)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.01510 (14)	0.02795 (15)	0.02151 (15)	-0.00134 (9)	-0.00195 (9)	0.00217 (9)
Cl1	0.0166 (2)	0.0604 (3)	0.0455 (3)	-0.0060 (2)	-0.0050 (2)	0.0169 (3)
O1	0.0233 (6)	0.0265 (7)	0.0265 (7)	-0.0020 (5)	-0.0014 (5)	0.0011 (5)
O2	0.0269 (7)	0.0294 (7)	0.0383 (8)	0.0047 (5)	-0.0040 (6)	-0.0023 (6)
O3	0.0217 (7)	0.0523 (9)	0.0220 (7)	-0.0095 (6)	-0.0014 (5)	-0.0024 (6)
O4	0.0336 (7)	0.0355 (7)	0.0223 (7)	-0.0109 (6)	0.0038 (6)	-0.0020 (5)
N1	0.0169 (7)	0.0310 (8)	0.0199 (7)	-0.0025 (6)	-0.0026 (6)	0.0015 (6)
C1	0.0263 (9)	0.0363 (10)	0.0330 (10)	0.0024 (8)	-0.0079 (8)	0.0062 (8)
C2	0.0228 (9)	0.0359 (10)	0.0337 (10)	0.0036 (8)	-0.0030 (8)	0.0022 (8)

C3	0.0367 (11)	0.0290 (10)	0.0445 (12)	-0.0038 (8)	-0.0073 (9)	-0.0057 (9)
C4	0.0280 (10)	0.0406 (11)	0.0323 (10)	-0.0047 (8)	-0.0069 (8)	-0.0066 (8)
C5	0.0247 (8)	0.0235 (8)	0.0238 (9)	-0.0031 (7)	-0.0006 (7)	0.0009 (7)
C6	0.0217 (8)	0.0443 (11)	0.0243 (9)	-0.0090 (8)	-0.0001 (7)	0.0027 (8)

*Geometric parameters (Å, °)*

Co1—C11	2.2701 (6)	C1—H1A	0.9700
Co1—O1	2.0482 (13)	C1—H1B	0.9700
Co1—O2	2.0463 (14)	C1—C2	1.507 (3)
Co1—O3	2.0095 (14)	C2—H2A	0.9700
Co1—N1	2.1626 (15)	C2—H2B	0.9700
O1—C1	1.441 (2)	C3—H3A	0.9700
O1—H1	0.79 (3)	C3—H3B	0.9700
O2—H2	0.852 (9)	C3—C4	1.518 (3)
O2—C3	1.440 (2)	C4—H4A	0.9700
O3—C5	1.259 (2)	C4—H4B	0.9700
O4—C5	1.259 (2)	C5—C6	1.514 (3)
N1—C2	1.498 (2)	C6—H6A	0.9700
N1—C4	1.465 (2)	C6—H6B	0.9700
N1—C6	1.473 (2)		
O1—Co1—C11	99.03 (4)	C2—C1—H1B	109.7
O1—Co1—N1	80.82 (6)	N1—C2—C1	110.66 (16)
O2—Co1—C11	101.63 (4)	N1—C2—H2A	109.5
O2—Co1—O1	110.19 (6)	N1—C2—H2B	109.5
O2—Co1—N1	79.33 (6)	C1—C2—H2A	109.5
O3—Co1—C11	99.28 (4)	C1—C2—H2B	109.5
O3—Co1—O1	122.70 (6)	H2A—C2—H2B	108.1
O3—Co1—O2	118.31 (6)	O2—C3—H3A	110.4
O3—Co1—N1	80.02 (6)	O2—C3—H3B	110.4
N1—Co1—C11	179.02 (5)	O2—C3—C4	106.80 (16)
Co1—O1—H1	115 (2)	H3A—C3—H3B	108.6
C1—O1—Co1	115.54 (11)	C4—C3—H3A	110.4
C1—O1—H1	104 (2)	C4—C3—H3B	110.4
Co1—O2—H2	123.6 (13)	N1—C4—C3	109.51 (16)
C3—O2—Co1	115.72 (12)	N1—C4—H4A	109.8
C3—O2—H2	108.8 (12)	N1—C4—H4B	109.8
C5—O3—Co1	118.44 (12)	C3—C4—H4A	109.8
C2—N1—Co1	104.78 (11)	C3—C4—H4B	109.8
C4—N1—Co1	108.01 (12)	H4A—C4—H4B	108.2
C4—N1—C2	113.16 (15)	O3—C5—C6	118.30 (16)
C4—N1—C6	113.21 (15)	O4—C5—O3	123.00 (17)
C6—N1—Co1	107.81 (11)	O4—C5—C6	118.69 (17)
C6—N1—C2	109.38 (15)	N1—C6—C5	111.54 (15)
O1—C1—H1A	109.7	N1—C6—H6A	109.3
O1—C1—H1B	109.7	N1—C6—H6B	109.3
O1—C1—C2	109.85 (15)	C5—C6—H6A	109.3



H1A—C1—H1B	108.2	C5—C6—H6B	109.3
C2—C1—H1A	109.7	H6A—C6—H6B	108.0
Co1—O1—C1—C2	-21.0 (2)	O3—C5—C6—N1	15.5 (3)
Co1—O2—C3—C4	-35.2 (2)	O4—C5—C6—N1	-165.10 (16)
Co1—O3—C5—O4	179.97 (13)	C2—N1—C4—C3	-156.64 (17)
Co1—O3—C5—C6	-0.7 (2)	C2—N1—C6—C5	92.75 (18)
Co1—N1—C2—C1	-44.47 (17)	C4—N1—C2—C1	73.0 (2)
Co1—N1—C4—C3	-41.12 (19)	C4—N1—C6—C5	-140.06 (17)
Co1—N1—C6—C5	-20.64 (19)	C6—N1—C2—C1	-159.82 (15)
O1—C1—C2—N1	44.4 (2)	C6—N1—C4—C3	78.2 (2)
O2—C3—C4—N1	49.9 (2)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O2—H2 $\cdots$ O4 <sup>i</sup>	0.85 (1)	1.79 (1)	2.6271 (19)	165 (2)
O1—H1 $\cdots$ O4 <sup>ii</sup>	0.79 (3)	1.89 (3)	2.6567 (19)	165 (3)

Symmetry codes: (i)  $x-1/2, -y+3/2, -z$ ; (ii)  $-x+1, -y+1, -z$ .