



High serum leptin levels are associated with central arterial stiffness in geriatric patients on hemodialysis

Chiu-Huang Kuo^a, Yu-Li Lin^a, Chih-Hsien Wang^a, Yu-Hsien Lai^a, Ru-Jiang Syu^a, Bang-Gee Hsu^{a, b*}

^aDivision of Nephrology, Buddhist Tzu Chi General Hospital, Hualien, Taiwan,
^bSchool of Medicine, Tzu Chi University, Hualien, Taiwan

ABSTRACT

Objective: Central arterial stiffness predicts cardiovascular (CV) mortality in hemodialysis (HD) patients. The aging process transforms lipid distribution and thus alters adipokine secretion. The harmful effects of leptin on CV events may change in the elderly. The purpose of this study was to investigate the relationship between leptin and central arterial stiffness markers through carotid–femoral pulse wave velocity (cfPWV) in geriatric HD patients. **Materials and Methods:** Patients over 65 years old on chronic HD were recruited. Blood samples were collected, and the cfPWV was measured with the SphygmoCor system. The patients with cfPWV values >10 m/s were defined as the high arterial stiffness group. **Results:** In total, 30 (51.7%) of the 58 geriatric patients on chronic HD in this study were in the high arterial stiffness group. The high arterial stiffness group had higher rates of diabetes mellitus ($P = 0.019$), hypertension ($P = 0.019$), and higher systolic blood pressure ($P = 0.018$), pulse pressure ($P = 0.019$), body mass index ($P = 0.018$), serum leptin levels ($P = 0.008$), and hemoglobin levels ($P = 0.040$) than those in the low arterial stiffness group. Multivariable forward stepwise linear regression analysis showed logarithmically transformed leptin (log-leptin, $\beta = 0.408$, adjusted R^2 change = 0.164; $P = 0.001$) and diabetes ($\beta = 0.312$, adjusted R^2 change = 0.085; $P = 0.009$) were associated with cfPWV values in geriatric HD patients. Moreover, an increased serum leptin level (odds ratio: 1.053; 95% confidence interval: 1.007–1.100; $P = 0.023$) was an independent factor for central arterial stiffness among geriatric HD patients after multivariate logistic regression analysis. **Conclusion:** In this study, a higher serum leptin level was correlated with central arterial stiffness in geriatric HD patients.

KEYWORDS: Carotid–femoral pulse wave velocity, Geriatric, Hemodialysis, Leptin

Received : 06-Mar-2017
Revised : 10-May-2017
Accepted : 14-Jul-2017

INTRODUCTION

Patients over 65 years old are a growing group worldwide in the chronic dialysis population [1]. The number of patients with prevalent end-stage renal disease (ESRD) continues to increase in all age groups, with the relative magnitude of increase greater in older age groups [2]. Older dialysis patients usually have numerous cardiovascular (CV) risks including diabetes, hypertension, and dyslipidemia, and atypical risks such as hyperhomocysteinemia and chronic inflammation [3]. Therefore, measurement of arterial elastic properties is becoming a method of risk stratification.

Central arterial stiffness is recognized as a surrogate end point for CV diseases and is the functional consequence of artery wall calcification. In the ESRD population, this disorder is closely linked to disturbances of calcium and phosphate homeostasis and to bone metabolism irregularities. Carotid–femoral pulse wave velocity (cfPWV) is a valid,

noninvasive, and inexpensive technique for measuring central arterial stiffness [4]. Independent of the classic CV risk factors, a higher cfPWV level represents a strong predictor of the future CV events and all-cause mortality, and the predictive ability is superior in people with a higher baseline CV risk [5,6].

Leptin is secreted by adipose tissue and the level is proportional to the body fat mass. It has a well-documented function in glucose homeostasis, insulin sensitivity, and energy expenditure [7]. An association of leptin and CV disease has been reported in conditions including hypertension, the atherosclerotic process, and endothelial cell dysfunction through direct sympathetic system stimulation, dysregulated vascular smooth

*Address for correspondence:

Dr. Bang-Gee Hsu,
Division of Nephrology, Buddhist Tzu Chi General Hospital,
707, Section 3, Chung-Yang Road, Hualien, Taiwan.
E-mail: geelily@tzuchi.com.tw

Access this article online

Quick Response Code:



Website: www.tcmjmed.com

DOI: 10.4103/tcmj.tcmj_10_18

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Kuo CH, Lin YL, Wang CH, Lai YH, Syu RJ, Hsu BG. High serum leptin levels are associated with central arterial stiffness in geriatric patients on hemodialysis. Tzu Chi Med J 2018;30(4):227-32.

muscle, and increased oxidative stress [8-10]. In the aging process, alterations in adipose cell size, number, and proliferation lead to adipokine dysregulation [11,12]. However, the association between leptin and central arterial stiffness in geriatric hemodialysis (HD) patients remains to be explored. The aim of the present study was to assess the association between serum leptin levels and central arterial stiffness measured by cfPWV in geriatric HD patients.

MATERIALS AND METHODS

Patients

The study participants were recruited in the HD department at Buddhist Tzu Chi General Hospital, Hualien, Taiwan, in mid-2015. The study was approved by the Protection of Human Subjects Institutional Review Board of Tzu Chi University and Hospital (IRB103-136-B). A total of 158 HD participants were enrolled and signed informed consents. Of these, 100 were excluded because of an HD duration <6 months ($n = 2$), each HD session >4 h ($n = 10$), each HD session <4-h ($n = 22$), HD 2 times per week ($n = 2$), acute infection ($n = 1$), acute myocardial infarction ($n = 1$), pulmonary edema ($n = 1$), missing clinical data ($n = 7$), and age under 65 years ($n = 54$). Finally, a total of 58 HD patients older than 65 years were included in the study. All patients in the study used the same high-flux polysulfone disposable artificial kidney (FX class dialyzer, Fresenius Medical Care, Bad Homburg, Germany), received standard 4-h dialysis 3 times/week using standard bicarbonate dialysate and had been on HD for more than half year. There were 27 men, and 31 women, with ages ranging from 66 to 92 years old. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were taken at the points of appearance and disappearance of the Korotkoff sounds before the start of the HD session. Pulse pressure was calculated by subtracting the DBP from the SBP.

Anthropometric analysis

All anthropometric analyses were performed at the midweek dialysis. Body height was measured to the nearest half centimeter, and body weight was measured to the nearest half-kilogram with the patient in light clothing and no shoes before and after the HD session. Waist circumference was measured to the nearest half centimeter at the shortest point below the lower rib margin and the iliac crest. Body mass index (BMI) was calculated as the weight (kilograms) divided by the height squared (meters) [13-16].

Biochemical investigations

All blood samplings were performed at the midweek dialysis. Blood samples of approximately 5 mL were collected just before the start of the HD session; about 0.5 mL was used to check hemoglobin levels and white blood cell counts (Sysmex K-1000, Sysmex American, Mundelein, IL, USA); the rest were immediately centrifuged at 3000 \times g for 10 min. Serum samples were stored at 4°C and used for biochemical analyses within 1 h after collection. Serum levels of creatinine, blood urea nitrogen (predialysis urea), glucose, total cholesterol, triglycerides (TG), iron, total iron-binding capacity, ferritin, total calcium, and phosphorus were measured through auto-analyzer (COBAS Integra 800, Roche Diagnostics, Basel,

Switzerland). Serum leptin (SPI-BIO, Montigny le Bretonneux, France) concentrations were determined using a commercially available enzyme immunoassay [13-16]. Serum intact parathyroid hormone levels (iPTH) were measured using enzyme-linked immunosorbent assays (Diagnostic Systems Laboratories, Webster, Texas, USA) [13-16]. At the end of the HD session, ultrafiltration was stopped, blood flow was slowed to 100 mL/min for 20 s, and blood samples were drawn from the arterial blood line sampling port for postdialysis serum urea measurements. The urea reduction rate was calculated as $([\text{predialysis urea} - \text{postdialysis urea}] \times 100) / [\text{predialysis urea}]$. The adequacy of dialysis was estimated by the measurement of the single-pool fractional clearance index for urea (Kt/Vurea) using Gotch's formula.

Carotid-femoral pulse wave velocity measurements

The cfPWV was measured transcutaneously with a pressure tonometer (SphygmoCor system, AtCor Medical, West Ryde, Australia) on the carotid and femoral sites, and the pressure pulse waveform was recorded, as previously described [14,15]. These measurements were performed in all patients in the supine position after a minimum 10 min rest in a quiet, temperature-controlled room before the start of the midweek HD session. Recordings were made simultaneously with an electrocardiographic (ECG) signal, which provided an R-timing reference. Pulse wave recordings were performed consecutively at the carotid and femoral segments. The integrated software (SphygmoCor system, AtCor Medical, West Ryde, Australia) was used to process each set of pulse wave and ECG data, to calculate the mean time difference between the R-wave and the pulse wave on a beat-to-beat basis, among an average of 10 consecutive cardiac cycles. The cfPWV was calculated using the distance and the mean time difference between the two recorded points (carotid-femoral segment). In this study, patients with cfPWV values above 10 m/s were defined as the high arterial stiffness group, while those with values below 10 m/s were regarded as the low arterial stiffness group. The cut-off value was determined according to the European Society of Hypertension and European Society of Cardiology guidelines in 2013 [17].

Statistical analysis

Data were tested with the Kolmogorov-Smirnov test for normal distribution. Data were expressed as means \pm standard deviation for normally distributed results and as medians and interquartile ranges for nonnormally distributed data. Comparisons between patients were performed using Student's independent *t*-test (two-tailed) for normally distributed data, or the Mann-Whitney U-test for parameters with a nonnormal distribution. The glucose, ferritin, TG, iPTH, leptin, and HD duration datasets showed skewed nonnormal distributions, and therefore were recalculated by logarithm transformation to base 10; after this transformation, the log-glucose, log-ferritin, log-TG, log-iPTH, log-leptin, and log-HD duration were normally distributed. Data expressed as the number of patients were analyzed by the Chi-square test. Clinical variables that correlated with cfPWV values in geriatric HD patients were first evaluated by univariable linear regression analysis. Variables that were significantly associated with cfPWV values in geriatric HD patients were tested for independence

by multivariate forward stepwise regression analysis (adopted factors: diabetes, age, waist circumference, BMI, body fat mass, post-HD body weight, SBP, pulse pressure, hemoglobin, log-TG, and log-leptin). Variables that were significantly associated with central arterial stiffness in HD patients were tested for independence by multivariate logistic regression analysis (adopted factors: Diabetes, hypertension, waist circumference, BMI, pulse pressure, hemoglobin, and leptin). All statistical analyses employed SPSS for Windows (version 19.0; SPSS Inc., Chicago, IL, USA). The $P < 0.05$ was considered statistically significant.

RESULTS

Demographic, biochemical, and clinical characteristics of the 58 geriatric HD patients are shown in Table 1. The mean age of participants was 75.14 years with a median HD duration of 4.14 years. Men accounted for 46.5% of participants. A total of 34 patients (58.6%) had diabetes mellitus and 33 (56.9%) had hypertension. Thirty patients (51.7%) were defined as having high arterial stiffness and 28 patients (48.3%) were defined as having low arterial stiffness. The high arterial stiffness group had higher BMI ($P = 0.018$), SBP ($P = 0.018$), and pulse pressure ($P = 0.019$) than the low arterial stiffness group. Serum leptin ($P = 0.008$) and hemoglobin ($P = 0.040$) levels were also significantly higher in the high arterial stiffness group.

Comorbidity and the use of antihypertensives and anti-lipid drugs are presented in Table 2. There were higher percentages of participants with diabetes ($P = 0.019$) and hypertension ($P = 0.019$) in the high than the low arterial stiffness group. The effects of the use of β -blockers, calcium channel blockers (CCB), angiotensin-converting enzyme inhibitors (ACEi), angiotensin receptor blockers (ARB), statins and fibrates on blood pressure, atherosclerosis, and lipid burden were also recorded. No statistically significant differences were recorded between groups in subgroup analysis for sex, use of ACEi, ARB, β -blockers, CCB, statins, or fibrates.

Table 3 shows the correlation between cfPWV values and clinical variables in these 58 geriatric HD patients. Univariate linear regression analyses revealed positive correlations between pre-HD body weight ($r = 0.293$; $P = 0.026$), post-HD body weight ($r = 0.300$; $P = 0.022$), BMI ($r = 0.313$; $P = 0.017$), pulse pressure ($r = 0.275$; $P = 0.036$), hemoglobin level ($r = 0.270$; $P = 0.040$), and log-leptin level ($r = 0.422$; $P = 0.001$). Table 4 shows the multivariate, forward stepwise linear regression analysis of the variables significantly associated with cfPWV value (diabetes, hypertension, post-HD body weight, BMI and hemoglobin, and log-leptin). Log-leptin ($\beta = 0.408$, adjusted R^2 change = 0.164, $P = 0.001$) and diabetes ($\beta = 0.312$, adjusted R^2 change = 0.085, $P = 0.009$) were independent factors of the cfPWV value in geriatric HD patients.

Table 1: Clinical variables of the 58 geriatric hemodialysis patients with high or low central arterial stiffness

Characteristics	All patients (n=58)	Low central arterial stiffness group (n=30)	High central arterial stiffness group (n=28)	P
Age (years)	75.14±6.03	74.93±6.39	75.36±5.73	0.792
HD duration (years)	4.14 (1.88-8.47)	4.35 (1.76-10.44)	4.09 (2.04-6.41)	0.565
Pre-HD body weight (kg)	57.38±15.93	55.08±12.69	60.98±11.03	0.065
Post-HD body weight (kg)	56.04±12.00	53.24±12.46	59.03±10.92	0.066
Waist circumference (cm)	90.24±11.17	87.93±11.74	92.71±10.16	0.104
BMI (kg/m ²)	22.64±4.09	21.43±3.63	23.94±4.23	0.018*
cfPWV (m/s)	9.79±4.17	6.58±2.30	13.23±2.70	<0.001*
SBP (mmHg)	136.97±24.52	129.67±24.76	144.79±22.09	0.018*
DBP (mmHg)	69.66±13.78	68.07±12.95	71.36±14.67	0.368
Pulse pressure (mmHg)	67.31±19.35	61.60±18.06	73.43±19.11	0.019*
White blood count (×1000/uL)	5.77±1.76	5.75±1.96	5.79±1.57	0.934
Hemoglobin (g/dL)	10.34±1.02	10.02±1.23	10.68±0.76	0.013*
Iron (µg/dL)	74.24±26.64	78.83±28.83	69.32±23.60	0.176
Total iron-binding capacity (µg/dL)	252.93±35.83	257.33±40.30	248.21±30.35	0.337
Ferritin (ng/mL)	203.90 (102.58-422.70)	245.10 (56.45-494.23)	190.95 (123.18-320.03)	0.950
Total cholesterol (mg/dL)	140.64±28.48	135.97±28.03	145.64±28.62	0.199
Triglycerides (mg/dL)	131.21±77.92	121.37±63.33	141.75±91.04	0.324
Glucose (mg/dL)	132.50 (114.00-179.00)	128.50 (109.00-163.50)	141.50 (120.75-210.00)	0.104
Blood urea nitrogen (mg/dL)	57.38±15.93	56.43±16.22	58.39±15.85	0.644
Creatinine (mg/dL)	8.48±1.72	8.53±1.58	8.42±1.89	0.807
Total calcium (mg/dL)	8.93±0.76	8.81±0.71	9.06±0.79	0.200
Phosphorus (mg/dL)	4.39±1.24	4.29±1.29	4.50±1.21	0.519
Intact parathyroid hormone (pg/mL)	240.97±210.83	248.65±194.66	232.74±230.22	0.777
Leptin (ng/mL)	12.49 (4.13-52.58)	7.90 (2.60-26.68)	21.87 (6.24-74.22)	0.008*
Urea reduction rate	0.75±0.04	0.75±0.05	0.74±0.03	0.805
Kt/V (Gotch)	1.38±0.16	1.39±0.19	1.37±0.13	0.643

* $P < 0.05$ is considered statistically significant. Values for continuous variables are given as means±SD and were tested by Student's t -test; variables not normally distributed are given as medians and interquartile range and were tested by Mann-Whitney U-test. HD: Hemodialysis, SBP: Systolic blood pressure, DBP: Diastolic blood pressure, Kt/V: Fractional clearance index for urea, SD: Standard deviation, cfPWV: Carotid-femoral pulse wave velocity, BMI: Body mass index

Table 2: Distribution of hemodialysis patients with high and low central arterial stiffness in subgroup analysis

Characteristics	Low central arterial stiffness group (%)	High central arterial stiffness group (%)	P
Gender			
Male	14 (46.7)	13 (46.4)	0.986
Female	16 (53.3)	15 (53.6)	
Diabetes			
No	22 (73.3)	12 (42.9)	0.019*
Yes	8 (26.7)	16 (57.1)	
Hypertension			
No	21 (70.0)	11 (39.3)	0.019*
Yes	9 (30.0)	17 (60.7)	
Hepatitis B carrier			
No	27 (90.0)	27 (96.4)	0.334
Yes	3 (10.0)	1 (3.6)	
Hepatitis C carrier			
No	24 (80.0)	24 (85.7)	0.565
Yes	6 (20.0)	4 (14.3)	
ARB use			
No	24 (80.0)	17 (60.7)	0.107
Yes	6 (20.0)	11 (39.3)	
β-blocker use			
No	24 (80.0)	16 (57.1)	0.060
Yes	6 (20.0)	12 (42.9)	
CCB use			
No	19 (63.3)	15 (53.6)	0.451
Yes	11 (36.7)	13 (46.4)	
Statin use			
No	27 (90.0)	23 (82.1)	0.386
Yes	3 (10.0)	5 (17.9)	
Fibrate use			
No	28 (93.3)	24 (85.7)	0.341
Yes	2 (6.7)	4 (14.3)	

* $P < 0.05$ is considered statistically significant. Data are expressed as n (%) of patients and analysis was done using the Chi-square test. ARB: Angiotensin receptor blockers, CCB: Calcium channel blockers

Adjustment of the factors significantly associated with central arterial stiffness (adopted factors: diabetes, hypertension, waist circumference, BMI, pulse pressure, hemoglobin, and leptin) in multivariate logistic regression analysis revealed that increased serum leptin levels (odds ratio [OR]: 1.053; 95% confidence interval [CI]: 1.007–1.100; $P = 0.023$), hemoglobin level (OR: 2.869; 95% CI: 1.176–6.999; $P = 0.021$), and pulse pressure (OR: 1.078; 95% CI: 1.003–1.159; $P = 0.041$) were independent factors of central arterial stiffness among geriatric HD patients [Table 5].

DISCUSSION

This study reported that the serum leptin level was positively associated with central arterial stiffness in geriatric HD patients and was an independent factor of the cfPWV value in geriatric HD patients.

The aging process is associated with fat redistribution, including peripheral subcutaneous fat loss and central fat accumulation [12,18]. Older people are susceptible to adipokine dysregulation caused by visceral obesity [12,19]. Several mechanisms have been proposed to explain the correlation of

Table 3: Correlation of carotid-femoral pulse wave velocity levels and clinical variables in univariate linear regression analysis among the 58 geriatric hemodialysis patients

Variable	r	P
Age (years)	0.084	0.532
Log-HD duration (years)	0.024	0.859
Pre-HD body weight (kg)	0.293	0.026*
Post-HD body weight (kg)	0.300	0.022*
Waist circumference (cm)	0.202	0.128
BMI (kg/m ²)	0.313	0.017*
SBP (mmHg)	0.230	0.082
DBP (mmHg)	0.023	0.866
Pulse pressure (mmHg)	0.275	0.036*
White blood count ($\times 1000/uL$)	0.077	0.568
Hemoglobin (g/dL)	0.270	0.040*
Iron ($\mu g/dL$)	-0.181	0.173
Total iron-binding capacity ($\mu g/dL$)	-0.204	0.124
Total cholesterol (mg/dL)	0.024	0.857
Triglycerides (mg/dL)	0.126	0.345
Blood urea nitrogen (mg/dL)	-0.098	0.464
Creatinine (mg/dL)	0.098	0.467
Total calcium (mg/dL)	0.180	0.175
Phosphorus (mg/dL)	-0.045	0.739
Intact parathyroid hormone (pg/mL)	-0.131	0.327
Log-leptin (ng/mL)	0.422	0.001*
Urea reduction rate	-0.019	0.885
Kt/V (Gotch)	-0.030	0.823

* $P < 0.05$ was considered statistically significant. Data for ferritin, glucose, and leptin levels showed skewed distributions, and therefore were log-transformed before analysis. HD: Hemodialysis, SBP: Systolic blood pressure, DBP: Diastolic blood pressure, Kt/V: Fractional clearance index for urea, BMI: Body mass index

hyperleptinemia with arterial stiffness. One possible mechanism is that hyperleptinemia enhances sympathetic tone and decreases kidney natriuresis, causing hypertension. Another possible mechanism is that hyperleptinemia stimulates intimal monocyte recruitment, macrophage cell transformation, and vascular smooth muscle cell proliferation, causing atherosclerotic changes; induces endothelial nitric oxide imbalance-mediated endothelial dysfunction, and finally prompts induction of platelet aggregation-related thrombosis induction [17]. Many cross-sectional studies have also proved a relationship between arterial stiffness and hyperleptinemia among healthy individuals, obese, diabetic, hypertensive, and elderly patients, kidney transplantation recipients, and coronary artery disease patients [14,15,20]. The results of the present study also showed that hyperleptinemia was positively associated with central arterial stiffness and was an independent factor for the cfPWV value in geriatric HD patients.

Our results revealed a positive association between central arterial stiffness and diabetes in geriatric HD patients. Diabetes is a traditional risk factor for arterial stiffness in elderly diabetic people [21]. Diabetes accelerates arterial aging, resulting in stiffer arteries and a steeper increase in pulse pressure in elderly participants [22]. Hyperinsulinemia accentuates renin-angiotensin-aldosterone axis activity, leads to vascular wall hypertrophy and alters vessels elasticity. In addition, advanced glycation end products further induce endothelium injury. These effects promote

Table 4: Multivariate forward stepwise regression analysis of the factors correlated with carotid-femoral pulse wave velocity levels among 58 geriatric hemodialysis patients

Items	Beta	Adjusted R ²	Adjusted R ² change	P
Log-leptin (ng/mL)	0.408	0.164	0.164	0.001*
Diabetes mellitus	0.312	0.249	0.085	0.009*

* $P < 0.05$ was considered statistically significant. Analysis was done using multivariate logistic regression analysis (adopted factors: Diabetes mellitus, hypertension, SBP, pulse pressure, post-HD body weight, BMI, hemoglobin, and log-leptin). HD: Hemodialysis, SBP: Systolic blood pressure, BMI: Body mass index

Table 5: Multivariate logistic regression analysis of the factors correlated with central arterial stiffness among 58 geriatric hemodialysis patients

Variable	OR	95% CI	P
Leptin (ng/mL) (each increase of 1 ng/mL)	1.053	1.007-1.100	0.023*
Hemoglobin (g/dL) (each increase of 1 g/dL)	2.869	1.176-6.999	0.021*
Pulse pressure (mmHg) (each increase of 1 mmHg)	1.078	1.003-1.159	0.041*

* $P < 0.05$ was considered statistically significant. Analysis was done using multivariate logistic regression analysis (adopted factors: Diabetes, hypertension, waist circumference, BMI, pulse pressure, hemoglobin, and leptin). OR: Odds ratio, CI: Confidence interval, BMI: Body mass index

atherosclerosis [23]. Our study revealed that diabetes mellitus was positively correlated with cfPWV values. After multiple regression analysis, diabetes was also an independent factor for the cfPWV value in geriatric HD patients.

BMI and waist circumference are commonly used to assess general and abdominal adiposity, and the increase in central arterial stiffness in patients with metabolic syndrome is strongly related to the amount of visceral adipose tissue [24,25]. Our results also noted BMI and waist circumference were positively associated with the cfPWV value in geriatric HD patients. Gender differences have been reported in the traditional vascular risk factors for hypertension [25]. Mean arterial pressure contributed independently to variations in cfPWV values in community participants in South Africa [26]. The prevalence of systolic hypertension for those over 60 years old is twice that of those between 40 and 59 years [27]. The SBP is elevated in both age groups, whereas a normal or low DBP in the elderly is due to aortic stiffening and increased pulse pressure [27]. Our results noted higher prevalence of hypertension, elevated SBP, and elevated pulse pressure in the high arterial stiffness group and pulse pressure was positively associated with the cfPWV value. Moreover, this difference in pulse pressure between central arterial stiffness groups remained after multivariate adjustment. The cfPWV value was significantly higher in men with type 2 diabetes mellitus in one study [28]. However, the BMI in women was associated with a higher cfPWV value in 1517 participants in the Nijmegen Biomedical Study, and age 50–70 years and female sex were associated with higher cfPWV values in community participants in South Africa [25,26]. Our results did not find statistically significant gender differences in central arterial stiffness among geriatric HD patients. The reasons for the sex differences in hypertension are multifactorial, implying different roles of sex hormones, the renin–angiotensin system, sympathetic activity,

and arterial stiffness [29]. Further larger prospective studies are needed to establish these relationships.

The current study also revealed a correlation between hemoglobin levels and high central arterial stiffness. There were similar findings in a large Japanese study in a healthy population with BMI < 25 kg/m² [30]. Another study in which the brachial-ankle PWV was measured found a lower hemoglobin level was independently beneficial for peripheral arterial stiffness in a healthy community population, but only in women [31]. Possible mechanisms accounting for this correlation include activation of nuclear factor-kappa B transcription, decreased nitric oxide to mediate vascular homeostasis by hemoglobin, and plaque instability and atherosclerosis progression caused by hemoglobin membranes [32-34]. Another study proposed a conflicting opinion that only the density score, composed of hemoglobin and total protein, would alter PWV instead of hemoglobin alone [35]. This relationship between the hemoglobin level and the severity of arterial stiffness is not conclusive.

There are a few limitations in our study. First, since our study participants were elderly HD participants from a single center at East Taiwan, and further assessment is needed to determine whether the results are applicable to other populations. Second, the current data lack comprehensive assessment and in-depth nutritional evaluation, such as the subjective global assessment score, to further analyze the relationship between nutrition and leptin in the geriatric dialysis population. Third, the study had a small sample size and a cross-sectional design, which may affect the statistical significance. Only 58 geriatric HD participants were enrolled in this study, and the power to predict central arterial stiffness was only 0.73. Further long-term prospective studies are needed to confirm the relationship between central arterial stiffness and hyperleptinemia in geriatric HD patients.

CONCLUSION

A higher serum leptin level was associated with central arterial stiffness in geriatric HD patients. Moreover, diabetes was also positively associated with the cfPWV value in these geriatric HD patients.

Financial support and sponsorship

This study was financially supported by a grant from the Ministry of Science and Technology, Taiwan (MOST-104-2314-B-303-010).

Conflict of interest

There are no conflicts of interest.

REFERENCES

- Williams ME, Sandeep J, Catic A. Aging and ESRD demographics: Consequences for the practice of dialysis. *Semin Dial* 2012;25:617-22.
- Saran R, Robinson B, Abbott KC, Agodoa LY, Ayanian J, Bragg-Gresham J, et al. US Renal Data System 2016 Annual Data Report – Epidemiology of kidney disease in the United States chapter 1: Incidence, prevalence, patient characteristics, and treatment modalities. *Am J Kidney Dis* 2017;69(3 Suppl 1):S261-300.
- Singh P, Germain MJ, Cohen L, Unruh M. The elderly patient on dialysis: Geriatric considerations. *Nephrol Dial Transplant* 2014;29:990-6.

4. Laurent S, Cockcroft J, Van Bortel L, Boutouyrie P, Giannattasio C, Hayoz D, et al. Expert consensus document on arterial stiffness: Methodological issues and clinical applications. *Eur Heart J* 2006;27:2588-605.
5. Blacher J, Pannier B, Guerin AP, Marchais SJ, Safar ME, London GM, et al. Carotid arterial stiffness as a predictor of cardiovascular and all-cause mortality in end-stage renal disease. *Hypertension* 1998;32:570-4.
6. Zoungas S, Cameron JD, Kerr PG, Wolfe R, Muske C, McNeil JJ, et al. Association of carotid intima-medial thickness and indices of arterial stiffness with cardiovascular disease outcomes in CKD. *Am J Kidney Dis* 2007;50:622-30.
7. Luo L, Liu M. Adipose tissue in control of metabolism. *J Endocrinol* 2016;231:R77-99.
8. Nasrallah MP, Ziyadeh FN. Overview of the physiology and pathophysiology of leptin with special emphasis on its role in the kidney. *Semin Nephrol* 2013;33:54-65.
9. Oda A, Taniguchi T, Yokoyama M. Leptin stimulates rat aortic smooth muscle cell proliferation and migration. *Kobe J Med Sci* 2001;47:141-50.
10. Bouloumie A, Marumo T, Lafontan M, Busse R. Leptin induces oxidative stress in human endothelial cells. *FASEB J* 1999;13:1231-8.
11. Mishra S, Harris TB, Hsueh WC, Hue T, Leak TS, Li R, et al. The association of serum leptin with mortality in older adults. *PLoS One* 2015;10:e0140763.
12. Arai Y, Takayama M, Abe Y, Hirose N. Adipokines and aging. *J Atheroscler Thromb* 2011;18:545-50.
13. Lee MC, Chen YC, Ho GJ, Shih MH, Chou KC, Hsu BG, et al. Serum leptin levels positively correlate with peripheral arterial stiffness in kidney transplantation patients. *Transplant Proc* 2014;46:353-8.
14. Tsai JP, Lee MC, Chen YC, Ho GJ, Shih MH, Hsu BG, et al. Hyperleptinemia is a risk factor for the development of central arterial stiffness in kidney transplant patients. *Transplant Proc* 2015;47:1825-30.
15. Tsai JP, Wang JH, Chen ML, Yang CF, Chen YC, Hsu BG, et al. Association of serum leptin levels with central arterial stiffness in coronary artery disease patients. *BMC Cardiovasc Disord* 2016;16:80.
16. Chen MC, Hsu BG, Lee CJ, Wang JH. Hyperleptinemia positively correlates with cardiometabolic syndrome in hypertensive patients. *Int J Clin Exp Pathol* 2016;9:12959-67.
17. Mancia G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Böhm M, et al. 2013 ESH/ESC guidelines for the management of arterial hypertension: The task force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2013;31:1281-357.
18. Sabbatini AR, Fontana V, Laurent S, Moreno H. An update on the role of adipokines in arterial stiffness and hypertension. *J Hypertens* 2015;33:435-44.
19. Arai Y, Takayama M, Gondo Y, Inagaki H, Yamamura K, Nakazawa S, et al. Adipose endocrine function, insulin-like growth factor-1 axis, and exceptional survival beyond 100 years of age. *J Gerontol A Biol Sci Med Sci* 2008;63:1209-18.
20. Sweeney G. Cardiovascular effects of leptin. *Nat Rev Cardiol* 2010;7:22-9.
21. Teoh WL, Price JF, Williamson RM, Payne RA, Van Look LA, Reynolds RM, et al. Metabolic parameters associated with arterial stiffness in older adults with type 2 diabetes: The Edinburgh type 2 diabetes study. *J Hypertens* 2013;31:1010-7.
22. Stehouwer CD, Henry RM, Ferreira I. Arterial stiffness in diabetes and the metabolic syndrome: A pathway to cardiovascular disease. *Diabetologia* 2008;51:527-39.
23. Shirwany NA, Zou MH. Arterial stiffness: A brief review. *Acta Pharmacol Sin* 2010;31:1267-76.
24. Wohlfahrt P, Somers VK, Cifkova R, Filipovsky J, Seidlerova J, Kraljovicheva A, et al. Relationship between measures of central and general adiposity with aortic stiffness in the general population. *Atherosclerosis* 2014;235:625-31.
25. van den Munckhof IC, Holewijn S, de Graaf J, Rutten JH. Sex differences in fat distribution influence the association between BMI and arterial stiffness. *J Hypertens* 2017;35:1219-25.
26. Norman G, Norton GR, Gomes M, Michel F, Majane OH, Sareli P, et al. Circulating resistin concentrations are independently associated with aortic pulse wave velocity in a community sample. *J Hypertens* 2016;34:274-81.
27. Smulyan H, Mookherjee S, Safar ME. The two faces of hypertension: Role of aortic stiffness. *J Am Soc Hypertens* 2016;10:175-83.
28. Sakura T, Morioka T, Shioi A, Kakutani Y, Miki Y, Yamazaki Y, et al. Lipopolysaccharide-binding protein is associated with arterial stiffness in patients with type 2 diabetes: A cross-sectional study. *Cardiovasc Diabetol* 2017;16:62.
29. Caddeu C, Franconi F, Cassisa L, Campesi I, Pepe A, Cugusi L, et al. Arterial hypertension in the female world: Pathophysiology and therapy. *J Cardiovasc Med (Hagerstown)* 2016;17:229-36.
30. Shimizu Y, Nakazato M, Sekita T, Kadota K, Yamasaki H, Takamura N, et al. Association between hemoglobin levels and arterial stiffness for general Japanese population in relation to body mass index status: The Nagasaki Islands Study. *Geriatr Gerontol Int* 2014;14:811-8.
31. Kawamoto R, Tabara Y, Kohara K, Miki T, Kusunoki T, Katoh T, et al. A slightly low hemoglobin level is beneficially associated with arterial stiffness in Japanese community-dwelling women. *Clin Exp Hypertens* 2012;34:92-8.
32. Tziakas DN, Chalikias GK, Stakos D, Boudoulas H. The role of red blood cells in the progression and instability of atherosclerotic plaque. *Int J Cardiol* 2010;142:2-7.
33. Moreno PR. Neovascularization in human atherosclerosis. *Circulation* 2006;113:2245-52.
34. Kim-Shapiro DB, Schechter AN, Gladwin MT. Unraveling the reactions of nitric oxide, nitrite, and hemoglobin in physiology and therapeutics. *Arterioscler Thromb Vasc Biol* 2006;26:697-705.
35. Kim JY, Yoon J, Cho M, Lee BK, Karimi A, Shin S, et al. Blood characteristics effect on pulse wave velocity. *Clin Hemorheol Microcirc* 2013;55:193-203.