

**[ $\mu$ -*N,N'*-Bis(2-aminoethyl)ethane-1,2-diamine- $\kappa^4 N^1, N^1': N^2, N^2'$ ]bis{[*N,N'*-bis(2-aminoethyl)ethane-1,2-diamine- $\kappa^4 N, N', N'', N'''$ ]cadmium} tetrakis-(perchlorate)**

Hamid Goudarziafshar,<sup>a\*</sup> Yunes Abbasityula,<sup>a</sup> Václav Eigner<sup>b,c</sup> and Michal Dušek<sup>c</sup>

<sup>a</sup>Faculty of Science, Department of Chemistry, Ilam University, Ilam, Iran, <sup>b</sup>Department of Solid State Chemistry, Institute of Chemical Technology, Technická 5, 166 28 Prague, Czech Republic, and <sup>c</sup>Institute of Physics AS CR, v.v.i., Na Slovance 2, 182 21 Prague 8, Czech Republic  
Correspondence e-mail: hamid\_gafshar@yahoo.com


Received 21 July 2012; accepted 28 July 2012

Key indicators: single-crystal X-ray study;  $T = 120$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in solvent or counterion;  $R$  factor = 0.030;  $wR$  factor = 0.106; data-to-parameter ratio = 18.4.

The centrosymmetric dinuclear cadmium title complex,  $[\text{Cd}_2(\text{C}_6\text{H}_{18}\text{N}_4)_3](\text{ClO}_4)_4$ , was obtained by the reaction of *N,N'*-bis(2-aminoethyl)ethane-1,2-diamine (trien) with  $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$  and sodium perchlorate in methanol. The  $\text{Cd}^{\text{II}}$  cation is coordinated by four N atoms of a non-bridging trien ligand and by two N atoms of a bridging trien ligand in a slightly distorted octahedral coordination geometry. The bridging ligand shares another two N atoms with a neighboring symmetry-equivalent  $\text{Cd}^{\text{II}}$  cation. The structure displays  $\text{C}-\text{H} \cdots \text{O}$  and  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonding. The perchlorate anion is disordered over two sets of sites in a 0.854 (7):0.146 (7) ratio.

## Related literature

Polyamines are an important class of *N*-donor ligands, particularly for transition metals, see: Patel *et al.* (2007); Blackman (2005). For polynuclear complexes, see: Gustafsson *et al.* (2010); Ambrosi *et al.* (2009); You *et al.* (2011). For polynuclear complexes of cadmium, see: Evans & Lin (2002); For background to the use of the trien ligand in complexation, see: Cai *et al.* (2001*a,b*); Buckingham *et al.* (1974, 1975); Chowdhury *et al.* (2007). Buckingham & Jones (1965); Shinohara *et al.* (1991); He (2009); Patel *et al.* (2008); Anderson *et al.* (1977); Shoukry *et al.* (1998); Hu *et al.* (2000). For dinuclear Cd complexes, see: Das *et al.* (2010); Nie *et al.* (2010); Wang *et al.* (2011); Sun *et al.* (2010). For details of the preparation, see: Harrowfield *et al.* (1996).



## Experimental

### Crystal data

$[\text{Cd}_2(\text{C}_6\text{H}_{18}\text{N}_4)_3](\text{ClO}_4)_4$   
 $M_r = 1061.3$   
Monoclinic,  $P2_1/n$   
 $a = 8.8056$  (2) Å  
 $b = 15.0259$  (3) Å  
 $c = 14.7516$  (3) Å  
 $\beta = 95.4420$  (17)°

$V = 1943.02$  (7) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 1.45$  mm<sup>-1</sup>  
 $T = 120$  K  
 $0.70 \times 0.51 \times 0.33$  mm

### Data collection

Agilent Xcalibur Atlas Gemini ultra diffractometer  
Absorption correction: analytical (*CrysAlis PRO*; Agilent, 2012)  
 $T_{\text{min}} = 0.509$ ,  $T_{\text{max}} = 0.738$

31295 measured reflections  
4952 independent reflections  
4373 reflections with  $I > 3\sigma(I)$   
 $R_{\text{int}} = 0.022$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.106$   
 $S = 2.02$   
4952 reflections  
269 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 1.13$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.92$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Cd1—N1	2.362 (2)	Cd1—N10	2.380 (3)
Cd1—N4	2.390 (3)	Cd1—N11	2.374 (3)
Cd1—N7	2.377 (3)	Cd1—N14	2.375 (2)

**Table 2**

Hydrogen-bond geometry (Å, °).

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C5—H1c5 $\cdots$ O4c <sup>i</sup>	0.96	2.49	3.449 (3)	172
N1—H2n1 $\cdots$ O3c <sup>ii</sup>	0.80 (4)	2.46 (4)	3.149 (4)	145 (3)
N11—H1n11 $\cdots$ O2c <sup>iii</sup>	0.82 (4)	2.40 (4)	3.140 (4)	149 (3)
N11—H2n11 $\cdots$ O4a <sup>ii</sup>	0.92 (4)	2.44 (4)	3.210 (5)	141 (3)
N4—H1n4 $\cdots$ O4a <sup>ii</sup>	0.98 (3)	2.36 (3)	3.271 (4)	154 (2)
N4—H1n4 $\cdots$ O5a <sup>ii</sup>	0.98 (3)	2.38 (3)	3.236 (4)	145 (2)
N4—H1n4 $\cdots$ O5b <sup>ii</sup>	0.98 (3)	2.22 (3)	3.113 (13)	151 (3)

Symmetry codes: (i)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + 1, -y + 1, -z$ ; (iii)  $-x, -y + 1, -z$ .

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis, 2007); program(s) used to refine structure: *JANA2006* (Petříček *et al.*, 2006); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *publCIF* (Westrip, 2010).

This work was supported by the University of Ilam, the Institutional research plan No. AVOZ10100521 of the Institute of Physics and the Praemium Academiae project of the Academy of Sciences (ASCR).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RU2040).

## References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Ambrosi, G., Formica, M., Fusi, V., Giorgi, L., Macedi, E., Micheloni, M., Paoli, P. & Rossi, P. (2009). *Inorg. Chem.* **48**, 10424–10434.
- Anderson, B. F., Bell, J. D., Buckingham, D. A., Cresswell, P. J., Gainsford, G. J., Marzilli, L. G., Robertson, G. B. & Sargeson, A. M. (1977). *Inorg. Chem.* **16**, 3233–3244.
- Blackman, A. G. (2005). *Polyhedron*, **24**, 1–39.
- Buckingham, D. A., Cresswell, P. J., Dellaca, R. J., Dwyer, M., Gainford, G. J., Marzilli, L. G., Maxwell, I. E., Robinson, W. T., Sargeson, A. M. & Turnbull, K. R. (1974). *J. Am. Chem. Soc.* **96**, 1713–1725.
- Buckingham, D. A., Dwyer, M., Gainsford, G. J., Ho, V. J., Marzilli, L. G., Robinson, W. T., Sargeson, A. M. & Turnbull, K. B. (1975). *Inorg. Chem.* **14**, 1739–1752.
- Buckingham, D. A. & Jones, D. (1965). *Inorg. Chem.* **4**, 1387–1392.
- Cai, J., Hu, X., Feng, X., Ji, L. & Bernal, I. (2001a). *Acta Cryst.* **B57**, 45–53.
- Cai, J., Hu, X., Yao, J. & Ji, L. (2001b). *Inorg. Chem. Commun.* **4**, 478–482.
- Chowdhury, H., Ghosh, R., Rahaman, Sk. H. & Ghosh, B. K., (2007). *Polyhedron*, **26**, 5023–5029.
- Das, S., Bhar, K., Fun, H. K., Chantrapromma, S. & Ghosh, B. K. (2010). *Inorg. Chim. Acta*, **363**, 784–792.
- Evans, O. R. & Lin, W. (2002). *Acc. Chem. Res.* **35**, 511–522.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gustafsson, M., Fischer, A., Ilyukhin, A., Maliarik, M. & Nordblad, P. (2010). *Inorg. Chem.* **49**, 5359–5361.
- Harrowfield, J. M., Miyamae, H., Skelton, B. W., Soudi, A. A. & White, A. H. (1996). *Aust. J. Chem.* **49**, 1165–1169.
- He, X. (2009). *Acta Cryst.* **E65**, m205.
- Hu, X., Cai, J., Feng, X. & Ji, L. (2000). *J. Chem. Crystallogr.* **30**, 27–34.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nie, F. M., Li, M., Li, G. X., Li, Z. & Li, W. (2010). *Inorg. Chim. Acta*, **363**, 3351–3358.
- Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
- Patel, R. N., Kesharwani, M. K., Singh, A., Patel, D. K. & Choudhary, M. (2008). *Transition Met. Chem.* **33**, 733–738.
- Patel, R. N., Singh, N., Patel, D. K. & Gundla, V. L. N. (2007). *Indian J. Chem. Sect. A*, **46**, 422–427.
- Petříček, V., Dušek, M. & Palatinus, L. (2006). *JANA2006*. Institute of Physics, Praha, Czech Republic.
- Shinohara, N., Matsufuji, S. & Okubo, W. (1991). *Polyhedron*, **10**, 107–112.
- Shoukry, M. M., Shehata, M. R. & Mohamed, M. M. A. (1998). *Mikrochim. Acta*, **129**, 107–113.
- Sun, S.-J., Deng, J.-H. & Liu, T.-L. (2010). *Acta Cryst.* **E66**, m981–m982.
- Wang, J., Xu, X.-J. & Tao, J.-Q. (2011). *Acta Cryst.* **C67**, m137–m139.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- You, Z. L., Wang, X. L., Zhang, J. C., Wang, C. & Zhou, X. S. (2011). *Struct. Chem.* **22**, 1297–1302.

## supplementary materials

*Acta Cryst.* (2012). E68, m1153–m1154 [doi:10.1107/S1600536812033880]

**[ $\mu$ -*N,N'*-Bis(2-aminoethyl)ethane-1,2-diamine- $\kappa^4 N^1, N^1': N^2, N^2'$ ]bis{[*N,N'*-bis(2-aminoethyl)ethane-1,2-diamine- $\kappa^4 N, N', N'', N'''$ ]cadmium} tetrakis(perchlorate)**

**Hamid Goudarziafshar, Yunes Abbasityula, Václav Eigner and Michal Dušek**

### Comment

Polyamines are an important class of N donor ligands, particularly for the transition metals (Patel *et al.*, 2007, 2008; Blackman, 2005). Considerable attention has been focused on the polynuclear complexes containing bridging ligands because of their interesting molecular topologies, as well as the fact that they may be designed with specific functionalities (Gustafsson *et al.* 2010;. Ambrosi *et al.* 2009; You *et al.* 2011). The investigation of polynuclear complexes of cadmium(II) is an important objective in view of their electronic and optoelectronic properties (Evans *et al.* 2002; Chowdhury *et al.* 2007).

The molecular structure of the title complex is shown in Fig. 1. The cadmium(II) centers are six-coordinate by four nitrogen atoms of the non-bridging tetradentate (trien) ligand and two nitrogen atoms of the bridging trien ligand, with a substantial departure from an ideal octahedral geometry [*cisoid* angles: 73.97 (8)–114.67 (9)°; *transoid* angles: 141.95 (1)–159.20 (6)°] (Table 1). The distance between the two cadmium(II) centers of the dinuclear complex is 7.735 Å, which is longer than the corresponding distance in dinickel(II) complex (7.497 Å) of the same ligand (Cai *et al.* 2001*b*) due to larger radius of cadmium. Cadmium atoms in the dinuclear complex are related by a 2 fold symmetry operation. Bond distance of Cd—N(trien) are in the range of 2.62 (3)- 2.90 (3) Å (Table 1). The structure exhibits disorder of one of the perchlorate anions in two positions with refined occupancy 0.854 (7) and 0.146 (7) for the major and minor component, respectively. The disorder was described using the rigid body approach. In the title complex the C—H...O and N—H...O hydrogen bonds have been found between the amine nitrogen/carbon donors and perchlorate acceptors (Fig.2),(Table 2).

### Experimental


*N,N'*-bis(2-aminoethyl)ethane-1,2-diamine (0.45 g, 3 mmol) was placed in one arm of a branched tube (Harrowfield *et al.*, 1996) and a mixture of Cd(NO<sub>3</sub>)<sub>2</sub>·4H<sub>2</sub>O (0.616 g, 2 mmol) and sodium perchlorate (0.488 g, 4 mmol) in the other. Methanol was then carefully added to fill both arms, the tube sealed and the ligand-containing arm immersed in a bath at 333 K, while the other was left at ambient temperature. After one week, colorless crystals were collected in the cooler arm. Then they were filtered off, washed with acetone and diethylether, and air dried. Yield: (53%).

### Refinement


All hydrogen atoms were discernible in difference Fourier maps and could be refined to reasonable geometry. According to common practice H atoms bonded to C were kept in ideal positions with C—H = 0.96 Å while positions of other H atoms were refined freely. In both cases  $U_{iso}(H)$  was set to  $1.2U_{eq}(C,N)$ . Disorder of perchlorate anion was refined using rigid body refinement, with occupancy ratio 0.85:0.15.

**Computing details**

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO* (Agilent, 2012); data reduction: *CrysAlis PRO* (Agilent, 2012); program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis, 2007); program(s) used to refine structure: *JANA2006* (Petricek *et al.*, 2006); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *publCIF* (Westrip, 2010).

**Figure 1**

The molecular structure of the title compound with 50% probability displacement ellipsoids.

**Figure 2**

Unit-cell packing diagram of the title compound viewed along the crystallographic  $\langle \rangle a \langle \rangle$  axis. Hydrogen bonds are indicated by dashed lines; (orange = cadmium; green = chlorine; violet = nitrogen; grey = carbon; light-grey = hydrogen).

$[\mu$ -*N,N'*-Bis(2-aminoethyl)ethane-1,2-diamine- $\kappa^4N^1,N^1':N^2,N^2'$ ]bis[[*N,N'*-bis(2-aminoethyl)ethane-1,2-diamine- $\kappa^4N,N',N'',N'''$ ]cadmium} tetrakis(perchlorate)

Crystal data

$[\text{Cd}_2(\text{C}_6\text{H}_{18}\text{N}_4)_3](\text{ClO}_4)_4$

$M_r = 1061.3$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 8.8056$  (2) Å

$b = 15.0259$  (3) Å

$c = 14.7516$  (3) Å

$\beta = 95.4420$  (17)°

$V = 1943.02$  (7) Å<sup>3</sup>

$Z = 2$

$F(000) = 1076$

$D_x = 1.814$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.7107$  Å

Cell parameters from 17686 reflections

$\theta = 2.9$ – $29.3$ °

$\mu = 1.45$  mm<sup>-1</sup>

$T = 120$  K

Prism, colourless

$0.70 \times 0.51 \times 0.33$  mm

Data collection

Agilent Xcalibur Atlas Gemini ultra diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: 10.3784 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: analytical

(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.509$ ,  $T_{\max} = 0.738$

31295 measured reflections

4952 independent reflections

4373 reflections with  $I > 3\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\max} = 29.4$ °,  $\theta_{\min} = 2.9$ °

$h = -11$ → $12$

$k = -19$ → $20$

$l = -20$ → $19$

Refinement

Refinement on  $F^2$

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.106$

$S = 2.02$

4952 reflections

269 parameters

0 restraints

82 constraints

H atoms treated by a mixture of independent and constrained refinement

Weighting scheme based on measured s.u.'s  $w =$

$1/(\sigma^2(I) + 0.0016I^2)$

$(\Delta/\sigma)_{\max} = 0.045$

$\Delta\rho_{\max} = 1.13$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.92$  e Å<sup>-3</sup>

Special details

**Experimental.** Absorption correction: analytical: *CrysAlisPro*, Agilent Technologies, Version 1.171.35.19 Analytical numeric absorption correction based on crystal shape

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cd1	0.286722 (18)	0.787742 (11)	0.064763 (11)	0.01848 (7)	
N1	0.5386 (3)	0.73266 (16)	0.08829 (18)	0.0272 (7)	
C2	0.5399 (3)	0.63634 (18)	0.0671 (2)	0.0292 (8)	
C3	0.4054 (3)	0.59042 (17)	0.10338 (19)	0.0283 (8)	
N4	0.2615 (2)	0.62928 (15)	0.06265 (14)	0.0228 (6)	
C5	0.1244 (3)	0.60375 (18)	0.10733 (18)	0.0300 (8)	
C6	-0.0058 (3)	0.6674 (2)	0.08095 (19)	0.0319 (9)	
N7	0.0352 (3)	0.76022 (18)	0.10385 (16)	0.0277 (7)	
C8	0.0284 (3)	0.78350 (18)	0.1997 (2)	0.0330 (9)	
C9	0.1157 (3)	0.8669 (2)	0.22447 (19)	0.0354 (9)	

N10	0.2773 (3)	0.85403 (16)	0.21066 (15)	0.0286 (7)	
N11	0.2454 (3)	0.77849 (15)	-0.09631 (18)	0.0296 (7)	
C12	0.2788 (3)	0.86355 (17)	-0.14108 (17)	0.0274 (8)	
C13	0.2345 (3)	0.94099 (17)	-0.08204 (17)	0.0231 (7)	
N14	0.3148 (2)	0.93478 (13)	0.00986 (14)	0.0183 (6)	
C15	0.4798 (3)	0.95387 (15)	0.01611 (16)	0.0200 (7)	
H1c2	0.632977	0.610336	0.094187	0.035*	
H2c2	0.535246	0.628241	0.002352	0.035*	
H1c3	0.407828	0.528128	0.089118	0.034*	
H2c3	0.411784	0.596978	0.168367	0.034*	
H1c5	0.147832	0.604608	0.172219	0.036*	
H2c5	0.094348	0.544411	0.089398	0.036*	
H1c6	-0.035184	0.662706	0.016775	0.0383*	
H2c6	-0.093123	0.650524	0.111322	0.0383*	
H1c8	0.068971	0.735436	0.2374	0.0396*	
H2c8	-0.076102	0.791229	0.211536	0.0396*	
H1c9	0.106623	0.881033	0.287172	0.0424*	
H2c9	0.074914	0.914972	0.18688	0.0424*	
H1c12	0.221103	0.866719	-0.199579	0.0329*	
H2c12	0.385736	0.866777	-0.14855	0.0329*	
H1c13	0.260242	0.996164	-0.109623	0.0277*	
H2c13	0.126385	0.940048	-0.07791	0.0277*	
H1c15	0.530028	0.909867	-0.017582	0.024*	
H2c15	0.523023	0.945412	0.077724	0.024*	
H1n10	0.331 (4)	0.819 (2)	0.267 (2)	0.0344*	
H2n10	0.340 (4)	0.909 (2)	0.227 (2)	0.0344*	
H1n1	0.574 (4)	0.746 (2)	0.148 (2)	0.0326*	
H1n7	-0.025 (5)	0.793 (2)	0.079 (3)	0.0332*	
H1n14	0.278 (3)	0.964 (2)	0.037 (2)	0.0219*	
H2n1	0.604 (4)	0.753 (3)	0.061 (2)	0.0326*	
H1n11	0.153 (5)	0.768 (2)	-0.104 (3)	0.0355*	
H2n11	0.285 (4)	0.736 (3)	-0.133 (3)	0.0355*	
H1n4	0.241 (3)	0.610 (2)	-0.001 (2)	0.0274*	
Cl1c	0.20589 (7)	0.15386 (4)	0.11243 (4)	0.03020 (19)	
O2c	0.0975 (3)	0.18921 (16)	0.16787 (16)	0.0431 (7)	
O3c	0.3065 (3)	0.22454 (16)	0.0906 (2)	0.0500 (9)	
O4c	0.2906 (3)	0.08460 (15)	0.16043 (15)	0.0451 (8)	
O5c	0.1347 (3)	0.11760 (15)	0.03083 (15)	0.0456 (7)	
Cl1a	0.8393 (2)	0.42448 (12)	0.19271 (14)	0.0246 (3)	0.854 (7)
O2a	0.8867 (6)	0.5086 (2)	0.2242 (3)	0.0768 (18)	0.854 (7)
O3a	0.8423 (5)	0.3641 (3)	0.2674 (3)	0.0378 (8)	0.854 (7)
O4a	0.6891 (4)	0.4247 (3)	0.1474 (3)	0.0468 (11)	0.854 (7)
O5a	0.9442 (5)	0.3933 (3)	0.1297 (3)	0.0388 (9)	0.854 (7)
Cl1b	0.8447 (14)	0.4319 (9)	0.2081 (8)	0.0246 (3)	0.146 (7)
O2b	0.9407 (15)	0.4949 (9)	0.2534 (9)	0.0768 (18)	0.146 (7)
O3b	0.8281 (14)	0.3574 (9)	0.2662 (9)	0.0378 (8)	0.146 (7)
O4b	0.6967 (14)	0.4661 (9)	0.1803 (9)	0.0468 (11)	0.146 (7)
O5b	0.9136 (14)	0.4022 (9)	0.1280 (9)	0.0388 (9)	0.146 (7)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cd1	0.01761 (13)	0.01696 (13)	0.02150 (13)	-0.00220 (5)	0.00519 (8)	0.00130 (5)
N1	0.0215 (11)	0.0231 (11)	0.0368 (13)	-0.0034 (9)	0.0021 (10)	-0.0032 (10)
C2	0.0242 (13)	0.0229 (12)	0.0407 (15)	0.0027 (10)	0.0053 (11)	-0.0014 (11)
C3	0.0302 (14)	0.0206 (12)	0.0338 (14)	-0.0002 (10)	0.0015 (11)	0.0043 (10)
N4	0.0252 (10)	0.0207 (11)	0.0227 (11)	-0.0071 (8)	0.0028 (8)	0.0030 (8)
C5	0.0324 (14)	0.0275 (13)	0.0306 (14)	-0.0112 (11)	0.0050 (11)	0.0044 (11)
C6	0.0200 (13)	0.0419 (17)	0.0339 (14)	-0.0112 (11)	0.0023 (10)	0.0042 (12)
N7	0.0187 (11)	0.0357 (13)	0.0289 (12)	0.0004 (9)	0.0041 (9)	0.0065 (10)
C8	0.0249 (14)	0.0429 (17)	0.0329 (15)	0.0045 (11)	0.0116 (12)	0.0036 (11)
C9	0.0357 (15)	0.0384 (16)	0.0336 (15)	0.0088 (12)	0.0118 (12)	-0.0014 (12)
N10	0.0310 (12)	0.0322 (12)	0.0231 (11)	0.0017 (10)	0.0047 (9)	-0.0011 (9)
N11	0.0434 (15)	0.0194 (11)	0.0265 (12)	-0.0092 (10)	0.0066 (11)	-0.0021 (9)
C12	0.0395 (15)	0.0222 (12)	0.0200 (11)	-0.0088 (11)	0.0001 (10)	0.0006 (10)
C13	0.0214 (11)	0.0208 (12)	0.0263 (12)	-0.0018 (9)	-0.0014 (9)	0.0026 (9)
N14	0.0182 (10)	0.0148 (9)	0.0224 (10)	-0.0017 (7)	0.0045 (8)	-0.0016 (7)
C15	0.0189 (11)	0.0173 (11)	0.0234 (12)	-0.0026 (9)	-0.0001 (9)	0.0023 (9)
Cl1c	0.0298 (3)	0.0308 (3)	0.0297 (3)	0.0041 (2)	0.0011 (3)	-0.0042 (2)
O2c	0.0390 (12)	0.0416 (11)	0.0498 (14)	0.0074 (10)	0.0101 (10)	-0.0168 (11)
O3c	0.0589 (17)	0.0402 (13)	0.0531 (14)	-0.0106 (10)	0.0159 (12)	-0.0020 (11)
O4c	0.0614 (15)	0.0400 (13)	0.0329 (11)	0.0157 (10)	0.0003 (10)	-0.0007 (9)
O5c	0.0517 (14)	0.0461 (13)	0.0361 (11)	0.0195 (10)	-0.0106 (10)	-0.0111 (10)
Cl1a	0.0346 (4)	0.0174 (5)	0.0224 (8)	-0.0034 (3)	0.0062 (4)	-0.0039 (4)
O2a	0.140 (4)	0.0407 (17)	0.060 (3)	-0.055 (2)	0.064 (3)	-0.0326 (18)
O3a	0.0459 (16)	0.0401 (14)	0.0267 (10)	-0.0048 (11)	0.0004 (10)	0.0130 (10)
O4a	0.0390 (14)	0.066 (2)	0.0348 (16)	0.0216 (14)	0.0008 (11)	0.0038 (15)
O5a	0.0396 (18)	0.0376 (16)	0.0420 (13)	-0.0078 (15)	0.0184 (13)	-0.0130 (11)
Cl1b	0.0310 (4)	0.0229 (5)	0.0196 (8)	-0.0082 (3)	0.0014 (4)	-0.0039 (4)
O2b	0.104 (4)	0.098 (2)	0.031 (3)	-0.084 (2)	0.023 (2)	-0.0276 (19)
O3b	0.0424 (15)	0.0356 (14)	0.0350 (11)	-0.0036 (11)	0.0020 (10)	0.0163 (10)
O4b	0.0542 (15)	0.049 (2)	0.0374 (17)	0.0258 (13)	0.0073 (12)	0.0057 (15)
O5b	0.0368 (18)	0.0497 (16)	0.0314 (14)	-0.0109 (14)	0.0112 (13)	-0.0133 (11)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Cd1—N1	2.362 (2)	C9—H1c9	0.96
Cd1—N4	2.390 (3)	C9—H2c9	0.96
Cd1—N7	2.377 (3)	N11—C12	1.481 (4)
Cd1—N10	2.380 (3)	N11—H1n11	0.82 (4)
Cd1—N11	2.374 (3)	N11—H2n11	0.92 (4)
Cd1—N14	2.375 (2)	C12—C13	1.526 (4)
N1—C2	1.481 (4)	C12—H1c12	0.96
N1—H1n1	0.93 (3)	C12—H2c12	0.96
N1—H2n1	0.80 (4)	C13—N14	1.471 (3)
C2—C3	1.512 (4)	C13—H1c13	0.96
C2—H1c2	0.96	C13—H2c13	0.96
C2—H2c2	0.96	N14—C15	1.475 (3)
C3—N4	1.471 (3)	N14—H1n14	0.70 (3)

C3—H1c3	0.96	C15—C15 <sup>i</sup>	1.519 (3)
C3—H2c3	0.96	C15—H1c15	0.96
N4—C5	1.480 (4)	C15—H2c15	0.96
C5—C6	1.515 (4)	Cl1c—O2c	1.418 (3)
C5—H1c5	0.96	Cl1c—O3c	1.439 (3)
C5—H2c5	0.96	Cl1c—O4c	1.428 (2)
C6—N7	1.472 (4)	Cl1c—O5c	1.413 (2)
C6—H1c6	0.96	Cl1a—O2a	1.397 (4)
C6—H2c6	0.96	Cl1a—O3a	1.425 (5)
N7—C8	1.463 (4)	Cl1a—O4a	1.424 (4)
N7—H1n7	0.78 (4)	Cl1a—O5a	1.449 (5)
C8—C9	1.497 (4)	Cl1b—O2b	1.397 (18)
C8—H1c8	0.96	Cl1b—O3b	1.425 (19)
C8—H2c8	0.96	Cl1b—O4b	1.424 (17)
C9—N10	1.469 (4)	Cl1b—O5b	1.449 (19)
N11—Cd1—N14	73.97 (8)	C8—C9—H2c9	109.47
N4—Cd1—N10	114.67 (9)	N10—C9—H1c9	109.47
N1—Cd1—N7	141.90 (9)	N10—C9—H2c9	109.47
N4—Cd1—N14	159.20 (6)	H1c9—C9—H2c9	109.17
Cd1—N1—C2	109.72 (16)	C12—N11—H1n11	110 (2)
Cd1—N1—H1n1	107 (2)	C12—N11—H2n11	103 (2)
Cd1—N1—H2n1	120 (3)	H1n11—N11—H2n11	102 (3)
C2—N1—H1n1	114 (2)	N11—C12—C13	109.4 (2)
C2—N1—H2n1	105 (3)	N11—C12—H1c12	109.47
H1n1—N1—H2n1	102 (3)	N11—C12—H2c12	109.47
N1—C2—C3	110.5 (2)	C13—C12—H1c12	109.47
N1—C2—H1c2	109.47	C13—C12—H2c12	109.47
N1—C2—H2c2	109.47	H1c12—C12—H2c12	109.59
C3—C2—H1c2	109.47	C12—C13—N14	110.6 (2)
C3—C2—H2c2	109.47	C12—C13—H1c13	109.47
H1c2—C2—H2c2	108.47	C12—C13—H2c13	109.47
C2—C3—N4	110.3 (2)	N14—C13—H1c13	109.47
C2—C3—H1c3	109.47	N14—C13—H2c13	109.47
C2—C3—H2c3	109.47	H1c13—C13—H2c13	108.36
N4—C3—H1c3	109.47	C13—N14—C15	115.44 (19)
N4—C3—H2c3	109.47	C13—N14—H1n14	106 (3)
H1c3—C3—H2c3	108.58	C15—N14—H1n14	111 (2)
C3—N4—C5	115.0 (2)	N14—C15—C15 <sup>i</sup>	114.62 (18)
N4—C5—C6	110.6 (2)	N14—C15—H1c15	109.47
N4—C5—H1c5	109.47	N14—C15—H2c15	109.47
N4—C5—H2c5	109.47	C15 <sup>i</sup> —C15—H1c15	109.47
C6—C5—H1c5	109.47	C15 <sup>i</sup> —C15—H2c15	109.47
C6—C5—H2c5	109.47	H1c15—C15—H2c15	103.78
H1c5—C5—H2c5	108.35	O2c—Cl1c—O3c	108.37 (16)
C5—C6—N7	112.1 (2)	O2c—Cl1c—O4c	109.59 (14)
C5—C6—H1c6	109.47	O2c—Cl1c—O5c	111.57 (14)
C5—C6—H2c6	109.47	O3c—Cl1c—O4c	110.20 (15)
N7—C6—H1c6	109.47	O3c—Cl1c—O5c	109.05 (16)



N7—C6—H2c6	109.47	O4c—C11c—O5c	108.05 (13)
H1c6—C6—H2c6	106.74	O2a—C11a—O3a	109.7 (3)
C6—N7—C8	114.6 (2)	O2a—C11a—O4a	112.9 (3)
C6—N7—H1n7	110 (2)	O2a—C11a—O5a	108.5 (3)
C8—N7—H1n7	102 (3)	O3a—C11a—O4a	108.2 (3)
N7—C8—C9	111.7 (2)	O3a—C11a—O5a	108.9 (3)
N7—C8—H1c8	109.47	O4a—C11a—O5a	108.6 (3)
N7—C8—H2c8	109.47	O2b—C11b—O3b	109.7 (11)
C9—C8—H1c8	109.47	O2b—C11b—O4b	112.9 (11)
C9—C8—H2c8	109.47	O2b—C11b—O5b	108.5 (11)
H1c8—C8—H2c8	107.19	O3b—C11b—O4b	108.2 (11)
C8—C9—N10	109.8 (2)	O3b—C11b—O5b	108.9 (11)
C8—C9—H1c9	109.47	O4b—C11b—O5b	108.6 (11)

Symmetry code: (i)  $-x+1, -y+2, -z$ .

*Hydrogen-bond geometry* ( $\text{\AA}, ^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C5—H1c5 $\cdots$ O4c <sup>ii</sup>	0.96	2.49	3.449 (3)	172
C8—H2c8 $\cdots$ O3b <sup>ii</sup>	0.96	2.48	3.412 (13)	163
N10—H1n10 $\cdots$ O2c <sup>ii</sup>	1.06 (3)	2.24 (4)	3.191 (3)	149 (3)
N10—H2n10 $\cdots$ O2b <sup>iii</sup>	1.01 (3)	2.32 (3)	3.268 (13)	156 (3)
N1—H1n1 $\cdots$ O3a <sup>iii</sup>	0.93 (3)	2.25 (3)	3.018 (5)	139 (3)
N1—H1n1 $\cdots$ O3b <sup>iii</sup>	0.93 (3)	2.22 (4)	3.004 (13)	141 (3)
N7—H1n7 $\cdots$ O5c <sup>iv</sup>	0.78 (4)	2.25 (4)	2.999 (3)	159 (4)
N1—H2n1 $\cdots$ O3c <sup>v</sup>	0.80 (4)	2.46 (4)	3.149 (4)	145 (3)
N11—H1n11 $\cdots$ O2c <sup>iv</sup>	0.82 (4)	2.40 (4)	3.140 (4)	149 (3)
N11—H2n11 $\cdots$ O4a <sup>v</sup>	0.92 (4)	2.44 (4)	3.210 (5)	141 (3)
N4—H1n4 $\cdots$ O4a <sup>v</sup>	0.98 (3)	2.36 (3)	3.271 (4)	154 (2)
N4—H1n4 $\cdots$ O5a <sup>v</sup>	0.98 (3)	2.38 (3)	3.236 (4)	145 (2)
N4—H1n4 $\cdots$ O5b <sup>v</sup>	0.98 (3)	2.22 (3)	3.113 (13)	151 (3)

Symmetry codes: (ii)  $-x+1/2, y+1/2, -z+1/2$ ; (iii)  $-x+3/2, y+1/2, -z+1/2$ ; (iv)  $-x, -y+1, -z$ ; (v)  $-x+1, -y+1, -z$ .