

COMMENTARY

Open Access

# Cilia, *tubby* mice, and obesity

Saikat Mukhopadhyay<sup>1,2\*</sup> and Peter K Jackson<sup>1</sup>

Please see related Research article by Li et al., <http://www.ciliajournal.com/content/1/1/21>

## Abstract

Primary cilia have been previously linked to the central regulation of satiety. The *tubby* mouse is characterized by maturity-onset obesity and blindness. A recent paper demonstrates molecular defects in trafficking of ciliary GPCRs in the central neurons of *tubby* mice, underscoring the role of ciliary signaling in the pathogenesis of this monogenic obesity syndrome.

## Background

Most neurons in the vertebrate nervous system elaborate primary cilia. Historically, neuronal primary cilia were first identified in neuroepithelial progenitor cells projecting into the neural tube lumen. Later on, they were described to be broadly present both in neurons and glia [1]. The primary cilia function as sensory antennae in a wide variety of cells. Cilia-localized receptors, which include certain G protein-coupled receptors (GPCRs), and their downstream effectors, determine the sensory modality of cells in specific contexts, especially during vertebrate photoreception or olfaction and for responding to morphogens, such as Sonic hedgehog (Shh). Although we have come to appreciate the function of the primary cilium in other tissues and organisms, the functional roles of this ubiquitous neuronal organelle in integrating neuroendocrine signals have remained enigmatic. Diseases resulting from disruption of primary cilia and the associated basal body complex, called ciliopathies, often have strong neurological components, emphasizing the role of this cellular compartment in neural development [2]. Interestingly, aside from the strong neurodevelopmental phenotypes, progressive obesity often affects patients with certain ciliopathies such as the Bardet-Biedel Syndrome (BBS) and Alström Syndrome [3]. Notably, conditional knockout of components of the cilia in the mice hypothalamus results in hyperphagia-induced obesity and underscores the role of ciliary signaling in the central

regulation of satiety [4]. Thus, it is imperative to achieve a better understanding of the ciliary signaling pathways in central satiety networks, which could lead to novel ways for treating the global obesity pandemic.

The *tubby* mouse was initially identified as a spontaneous maturity-onset obesity syndrome [5], and positional cloning strategies in the 1990s mapped the causative mutation to a novel gene of unknown function called *Tub* [6,7]. In nematodes, *tub-1*, the canonical *Tub* homolog was identified in an RNAi screen for fat storage defects [8], and was found to be expressed in the ciliated neurons [9], highlighting the role of neuroendocrine signals in maintenance of systemic fat homeostasis even in these distant evolutionary relatives. Thanks to a recent paper from Sun *et al.* [10] the *tubby* mouse can now be added to the growing list of monogenic obesity syndromes with a strong ciliary functional component in the central nervous system [3]. The authors demonstrate molecular defects in ciliary GPCR signaling in the *tubby* mice, suggesting the importance of ciliary GPCR trafficking in central neurons implicated in satiety circuits.

The authors show that in the *tubby* mice, the primary cilia in the neurons show no obvious structural defects. However, two ciliary GPCRs, melanin-concentrating hormone receptor 1 (*Mchr1*) and somatostatin receptor subtype 3 (*Sstr3*), known to localize to distinct regions of the brain [11,12], are strongly prevented from trafficking to the primary cilia. This phenotype is strongly reminiscent of a previous study showing defective ciliary targeting of these receptors in BBS mice [12]. Similar to the BBS mice, the *tubby* mice also display retinal degeneration, and a defect in trafficking of rhodopsin to the outer segment of

\* Correspondence: Saikat.Mukhopadhyay@utsouthwestern.edu

<sup>1</sup>Department of Research Oncology, Genentech Inc, South San Francisco, CA 94080, USA

<sup>2</sup>Department of Cell Biology, UT Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, TX 75390, USA

the photoreceptor, an extension of the connecting cilia in these cells. However, in distinction from the BBS mice that have defective olfactory cilia and are anosmic [13], *tubby* mice do not show defects in either the structure of these specialized cilia or in localization of olfactory GPCRs. This difference could be because *Tub* is not expressed and does not play a major role in these specific neurons, or because the presence of other *tubby* family homologs (such as *Tulp3*) compensates for the loss of *Tub* activity. The authors also detect the defects in ciliary GPCR trafficking well in advance of the development of obesity and retinal degeneration, implying that these trafficking defects could be causative for the development of these phenotypes.

How does *Tubby* affect ciliary GPCR trafficking? The *Tub* gene is the founding member of a family of proteins [14], characterized by a C-terminal *tubby* domain, which is highly specific for binding to 4,5 phosphoinositides (PIP<sub>2</sub>) [15]. This domain is likely to participate in binding to specific membrane compartments, which for *Tubby* may be the ciliary membrane. Some of the *tubby* family members (including *Tub*, *Tulp2* and *Tulp3*) also have a signature motif in the divergent N-terminus that binds to the core subunits of the ciliary intraflagellar transport complex-A (IFT-A) [16]. *Tulp3* mutant mice are embryonic lethal by mid-gestation [17], but previous *in vitro* studies with heterologous cultured ciliated cell lines suggested that both *Tulp3* and IFT-A core subcomplex direct GPCR trafficking to the cilia [16]. Careful mutational analysis of both the IFT-A binding N-terminal and PIP<sub>2</sub>-binding C-terminal domains suggest that both the IFT-A- and membrane phosphoinositide-binding properties of TULP3 are necessary for ciliary GPCR localization. TULP3 thus bridges the IFT-A complex to the membrane compartment in gating ciliary GPCR trafficking, although the specific mechanism of ciliary GPCR recruitment remains to be determined. Most importantly, in the context of neuronal ciliary GPCR trafficking, the *Tulp3* N-terminal fragment can act as a dominant negative reagent, preventing GPCR trafficking in cultured hippocampal neurons [16]. *Tub* also shares the IFT-A binding motif with *Tulp3*, and binds to the IFT-A complex [16], although possibly less efficiently. Thus, similar to *Tulp3*, *Tub* could be directing ciliary GPCR trafficking through its simultaneous binding to the IFT-A complex, and membrane phosphoinositides. Presumably, higher levels of *Tub* in the brain could compensate for the lower binding or weaker affinity of *Tub* for the IFT-A complex. According to the Allen Brain Atlas, hypothalamic *Tub* transcript levels are about 26 times that of *Tulp3*. However, *Tub*/IFT-A might also require additional factors. Besides, the dominant negative IFT-A-binding N-terminal fragment of *Tulp3* would be expected to inhibit both *Tulp3* and *Tub* binding to the IFT-A complex in these neurons, effectively shutting down complementary effects

of these proteins in trafficking ciliary GPCRs. Thus, based on the spatial and temporal expression of these specific *tubby* family proteins in different tissues, and their affinity for the IFT-A complex, we might expect to observe a differential effect in their relative capacities for gating ciliary GPCRs. These differences could create a combinatorial code by utilizing an identical molecular mechanism for fine-tuning levels of ciliary receptors.

A suggestion implicit in the authors' findings is that the GPCR trafficking defects into the neuronal cilia, especially *Mchr1*, could underlie the obesity phenotype in the *tubby* mice. *Mchr1*, the receptor for melanin-concentrating hormone (MCH), is involved in the regulation of feeding and energy balance [18,19]. However, *Mchr1* knockout mice are lean [18,19], whereas MCH overexpression results in obesity [20]. Thus, in the simplest model, *Mchr1* trafficking defect to the cilia should mirror its effect on energy balance and cause leanness, rather than obesity, as evident in the *tubby* mouse. Dissecting the downstream effectors of *Mchr1* in regulating energy balance could address the conflicting effects of *Mchr1* trafficking on obesity. The best downstream effector implicated in neuronal satiety pathways is the adenylyl cyclase, type 3 (ACIII). Mice deficient in ACIII become obese with age, suggesting that ACIII-mediated cAMP signals are critical in the hypothalamus [21]. In line with this observation, downstream effectors of MCHR1 signaling include multiple G proteins including G<sub>i</sub>, G<sub>o</sub> and G<sub>q</sub> [22]. Thus, MCHR1 inhibits cAMP production stimulated by forskolin and increases intracellular Ca<sup>2+</sup> levels. However, in metabolically active brain slices, it paradoxically increases extracellular signal-regulated kinase (ERK) phosphorylation to levels above those observed with forskolin alone [23]. Thus, the synergistic effects of *Mchr1* signaling on cAMP, Ca<sup>2+</sup>, and ERK phosphorylation could be important in determining the final outcome on promoting energy intake.

Another possibility is the role of additional ciliary GPCRs in neuronal satiety centers, and a combination of trafficking defects of these receptors could result in the final maturity-onset obesity phenotype. For example, other neuronal GPCRs such as D1, D2, and D5 dopamine receptors are also expressed in neuronal cilia [24], and were not examined in this study. Besides, it is important to note that our catalog of GPCRs expressed in neuronal cilia is mostly incomplete. Thus, although the exact molecular explanation for obesity in the *tubby* (and BBS) mice is far from clear, we still favor the hypothesis that mislocalization of other novel, yet-unidentified GPCRs could provide us with a more complete answer in the future. Nevertheless, the final acid test for dissecting the role of ciliary trafficking of these individual receptors on neuronal phenotypes would entail detailed engineering of *knock-in* mice, expressing ciliary

localization-defective variants into the endogenous genomic loci of these receptors.

Another central question is the means by which ciliary signaling impacts neurons, and the reason why neurons need this signaling organelle in the first place. Currently, this is best answered in the case of morphogenetic developmental processes involving Shh signaling, which impacts neuronal differentiation both during embryogenesis and later stages. For example, Shh signaling in the cilia is fundamentally important in the neural progenitor cells during neural tube patterning [25]. Many of the Shh signaling components are localized to the cilia-basal body complex, and downstream signaling mediated by protein kinase A (PKA) and Gli3 processing are intricately linked to this organelle. In a broader developmental context, primary cilia are also fundamentally important in neurogenesis in cerebellar granule neurons [26,27], hippocampal neurogenesis in the dentate gyrus (DG) [28,29], adult DG neural stem cells [30], and in cerebral cortical development [31,32]. At least, in some of these neuronal cells, the primary cilium probably acts as a subcellular compartment for efficiently amplifying extracellular Shh signals for intracellular signal transduction. However, neither *tubby* nor BBS mice demonstrate gross defects in the neuroanatomical networks that regulate satiety, suggesting by extension that a lack of GPCR trafficking in these neurons probably would not cause apparent deficits during development of these circuits.

Apart from the role of cilia in Shh signaling and differentiation, recent studies have begun to provide intriguing molecular insights into other neuron-dependent processes dependent on the presence of cilia, and similar mechanisms could impact the satiety networks in a cilia-dependent manner. First of all, primary cilia function in glutamatergic synaptic integration of adult-born neurons [33]. Conditional deletion of cilia from adult-born neurons induces severe defects in dendritic refinement and synapse formation, which is partially correlated with an enhancement of Wnt and  $\beta$ -catenin signaling [33]. Signaling in the context of primary cilia could thus eventually impinge upon the subsequent efficient integration of neurons into neural networks. Second, primary ciliary signaling has also been shown to have an effect on long-term potentiation (LTP) and plasticity [34]. *Sstr3* signaling in the hippocampus is important in novelty detection in mice, and adenylyl cyclase/cAMP-mediated LTP is impaired in hippocampal slices from the *Sstr3* knockout and upon addition of *Sstr3* antagonists into wild-type sections. In this case, cilia could act as coincidence detectors and affect synaptic plasticity by affecting downstream signaling pathways. On a similar note, dopamine produces a synapse-specific enhancement of early LTP through D1/D5 receptors and cAMP signaling [35]. Future work is needed to establish if efficient targeting of

these receptors to the neuronal cilia is important in these processes. Identifying downstream pathways regulating synaptic plasticity particularly promises to be an important future avenue of research for understanding the puzzling role of cilia in neuronal function.

#### Competing interest

The authors declare that they have no competing interests.

#### Authors' contribution

Both authors read and approved the final manuscript.

Received: 6 November 2012 Accepted: 7 November 2012

Published: 3 January 2013

#### References

1. Whitfield JF (2004) The neuronal primary cilium—an extrasynaptic signaling device. *Cell Signal* 16:763–767
2. Hildebrandt F, Benzing T, Katsanis N (2011) Ciliopathies. *N Engl J Med* 364:1533–1543
3. Beales PL, Farooqi IS, O'Rahilly S (2009) *The Genetics of Obesity Syndromes*. Oxford University Press, New York
4. Davenport JR, Watts AJ, Roper VC, Croyle MJ, van Groen T, Wyss JM, Nagy TR, Kesterson RA, Yoder BK (2007) Disruption of intraflagellar transport in adult mice leads to obesity and slow-onset cystic kidney disease. *Curr Biol* 17:1586–1594
5. Coleman DL, Eicher EM (1990) Fat (fat) and tubby (tubby): two autosomal recessive mutations causing obesity syndromes in the mouse. *J Hered* 81:424–427
6. Kleyn PW, Fan W, Kovats SG, Lee JJ, Pulido JC, Wu Y, Berkemeier LR, Misumi DJ, Holmgren L, Charlat O, Woolf EA, Tayber O, Brody T, Shu P, Hawkins F, Kennedy B, Baldini L, Ebeling C, Alperin GD, Deeds J, Lakey ND, Culpepper J, Chen H, Glücksmann-Kuis MA, Carlson GA, Duyk GM, Moore KJ (1996) Identification and characterization of the mouse obesity gene *tubby*: a member of a novel gene family. *Cell* 85(2):281–290
7. Noben-Trauth K, Naggert JK, North MA, Nishina PM (1996) A candidate gene for the mouse mutation *tubby*. *Nature* 380:534–538
8. Ashrafi K, Chang FY, Watts JL, Fraser AG, Kamath RS, Ahringer J, Ruvkun G (2003) Genome-wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature* 421:268–272
9. Mak HY, Nelson LS, Basson M, Johnson CD, Ruvkun G (2006) Polygenic control of *Caenorhabditis elegans* fat storage. *Nat Genet* 38:363–368
10. Sun X, Haley J, Bulgakovolev OV, Cai X, McGinnis J, Li T (2012) *Tubby* is required for trafficking g protein-coupled receptors to neuronal cilia. *Cilia* 1:21
11. Handel M, Schulz S, Stanarius A, Schreff M, Erdtmann-Vourliotis M, Schmidt H, Wolf G, Hollt V (1999) Selective targeting of somatostatin receptor 3 to neuronal cilia. *Neuroscience* 89:909–926
12. Berbari NF, Lewis JS, Bishop GA, Askwith CC, Mykityn K (2008) Bardet-Biedl syndrome proteins are required for the localization of G protein-coupled receptors to primary cilia. *Proc Natl Acad Sci USA* 105:4242–4246
13. Kulaga HM, Leitch CC, Eichers ER, Badano JL, Lesemann A, Hoskins BE, Lupski JR, Beales PL, Reed RR, Katsanis N (2004) Loss of BBS proteins causes anosmia in humans and defects in olfactory cilia structure and function in the mouse. *Nat Genet* 36:994–998
14. Mukhopadhyay S, Jackson PK (2011) The *tubby* family proteins. *Genome Biol* 12:225
15. Santagata S, Boggon TJ, Baird CL, Gomez CA, Zhao J, Shan WS, Myszka DG, Shapiro L (2001) G-protein signaling through *tubby* proteins. *Science* 292:2041–2050
16. Mukhopadhyay S, Wen X, Chih B, Nelson CD, Lane WS, Scales SJ, Jackson PK (2010) TULP3 bridges the IFT-A complex and membrane phosphoinositides to promote trafficking of G protein-coupled receptors into primary cilia. *Genes Dev* 24:2180–2193
17. Norman RX, Ko HW, Huang V, Eun CM, Ablor LL, Zhang Z, Sun X, Eggenschwiler JT (2009) *Tubby*-like protein 3 (TULP3) regulates patterning in the mouse embryo through inhibition of Hedgehog signaling. *Hum Mol Genet* 18:1740–1754

18. Shimada M, Tritos NA, Lowell BB, Flier JS, Maratos-Flier E (1998) Mice lacking melanin-concentrating hormone are hypophagic and lean. *Nature* 396:670–674
19. Chen Y, Hu C, Hsu CK, Zhang Q, Bi C, Asnicar M, Hsiung HM, Fox N, Sliker LJ, Yang DD, Heiman ML, Shi Y (2002) Targeted disruption of the melanin-concentrating hormone receptor-1 results in hyperphagia and resistance to diet-induced obesity. *Endocrinology* 143:2469–2477
20. Ludwig DS, Tritos NA, Mastaitis JW, Kulkarni R, Kokkotou E, Elmquist J, Lowell B, Flier JS, Maratos-Flier E (2001) Melanin-concentrating hormone overexpression in transgenic mice leads to obesity and insulin resistance. *J Clin Invest* 107:379–386
21. Wang Z, Li V, Chan GC, Phan T, Nudelman AS, Xia Z, Storm DR (2009) Adult type 3 adenylyl cyclase-deficient mice are obese. *PLoS One* 4:e6979
22. Hawes BE, Kil E, Green B, O'Neill K, Fried S, Graziano MP (2000) The melanin-concentrating hormone receptor couples to multiple G proteins to activate diverse intracellular signaling pathways. *Endocrinology* 141:4524–4532
23. Pissios P, Trombly DJ, Tzamei I, Maratos-Flier E (2003) Melanin-concentrating hormone receptor 1 activates extracellular signal-regulated kinase and synergizes with G(s)-coupled pathways. *Endocrinology* 144:3514–3523
24. Marley A, von Zastrow M (2010) DISC1 regulates primary cilia that display specific dopamine receptors. *PLoS One* 5:e10902
25. Goetz SC, Anderson KV (2010) The primary cilium: a signalling centre during vertebrate development. *Nat Rev Genet* 11:331–344
26. Chizhikov W, Davenport J, Zhang Q, Shih EK, Cabello OA, Fuchs JL, Yoder BK, Millen KJ (2007) Cilia proteins control cerebellar morphogenesis by promoting expansion of the granule progenitor pool. *J Neurosci* 27:9780–9789
27. Spassky N, Merkle FT, Flames N, Tramontin AD, Garcia-Verdugo JM, Alvarez-Buylla A (2005) Adult ependymal cells are postmitotic and are derived from radial glial cells during embryogenesis. *J Neurosci* 25:10–18
28. Breunig JJ, Sarkisian MR, Arellano JI, Morozov YM, Ayoub AE, Sojitra S, Wang B, Flavell RA, Rakic P, Town T (2008) Primary cilia regulate hippocampal neurogenesis by mediating sonic hedgehog signaling. *Proc Natl Acad Sci USA* 105:13127–13132
29. Han YG, Kim HJ, Dlugosz AA, Ellison DW, Gilbertson RJ, Alvarez-Buylla A (2009) Dual and opposing roles of primary cilia in medulloblastoma development. *Nat Med* 15:1062–1065
30. Amador-Arjona A, Elliott J, Miller A, Ginbey A, Pazour GJ, Enikolopov G, Roberts AJ, Tersikh AV (2011) Primary cilia regulate proliferation of amplifying progenitors in adult hippocampus: implications for learning and memory. *J Neurosci* 31:9933–9944
31. Stottmann RW, Tran PV, Turbe-Doan A, Beier DR (2009) Ttc21b is required to restrict sonic hedgehog activity in the developing mouse forebrain. *Dev Biol* 335:166–178
32. Willaredt MA, Hasenpusch-Theil K, Gardner HA, Kitanovic I, Hirschfeld-Warneken VC, Gojak CP, Gorgas K, Bradford CL, Spatz J, Wolf S, Theil T, Tucker KL (2008) A crucial role for primary cilia in cortical morphogenesis. *J Neurosci* 28:12887–12900
33. Kumamoto N, Gu Y, Wang J, Janoschka S, Takemaru K, Levine J, Ge S (2012) A role for primary cilia in glutamatergic synaptic integration of adult-born neurons. *Nat Neurosci* 15:399–405. S1
34. Einstein EB, Patterson CA, Hon BJ, Regan KA, Reddi J, Melnikoff DE, Mateer MJ, Schulz S, Johnson BN, Tallent MK (2010) Somatostatin signaling in neuronal cilia is critical for object recognition memory. *J Neurosci* 30:4306–4314
35. Otmakhova NA, Lisman JE (1996) D1/D5 dopamine receptor activation increases the magnitude of early long-term potentiation at CA1 hippocampal synapses. *J Neurosci* 16:7478–7486

doi:10.1186/2046-2530-2-1

Cite this article as: Mukhopadhyay and Jackson: *Cilia, tubby mice, and obesity*. *Cilia* 2013 **2**:1.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

