



Crystal structure of 4-bromoanilinium 4-methylbenzenesulfonate

P. K. Sivakumar,^a M. Krishna Kumar,^b R. Mohan Kumar,^b
G. Chakkaravarthi^{c*} and R. Kanagadurai^{b*}

^aDepartment of Physics, M. N. M. Jain Engineering College, Chennai 600 097, India,

^bDepartment of Physics, Presidency College, Chennai 600 005, India, and

^cDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India.

*Correspondence e-mail: chakkaravarthi_2005@yahoo.com, srkanagadurai@yahoo.co.in

Received 5 February 2015; accepted 8 February 2015

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


In the crystal of the title molecular salt, C₆H₇BrN⁺·C₇H₇O₃S⁻, the anions and cations are linked *via* N—H···O hydrogen bonds forming layers, enclosing R₂²(4) ring motifs, lying parallel to (001). Within the layers there are short O···O contacts of 2.843 (2) Å.

Keywords: crystal structure; anilinium; 4-methylbenzenesulfonate; hydrogen bonding.

CCDC reference: 1048164

1. Related literature

For the crystal structure of 4-chloroanilinium 4-methylbenzenesulfonate, isostructural with the title salt, see: Jasinski *et al.* (2011). For the crystal structure of other 4-methylbenzenesulfonate salts, see, for example: Krishnakumar *et al.* (2012); Sudhahar *et al.* (2013).



2. Experimental

2.1. Crystal data

C₆H₇BrN⁺·C₇H₇O₃S⁻

$M_r = 344.22$

Triclinic, $P\bar{1}$

$a = 5.7908$ (3) Å

$b = 7.6004$ (4) Å

$c = 15.9073$ (7) Å

$\alpha = 94.716$ (2)°

$\beta = 96.520$ (3)°

$\gamma = 92.732$ (2)°

$V = 692.09$ (6) Å³

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 3.12$ mm⁻¹

$T = 295$ K

0.24 × 0.20 × 0.18 mm

2.2. Data collection

Bruker Kappa APEXII CCD
diffractometer

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.521$, $T_{\max} = 0.603$

11503 measured reflections

3001 independent reflections

2526 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.083$

$S = 1.03$

3001 reflections

175 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.45$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.56$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1A···O2	0.89	2.07	2.879 (2)	151
N1—H1C···O1 ⁱ	0.89	2.46	2.971 (2)	117
N1—H1A···O2 ⁱⁱ	0.89	2.46	3.096 (2)	129
N1—H1B···O3 ⁱⁱⁱ	0.89	1.91	2.794 (2)	172
N1—H1C···O1 ⁱⁱⁱ	0.89	2.04	2.904 (2)	165

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $x, y - 1, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

Acknowledgements

The authors thank the SAIF, IIT Madras, for the data collection.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU5079).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jasinski, J. P., Golen, J. A., Praveen, A. S., Yathirajan, H. S. & Narayana, B. (2011). *Acta Cryst.* **E67**, o3288–o3289.
- Krishnakumar, M., Sudhahar, S., Silambarasan, A., Chakkaravarthi, G. & Mohankumar, R. (2012). *Acta Cryst.* **E68**, o3268.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

Sudhahar, S., Krishnakumar, M., Sornamurthy, B. M., Chakkaravarthi, G. & Mohankumar, R. (2013). *Acta Cryst.* **E69**, o279.

supporting information

Acta Cryst. (2015). E71, o163–o164 [doi:10.1107/S2056989015002686]

Crystal structure of 4-bromoanilinium 4-methylbenzenesulfonate

P. K. Sivakumar, M. Krishna Kumar, R. Mohan Kumar, G. Chakkaravarthi and R. Kanagadurai

S1. Structural commentary

The molecular structure of the title salt is illustrated in Fig. 1. The bond lengths and angles are similar to those reported for the 4-chloroanilinium analogue (Jasinski *et al.*, 2011), and for other similar 4-methylbenzenesulfonate salts (Krishnakumar *et al.*, 2012; Sudhahar *et al.*, 2013).

In the crystal, the benzene rings (C1—C6) and (C8—C13) make a dihedral angle of 50.89 (11) °. The cations and anions are linked via N—H···O hydrogen bonds which generates $R_2^2(4)$ ring motifs, and form layers parallel to (001); see Table 1 and Fig. 2. Within the layers there are short O1···O1ⁱ contacts of 2.843 (2) Å [symmetry code: (i) -x+2, -y+2, -z+1].

S2. Synthesis and crystallization

The title compound was synthesized in ethanol by mixing 4-bromoaniline (2.37 g) and *p*-toluenesulfonic acid (2.62 g) in an equimolar ratio. The saturated solution was allowed to evaporating slowly at room temperature. After a period of three weeks colourless block-like crystals appeared.

S3. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms were positioned geometrically and refined using a riding model: N—H = 0.89 Å, C—H = 0.93 - 96 Å with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N,C})$ for the ammonium and methyl H atoms and = $1.2U_{\text{eq}}(\text{C})$ for other H atoms.




Figure 1

The molecular structure of the title molecular salt, with atom labelling. Displacement ellipsoids are drawn at the 30% probability level.




Figure 2

The crystal packing of the title molecular salt, viewed along the a axis. Hydrogen bonds are shown as dashed lines (see Table 1 for details; H atoms not involved in hydrogen bonding have been omitted for clarity).

4-Bromoanilinium 4-methylbenzenesulfonate

Crystal data

$C_6H_7BrN^+ \cdot C_7H_7O_3S^-$

$M_r = 344.22$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 5.7908\ (3)\ \text{\AA}$

$b = 7.6004\ (4)\ \text{\AA}$

$c = 15.9073\ (7)\ \text{\AA}$

$\alpha = 94.716\ (2)^\circ$

$\beta = 96.520\ (3)^\circ$

$\gamma = 92.732\ (2)^\circ$

$V = 692.09\ (6)\ \text{\AA}^3$

$Z = 2$

$F(000) = 3048$

$D_x = 1.652\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 348 reflections

$\theta = 1.3\text{--}27.1^\circ$

$\mu = 3.12\ \text{mm}^{-1}$

$T = 295$ K $0.24 \times 0.20 \times 0.18$ mm
 Block, colourless

Data collection

Bruker Kappa APEXII CCD diffractometer Radiation source: fine-focus sealed tube Graphite monochromator ω and φ scan Absorption correction: multi-scan (SADABS; Sheldrick, 1996) $T_{\min} = 0.521$, $T_{\max} = 0.603$	11503 measured reflections 3001 independent reflections 2526 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.030$ $\theta_{\text{max}} = 27.1^\circ$, $\theta_{\text{min}} = 1.3^\circ$ $h = -7 \rightarrow 7$ $k = -9 \rightarrow 9$ $l = -20 \rightarrow 20$
---	--

Refinement

Refinement on F^2 Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.033$ $wR(F^2) = 0.083$ $S = 1.03$ 3001 reflections 175 parameters 0 restraints Primary atom site location: structure-invariant direct methods Secondary atom site location: difference Fourier map	Hydrogen site location: inferred from neighbouring sites H-atom parameters constrained $w = 1/[\sigma^2(F_o^2) + (0.0406P)^2 + 0.3371P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.45 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.56 \text{ e } \text{\AA}^{-3}$ Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$ Extinction coefficient: 0.034 (2)
---	--

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.0889 (4)	0.8014 (3)	0.89050 (15)	0.0401 (6)
C2	1.2329 (4)	0.8603 (4)	0.83410 (17)	0.0428 (6)
H2	1.3820	0.9068	0.8546	0.051*
C3	1.1611 (4)	0.8518 (3)	0.74779 (15)	0.0353 (5)
H3	1.2613	0.8909	0.7106	0.042*
C4	0.9385 (3)	0.7845 (3)	0.71760 (12)	0.0233 (4)
C5	0.7913 (4)	0.7260 (3)	0.77291 (15)	0.0351 (5)
H5	0.6416	0.6807	0.7525	0.042*
C6	0.8673 (4)	0.7348 (4)	0.85901 (15)	0.0439 (6)
H6	0.7674	0.6952	0.8962	0.053*
C7	1.1741 (6)	0.8053 (5)	0.98404 (18)	0.0686 (9)
H7A	1.2377	0.6944	0.9958	0.103*

H7B	1.0464	0.8251	1.0166	0.103*
H7C	1.2922	0.8991	0.9992	0.103*
C8	0.4302 (4)	0.3193 (3)	0.77075 (14)	0.0317 (5)
C9	0.6357 (4)	0.2352 (3)	0.77106 (15)	0.0342 (5)
H9	0.7066	0.1958	0.8209	0.041*
C10	0.7353 (4)	0.2102 (3)	0.69649 (14)	0.0296 (5)
H10	0.8749	0.1548	0.6958	0.036*
C11	0.6259 (3)	0.2681 (3)	0.62306 (13)	0.0253 (4)
C12	0.4203 (4)	0.3529 (3)	0.62302 (14)	0.0309 (5)
H12	0.3492	0.3920	0.5731	0.037*
C13	0.3209 (4)	0.3792 (3)	0.69759 (15)	0.0329 (5)
H13	0.1826	0.4363	0.6986	0.039*
N1	0.7352 (3)	0.2443 (2)	0.54491 (11)	0.0301 (4)
H1A	0.8264	0.3397	0.5401	0.045*
H1B	0.6256	0.2287	0.5006	0.045*
H1C	0.8204	0.1499	0.5464	0.045*
O1	0.9616 (3)	0.9187 (2)	0.57494 (10)	0.0387 (4)
O2	0.9290 (3)	0.6031 (2)	0.57251 (10)	0.0372 (4)
O3	0.5965 (3)	0.7733 (3)	0.59796 (10)	0.0460 (4)
Br1	0.29548 (5)	0.35751 (4)	0.873294 (18)	0.05714 (14)
S1	0.84823 (9)	0.76863 (7)	0.60718 (3)	0.02575 (14)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0476 (14)	0.0480 (14)	0.0238 (12)	0.0154 (11)	-0.0024 (10)	-0.0011 (10)
C2	0.0341 (12)	0.0518 (15)	0.0382 (14)	0.0012 (11)	-0.0083 (10)	-0.0035 (11)
C3	0.0298 (11)	0.0428 (13)	0.0327 (12)	-0.0022 (9)	0.0030 (9)	0.0032 (10)
C4	0.0287 (10)	0.0214 (9)	0.0200 (10)	0.0048 (8)	0.0029 (7)	0.0010 (8)
C5	0.0315 (11)	0.0463 (14)	0.0273 (12)	-0.0034 (10)	0.0038 (9)	0.0050 (10)
C6	0.0460 (14)	0.0625 (17)	0.0261 (12)	0.0034 (12)	0.0111 (10)	0.0106 (11)
C7	0.077 (2)	0.098 (3)	0.0278 (14)	0.0243 (19)	-0.0093 (14)	-0.0014 (16)
C8	0.0353 (11)	0.0310 (11)	0.0287 (12)	0.0018 (9)	0.0061 (9)	-0.0007 (9)
C9	0.0399 (12)	0.0358 (12)	0.0267 (11)	0.0083 (10)	-0.0007 (9)	0.0055 (9)
C10	0.0289 (10)	0.0283 (11)	0.0315 (12)	0.0075 (8)	0.0003 (8)	0.0034 (9)
C11	0.0263 (10)	0.0220 (10)	0.0262 (11)	-0.0013 (8)	0.0009 (8)	-0.0012 (8)
C12	0.0300 (11)	0.0321 (11)	0.0299 (12)	0.0054 (9)	-0.0020 (9)	0.0046 (9)
C13	0.0273 (11)	0.0333 (12)	0.0379 (13)	0.0070 (9)	0.0020 (9)	0.0015 (10)
N1	0.0298 (9)	0.0316 (9)	0.0286 (10)	0.0032 (7)	0.0019 (7)	0.0029 (8)
O1	0.0566 (10)	0.0325 (8)	0.0315 (9)	0.0091 (7)	0.0160 (7)	0.0109 (7)
O2	0.0531 (10)	0.0281 (8)	0.0297 (9)	0.0057 (7)	0.0065 (7)	-0.0053 (7)
O3	0.0321 (9)	0.0787 (13)	0.0263 (9)	0.0106 (8)	-0.0017 (7)	0.0025 (8)
Br1	0.0649 (2)	0.0722 (2)	0.03948 (19)	0.02182 (15)	0.02132 (13)	0.00502 (14)
S1	0.0313 (3)	0.0287 (3)	0.0179 (3)	0.0066 (2)	0.00382 (19)	0.0019 (2)

Geometric parameters (Å, °)

C1—C2	1.379 (4)	C8—Br1	1.894 (2)
C1—C6	1.380 (4)	C9—C10	1.380 (3)
C1—C7	1.511 (3)	C9—H9	0.9300
C2—C3	1.384 (3)	C10—C11	1.379 (3)
C2—H2	0.9300	C10—H10	0.9300
C3—C4	1.382 (3)	C11—C12	1.381 (3)
C3—H3	0.9300	C11—N1	1.460 (3)
C4—C5	1.377 (3)	C12—C13	1.380 (3)
C4—S1	1.767 (2)	C12—H12	0.9300
C5—C6	1.385 (3)	C13—H13	0.9300
C5—H5	0.9300	N1—H1A	0.8900
C6—H6	0.9300	N1—H1B	0.8900
C7—H7A	0.9600	N1—H1C	0.8900
C7—H7B	0.9600	O1—S1	1.4480 (16)
C7—H7C	0.9600	O2—S1	1.4513 (16)
C8—C9	1.377 (3)	O3—S1	1.4509 (17)
C8—C13	1.382 (3)		
C2—C1—C6	118.3 (2)	C8—C9—C10	119.2 (2)
C2—C1—C7	120.7 (3)	C8—C9—H9	120.4
C6—C1—C7	120.9 (3)	C10—C9—H9	120.4
C1—C2—C3	121.6 (2)	C11—C10—C9	119.39 (19)
C1—C2—H2	119.2	C11—C10—H10	120.3
C3—C2—H2	119.2	C9—C10—H10	120.3
C4—C3—C2	119.1 (2)	C10—C11—C12	121.2 (2)
C4—C3—H3	120.5	C10—C11—N1	118.94 (18)
C2—C3—H3	120.5	C12—C11—N1	119.80 (18)
C5—C4—C3	120.20 (19)	C13—C12—C11	119.56 (19)
C5—C4—S1	120.52 (16)	C13—C12—H12	120.2
C3—C4—S1	119.26 (16)	C11—C12—H12	120.2
C4—C5—C6	119.7 (2)	C12—C13—C8	118.9 (2)
C4—C5—H5	120.1	C12—C13—H13	120.5
C6—C5—H5	120.1	C8—C13—H13	120.5
C1—C6—C5	121.0 (2)	C11—N1—H1A	109.5
C1—C6—H6	119.5	C11—N1—H1B	109.5
C5—C6—H6	119.5	H1A—N1—H1B	109.5
C1—C7—H7A	109.5	C11—N1—H1C	109.5
C1—C7—H7B	109.5	H1A—N1—H1C	109.5
H7A—C7—H7B	109.5	H1B—N1—H1C	109.5
C1—C7—H7C	109.5	O1—S1—O3	113.04 (11)
H7A—C7—H7C	109.5	O1—S1—O2	111.33 (10)
H7B—C7—H7C	109.5	O3—S1—O2	113.23 (11)
C9—C8—C13	121.7 (2)	O1—S1—C4	106.17 (10)
C9—C8—Br1	119.41 (16)	O3—S1—C4	106.11 (10)
C13—C8—Br1	118.93 (17)	O2—S1—C4	106.31 (9)

C6—C1—C2—C3	-0.8 (4)	C9—C10—C11—C12	0.9 (3)
C7—C1—C2—C3	177.6 (3)	C9—C10—C11—N1	178.73 (19)
C1—C2—C3—C4	0.7 (4)	C10—C11—C12—C13	-0.5 (3)
C2—C3—C4—C5	-0.3 (3)	N1—C11—C12—C13	-178.31 (19)
C2—C3—C4—S1	-178.54 (19)	C11—C12—C13—C8	-0.1 (3)
C3—C4—C5—C6	0.0 (3)	C9—C8—C13—C12	0.4 (3)
S1—C4—C5—C6	178.18 (19)	Br1—C8—C13—C12	179.35 (16)
C2—C1—C6—C5	0.4 (4)	C5—C4—S1—O1	147.56 (18)
C7—C1—C6—C5	-178.0 (3)	C3—C4—S1—O1	-34.23 (19)
C4—C5—C6—C1	0.0 (4)	C5—C4—S1—O3	27.0 (2)
C13—C8—C9—C10	0.0 (3)	C3—C4—S1—O3	-154.76 (18)
Br1—C8—C9—C10	-178.95 (17)	C5—C4—S1—O2	-93.79 (19)
C8—C9—C10—C11	-0.7 (3)	C3—C4—S1—O2	84.42 (19)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots O2	0.89	2.07	2.879 (2)	151
N1—H1C \cdots O1 ⁱ	0.89	2.46	2.971 (2)	117
N1—H1A \cdots O2 ⁱ	0.89	2.46	3.096 (2)	129
N1—H1B \cdots O3 ⁱⁱ	0.89	1.91	2.794 (2)	172
N1—H1C \cdots O1 ⁱⁱⁱ	0.89	2.04	2.904 (2)	165

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, y-1, z$.