

CRYSTALLOGRAPHIC
COMMUNICATIONS

ISSN 2056-9890

Crystal structure of 2-isopropyl-5,7'-dimethyl-1',3',3a',6',8a',8b'-hexahydrospiro[cyclohexane-1,6'-furo[3,4-*d*]imidazo[1,5-*b*]isoxazol]-8'(7'*H*)-one

Heithem Abda,^{a*} Khaireddine Ezzayani,^b Kaiss Aouadi,^a Taha Guerfel,^c Sebastien Vidal^d and Moncef Msaddek^a

Received 4 June 2016

Accepted 30 June 2016

Edited by G. Smith, Queensland University of Technology, Australia

Keywords: crystal structure; isoxazolidine; hydrogen bonding.

CCDC reference: 1489217



Supporting information: this article has supporting information at journals.iucr.org/e

^aUniversity of Monastir, Heterocyclic Chemistry Laboratory, Products, Natural and Reactivity, Faculty of Sciences of Monastir, Avenue of the Environment, 5000 Monastir, Tunisia, ^bUniversity of Monastir, Laboratory of Physical Chemistry of Materials, Faculty of Sciences of Monastir, Avenue of the Environment, 5019 Monastir, Tunisia, ^cLaboratory of Electrochemistry, Materials and Environment, Kairouan University, 3100 Kairouan, Tunisia, and ^dUniversity of Lyon CNRS, Institute of Chemistry and Biochemistry and Molecular Supramolecular, UMR 5246, Laboratoire de Chimie Organique 2-Glycochemistry, Curien Building, 43 Boulevard du 11 Novembre 1918, F-69622 Villeurbanne, France.
*Correspondence e-mail: abda_he@hotmail.fr

In the title compound, C₁₇H₂₈N₂O₃, the isoxazolidine ring adopts an envelope conformation with the O atom deviating from the mean plane of the other four ring atoms by 0.617 (1) Å. In the crystal, molecules are linked *via* weak C—H⋯O hydrogen bonds, forming chains which extend along the *b*-axis direction.

1. Chemical context

The 1,3-dipolar cycloaddition of nitrones to alkenes has been applied to produce substituted isoxazolidines (Gothelf & Jørgensen, 1998). These compounds can be converted into β -amino alcohols (Padwa *et al.*, 2002), β -lactams (Hanselmann *et al.*, 2003) and α -amino acids (Aouadi *et al.*, 2006), by reductive cleavage of the N—O bond. Consequently, isoxazolidines have been used as key intermediates for the synthesis of various natural products or antifungal, anti-inflammatory, anti-mycobacterial, anti-tuberculosis and antiviral agents. The previously mentioned importance of the isoxazolidine substructure led us to investigate the cycloaddition of chiral nitronne [(5(*S*),6(*S*),9(*R*)-6-isopropyl-4,9-dimethyl-3-oxo-1,4-diazaspiro[4.5]dec-1-ene-1-oxide] with 2,5-dihydrofuran. The present work reports the synthesis and the X-ray crystallographic study of this substituted isoxazolidine, the title compound, C₁₇H₂₈N₂O₃, (I).



OPEN ACCESS




Figure 1
The molecular conformation in the molecules of (I), showing the atom labelling. Displacement ellipsoids are drawn at the 35% probability level. H atoms have been omitted for clarity.

2. Structural commentary

In the title compound (I), the asymmetric unit comprises a single molecule (Fig. 1). Each molecule has six stereogenic centres (Abda *et al.*, 2014) although the absolute configuration

Table 1
Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|----------------------|-------|-------------|-------------|---------------|
| $C5-H52\cdots O13^i$ | 0.97 | 2.57 | 3.536 (3) | 172 |

Symmetry code: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

for the molecule was not determined definitively in this analysis. The isoxazolidine ring (O1/N2/C7–C9) adopts an envelope conformation with atom O1 displaced by 0.617 (1) Å from the mean plane through atoms N2/C7–C9. The N–O bond lengths of the isoxazolidine rings O1–N2 = 1.482 (2) Å, close to values reported for related compounds (Loh *et al.*, 2010; Molander *et al.*, 2013).

3. Supramolecular features

In the crystal, the molecules are linked *via* non-classical weak $C5-H52\cdots O13^i$ hydrogen bonds, forming zigzag chains, which extend along the *b*-axis direction (Table 1 and Fig. 2).

4. Synthesis and crystallization

In a Biotage Initiator 10 ml vial, nitron [(5(*S*),6(*S*),9(*R*))-6-isopropyl-4,9-dimethyl-3-oxo-1,4-diazaspiro[4.5]dec-1-ene-1-oxide] (1 eq.) in anhydrous toluene (4 ml) was introduced.




Figure 2
The $C-H\cdots O$ hydrogen-bonded chains extending along the *b* axis in the crystal structure of (I). Dashed lines indicate hydrogen bonds. Non-associated H atoms have been omitted.




Figure 3
The cycloaddition reaction in the synthesis of (I).

The vial was flushed with argon and 2,5-dihydrofuran (3 eq.) was added. The vial was sealed with a septum cap and was irradiated with microwaves (temperature: 373 K) (Fig. 3). TLC monitoring (EtOAc/PE 5/5) showed full conversion after 2 h. After the crude mixture was concentrated and purified by flash column chromatography (silica gel, EtOAc/PE 5/5), the desired isoxazolidine (I) was obtained (m.p. = 410–411 K).

5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. The H atoms were located in a difference map, but these were repositioned geometrically and were initially refined with soft restraints on the bond lengths and angles to regularize their geometry (C–H in the range 0.93–0.98 Å) and $U_{\text{iso}}(\text{H})$ (in the range 1.2–1.5 times U_{eq} of the parent atom). These were subsequently refined with riding constraints (Cooper *et al.*, 2010). Although not definitive for this chiral structure, the Flack (1983) absolute structure parameter obtained [0.60 (3) for 1261 Friedel pairs] gave C3(*S*), C7(*S*), C8(*S*), C9(*S*), C14(*S*), C20(*R*) assignments for the six arbitrarily named chiral centres in the molecule. The inverted structure gave a similarly high Flack factor.

Acknowledgements

The authors are grateful to the Ministry of Higher Education and Scientific Research of Tunisia for financial support.

References

Abda, H., Aouadi, K., Perrin, L., Msaddek, M., Praly, J.-P. & Vidal, S. (2014). *Eur. J. Org. Chem.* pp. 6017–6024.
 Agilent (2013). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
 Aouadi, K., Vidal, S., Msaddek, M. & Praly, J.-P. (2006). *Synlett*, **2006**, 3299–3303.
 Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
 Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.

Table 2
Experimental details.

| | |
|------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Crystal data | |
| Chemical formula | C ₁₇ H ₂₈ N ₂ O ₃ |
| M_r | 308.42 |
| Crystal system, space group | Orthorhombic, $P2_12_12_1$ |
| Temperature (K) | 150 |
| a, b, c (Å) | 7.7474 (6), 11.1404 (8), 19.208 (2) |
| V (Å ³) | 1657.8 (2) |
| Z | 4 |
| Radiation type | Cu $K\alpha$ |
| μ (mm ⁻¹) | 0.68 |
| Crystal size (mm) | 0.49 × 0.43 × 0.25 |
| Data collection | |
| Diffractometer | Oxford Diffraction Xcalibur (Atlas, Gemini Ultra) |
| Absorption correction | Analytical [<i>CrysAlis PRO</i> (Agilent, 2013) based on expressions derived by Clark & Reid (1995); changes in illuminated volume were kept to a minimum, and were taken into account (Görlitz, 1999)] |
| $T_{\text{min}}, T_{\text{max}}$ | 0.782, 0.866 |
| No. of measured, independent and observed [$I > 2.0\sigma(I)$] reflections | 10374, 2879, 2680 |
| R_{int} | 0.059 |
| $(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹) | 0.596 |
| Refinement | |
| $R[F^2 > 2\sigma(F^2)], wR(F^2), S$ | 0.042, 0.096, 1.03 |
| No. of reflections | 2866 |
| No. of parameters | 201 |
| H-atom treatment | H-atom parameters constrained |
| $\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³) | 0.16, -0.17 |
| Absolute structure | Flack (1983), 1261 Friedel pairs |
| Absolute structure parameter | 0.6 (3) |

Computer programs: *CrysAlis PRO* (Agilent, 2013), *SIR97* (Altomare *et al.*, 1999), *CRYSTALS* (Betteridge *et al.*, 2003), *CAMERON* (Watkin *et al.*, 1996), Larson (1970), Prince (1982) and Watkin (1994).

Cooper, R. I., Thompson, A. L. & Watkin, D. J. (2010). *J. Appl. Cryst.* **43**, 1100–1107.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Görlitz, C. H. (1999). *Acta Cryst.* **B55**, 1090–1098.
 Gothelf, K. V. & Jørgensen, K. A. (1998). *Chem. Rev.* **98**, 863–910.
 Hanselmann, R., Zhou, J., Ma, P. & Confalone, P. N. (2003). *J. Org. Chem.* **68**, 8739–8741.
 Larson, A. C. (1970). *Crystallographic Computing*, edited by F. R. Ahmed, pp. 291–294. Copenhagen: Munksgaard.
 Loh, B., Vozzolo, L., Mok, B. J., Lee, C. C., Fitzmaurice, R. J., Caddick, S. & Fassati, A. (2010). *Chem. Biol. Drug Des.* **75**, 461–474.
 Molander, G. A., Cavalcanti, L. N. & García-García, C. (2013). *J. Org. Chem.* **78**, 6427–6439.
 Padwa, A. & Pearson, W. H. (2002). Editors. *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*, Vol. 59, pp. 1–81. Chichester: Wiley.
 Prince, E. (1982). In *Mathematical Techniques in Crystallography and Materials Science*. New York: Springer-Verlag.
 Watkin, D. (1994). *Acta Cryst.* **A50**, 411–437.
 Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). *CAMERON*. Chemical Crystallography Laboratory, Oxford, England.

supporting information

Acta Cryst. (2016). E72, 1150-1152 [doi:10.1107/S2056989016010641]

Crystal structure of 2-isopropyl-5,7'-dimethyl-1',3',3a',6',8a',8b'-hexahydro-spiro[cyclohexane-1,6'-furo[3,4-d]imidazo[1,5-b]isoxazol]-8'(7'H)-one

Heithem Abda, Khaireddine Ezzayani, Kaiss Aouadi, Taha Guerfel, Sebastien Vidal and Moncef Msaddek

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO* (Agilent, 2013); data reduction: *CrysAlis PRO* (Agilent, 2013); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *CAMERON* (Watkin *et al.*, 1996); software used to prepare material for publication: *CRYSTALS* (Betteridge *et al.*, 2003).

(I)

Crystal data

$C_{17}H_{28}N_2O_3$

$M_r = 308.42$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.7474$ (6) Å

$b = 11.1404$ (8) Å

$c = 19.208$ (2) Å

$V = 1657.8$ (2) Å³

$Z = 4$

$F(000) = 672$

$D_x = 1.236$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.5418$ Å

Cell parameters from 5548 reflections

$\theta = 4.5$ – 66.7°

$\mu = 0.68$ mm⁻¹

$T = 150$ K

Block, colorless

$0.49 \times 0.43 \times 0.25$ mm

Data collection

Oxford Diffraction Xcalibur (Atlas, Gemini Ultra)

diffractometer

Radiation source: Enhance Ultra (Cu) X-ray source

Mirror monochromator

Detector resolution: 10.4678 pixels mm⁻¹

ω scans

Absorption correction: analytical

[*CrysAlis PRO* (Agilent, 2013) based on expressions derived by Clark & Reid (1995); changes in illuminated volume were kept to a minimum, and were taken into account (Görbitz, 1999)]

$T_{\min} = 0.782$, $T_{\max} = 0.866$

10374 measured reflections

2879 independent reflections

2680 reflections with $I > 2.0\sigma(I)$

$R_{\text{int}} = 0.059$

$\theta_{\max} = 66.8^\circ$, $\theta_{\min} = 11^\circ$

$h = -9 \rightarrow 8$

$k = -13 \rightarrow 12$

$l = -22 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.096$
 $S = 1.03$
 2866 reflections
 201 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Hydrogen site location: difference Fourier map
 H-atom parameters constrained

Method, part 1, Chebychev polynomial,
 (Watkin, 1994; Prince, 1982) [weight] =
 $1.0/[A_0*T_0(x) + A_1*T_1(x) \dots + A_{n-1}*T_{n-1}(x)]$
 where A_i are the Chebychev coefficients listed
 below and $x = F/F_{max}$ Method = Robust
 Weighting (Prince, 1982) $W = [weight]^*$
 $[1-(\Delta F/6*\sigma F)^2]^2$ A_i are: 0.124E + 04
 0.195E + 04 0.105E + 04 304.
 $(\Delta/\sigma)_{max} = 0.0002$
 $\Delta\rho_{max} = 0.16 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{min} = -0.17 \text{ e } \text{Å}^{-3}$
 Extinction correction: Larson (1970), Equation
 22
 Extinction coefficient: 74 (4)
 Absolute structure: Flack (1983), 1261 Friedel
 pairs
 Absolute structure parameter: 0.6 (3)

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems open-flow nitrogen cryostat with a nominal stability of 0.1K.

Refinement. The analytical numeric absorption correction using a multi-faceted crystal model is based on expressions derived by Clark & Reid (1995). The relatively large ratio of minimum to maximum corrections applied in the multi-scan process (1:nnn) reflect changes in the illuminated volume of the crystal. Changes in illuminated volume were kept to a minimum, and were taken into account (Görbitz, 1999).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

| | x | y | z | U_{iso}^*/U_{eq} |
|------|------------|--------------|--------------|--------------------|
| O1 | 0.3924 (2) | 0.46528 (13) | 0.43226 (8) | 0.0295 |
| N2 | 0.2276 (2) | 0.48439 (15) | 0.39468 (9) | 0.0248 |
| C3 | 0.2216 (3) | 0.39172 (18) | 0.33856 (11) | 0.0253 |
| N4 | 0.3084 (3) | 0.44977 (16) | 0.27929 (9) | 0.0257 |
| C5 | 0.3388 (3) | 0.3911 (2) | 0.21315 (11) | 0.0309 |
| H51 | 0.3934 | 0.4467 | 0.1812 | 0.0471* |
| H53 | 0.2321 | 0.3682 | 0.1919 | 0.0466* |
| H52 | 0.4129 | 0.3213 | 0.2193 | 0.0470* |
| C6 | 0.3369 (3) | 0.56800 (19) | 0.28913 (11) | 0.0269 |
| C7 | 0.2626 (3) | 0.60075 (18) | 0.35923 (11) | 0.0257 |
| C8 | 0.3798 (3) | 0.67280 (19) | 0.40726 (12) | 0.0279 |
| C9 | 0.4274 (3) | 0.5792 (2) | 0.46276 (11) | 0.0271 |
| C10 | 0.3137 (4) | 0.6110 (2) | 0.52465 (12) | 0.0372 |
| O11 | 0.1969 (2) | 0.70248 (16) | 0.50158 (9) | 0.0397 |
| C12 | 0.2873 (3) | 0.7679 (2) | 0.44941 (13) | 0.0352 |
| H122 | 0.2094 | 0.8163 | 0.4202 | 0.0417* |
| H121 | 0.3689 | 0.8236 | 0.4714 | 0.0416* |
| H102 | 0.2495 | 0.5396 | 0.5410 | 0.0447* |
| H101 | 0.3821 | 0.6433 | 0.5635 | 0.0448* |
| H91 | 0.5499 | 0.5839 | 0.4753 | 0.0337* |
| H81 | 0.4758 | 0.7027 | 0.3812 | 0.0341* |

| | | | | |
|------|-------------|--------------|--------------|---------|
| H71 | 0.1551 | 0.6415 | 0.3518 | 0.0307* |
| O13 | 0.4052 (2) | 0.63710 (15) | 0.24780 (8) | 0.0357 |
| C14 | 0.0307 (3) | 0.3616 (2) | 0.32135 (11) | 0.0274 |
| C15 | -0.0806 (3) | 0.4686 (2) | 0.29604 (12) | 0.0310 |
| C16 | -0.2129 (4) | 0.4250 (3) | 0.24307 (14) | 0.0435 |
| H162 | -0.2845 | 0.4937 | 0.2288 | 0.0654* |
| H163 | -0.1594 | 0.3903 | 0.2017 | 0.0653* |
| H161 | -0.2850 | 0.3629 | 0.2652 | 0.0658* |
| C17 | -0.1731 (4) | 0.5369 (2) | 0.35403 (14) | 0.0406 |
| H171 | -0.2186 | 0.6122 | 0.3336 | 0.0606* |
| H173 | -0.2672 | 0.4908 | 0.3721 | 0.0604* |
| H172 | -0.0979 | 0.5542 | 0.3931 | 0.0602* |
| H151 | -0.0031 | 0.5263 | 0.2711 | 0.0372* |
| C18 | -0.0536 (3) | 0.2924 (2) | 0.38122 (13) | 0.0346 |
| C19 | 0.0460 (4) | 0.1795 (2) | 0.40142 (14) | 0.0379 |
| C20 | 0.2339 (3) | 0.2100 (2) | 0.41916 (12) | 0.0327 |
| C21 | 0.3154 (3) | 0.27553 (19) | 0.35817 (12) | 0.0288 |
| H211 | 0.3096 | 0.2212 | 0.3171 | 0.0339* |
| H212 | 0.4342 | 0.2949 | 0.3684 | 0.0350* |
| C22 | 0.3387 (4) | 0.0976 (2) | 0.43654 (13) | 0.0429 |
| H222 | 0.4583 | 0.1173 | 0.4469 | 0.0645* |
| H221 | 0.3358 | 0.0416 | 0.3963 | 0.0629* |
| H223 | 0.2912 | 0.0549 | 0.4787 | 0.0629* |
| H201 | 0.2358 | 0.2644 | 0.4599 | 0.0385* |
| H192 | 0.0469 | 0.1237 | 0.3607 | 0.0463* |
| H191 | -0.0113 | 0.1390 | 0.4418 | 0.0451* |
| H182 | -0.1724 | 0.2721 | 0.3658 | 0.0423* |
| H181 | -0.0606 | 0.3469 | 0.4217 | 0.0410* |
| H141 | 0.0390 | 0.3055 | 0.2810 | 0.0325* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| O1 | 0.0344 (8) | 0.0232 (7) | 0.0308 (8) | 0.0026 (7) | -0.0085 (7) | -0.0014 (6) |
| N2 | 0.0304 (10) | 0.0202 (9) | 0.0239 (9) | 0.0002 (8) | -0.0028 (8) | 0.0005 (7) |
| C3 | 0.0324 (12) | 0.0197 (10) | 0.0237 (10) | 0.0008 (9) | 0.0029 (9) | -0.0011 (9) |
| N4 | 0.0317 (9) | 0.0218 (9) | 0.0236 (9) | -0.0001 (8) | 0.0035 (8) | -0.0015 (7) |
| C5 | 0.0377 (12) | 0.0307 (11) | 0.0242 (11) | -0.0039 (10) | 0.0039 (9) | -0.0034 (9) |
| C6 | 0.0290 (11) | 0.0233 (10) | 0.0284 (11) | -0.0006 (9) | -0.0018 (9) | 0.0016 (9) |
| C7 | 0.0302 (11) | 0.0206 (11) | 0.0262 (10) | 0.0023 (10) | 0.0024 (9) | 0.0014 (8) |
| C8 | 0.0324 (12) | 0.0204 (10) | 0.0307 (12) | -0.0016 (9) | 0.0028 (10) | -0.0014 (9) |
| C9 | 0.0313 (11) | 0.0233 (10) | 0.0267 (10) | 0.0000 (9) | -0.0042 (9) | -0.0040 (9) |
| C10 | 0.0453 (14) | 0.0361 (13) | 0.0302 (12) | -0.0014 (12) | -0.0002 (11) | -0.0035 (10) |
| O11 | 0.0392 (10) | 0.0387 (9) | 0.0413 (9) | 0.0061 (8) | 0.0076 (8) | -0.0078 (8) |
| C12 | 0.0421 (15) | 0.0257 (11) | 0.0377 (13) | 0.0022 (11) | -0.0016 (11) | -0.0076 (10) |
| O13 | 0.0489 (10) | 0.0280 (8) | 0.0303 (8) | -0.0062 (8) | 0.0057 (8) | 0.0044 (6) |
| C14 | 0.0329 (11) | 0.0228 (11) | 0.0265 (11) | -0.0031 (9) | 0.0021 (9) | -0.0005 (9) |
| C15 | 0.0295 (11) | 0.0298 (12) | 0.0335 (12) | -0.0022 (10) | -0.0007 (10) | 0.0024 (9) |

| | | | | | | |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| C16 | 0.0430 (14) | 0.0423 (14) | 0.0453 (14) | -0.0056 (13) | -0.0109 (13) | 0.0027 (12) |
| C17 | 0.0370 (13) | 0.0368 (13) | 0.0481 (15) | 0.0038 (12) | 0.0041 (12) | -0.0022 (11) |
| C18 | 0.0371 (13) | 0.0322 (13) | 0.0344 (12) | -0.0047 (11) | 0.0039 (11) | 0.0025 (10) |
| C19 | 0.0533 (16) | 0.0238 (12) | 0.0365 (13) | -0.0085 (11) | 0.0037 (12) | 0.0039 (10) |
| C20 | 0.0495 (15) | 0.0209 (11) | 0.0278 (11) | -0.0015 (11) | -0.0011 (11) | -0.0001 (9) |
| C21 | 0.0360 (12) | 0.0211 (11) | 0.0292 (11) | 0.0013 (10) | 0.0009 (10) | -0.0008 (9) |
| C22 | 0.0680 (19) | 0.0239 (12) | 0.0369 (13) | 0.0025 (12) | -0.0068 (13) | 0.0040 (10) |

Geometric parameters (Å, °)

| | | | |
|------------|-------------|--------------|-------------|
| O1—N2 | 1.482 (2) | C14—C15 | 1.550 (3) |
| O1—C9 | 1.424 (3) | C14—C18 | 1.531 (3) |
| N2—C3 | 1.493 (3) | C14—H141 | 0.997 |
| N2—C7 | 1.489 (3) | C15—C16 | 1.524 (3) |
| C3—N4 | 1.472 (3) | C15—C17 | 1.527 (3) |
| C3—C14 | 1.552 (3) | C15—H151 | 1.001 |
| C3—C21 | 1.531 (3) | C16—H162 | 0.983 |
| N4—C5 | 1.448 (3) | C16—H163 | 0.977 |
| N4—C6 | 1.349 (3) | C16—H161 | 0.986 |
| C5—H51 | 0.969 | C17—H171 | 0.991 |
| C5—H53 | 0.956 | C17—H173 | 0.957 |
| C5—H52 | 0.974 | C17—H172 | 0.970 |
| C6—C7 | 1.509 (3) | C18—C19 | 1.526 (4) |
| C6—O13 | 1.226 (3) | C18—H182 | 0.992 |
| C7—C8 | 1.523 (3) | C18—H181 | 0.988 |
| C7—H71 | 0.959 | C19—C20 | 1.533 (4) |
| C8—C9 | 1.536 (3) | C19—H192 | 0.999 |
| C8—C12 | 1.513 (3) | C19—H191 | 1.001 |
| C8—H81 | 0.957 | C20—C21 | 1.518 (3) |
| C9—C10 | 1.521 (3) | C20—C22 | 1.529 (3) |
| C9—H91 | 0.980 | C20—H201 | 0.990 |
| C10—O11 | 1.433 (3) | C21—H211 | 0.996 |
| C10—H102 | 0.989 | C21—H212 | 0.966 |
| C10—H101 | 0.983 | C22—H222 | 0.972 |
| O11—C12 | 1.423 (3) | C22—H221 | 0.995 |
| C12—H122 | 0.985 | C22—H223 | 1.009 |
| C12—H121 | 0.981 | | |
| N2—O1—C9 | 103.68 (14) | C3—C14—C18 | 110.82 (19) |
| O1—N2—C3 | 106.18 (15) | C15—C14—C18 | 112.69 (19) |
| O1—N2—C7 | 100.99 (15) | C3—C14—H141 | 103.9 |
| C3—N2—C7 | 106.11 (15) | C15—C14—H141 | 105.9 |
| N2—C3—N4 | 103.92 (16) | C18—C14—H141 | 107.2 |
| N2—C3—C14 | 109.41 (17) | C14—C15—C16 | 109.79 (19) |
| N4—C3—C14 | 111.43 (18) | C14—C15—C17 | 114.5 (2) |
| N2—C3—C21 | 113.10 (17) | C16—C15—C17 | 109.3 (2) |
| N4—C3—C21 | 110.18 (18) | C14—C15—H151 | 108.1 |
| C14—C3—C21 | 108.77 (18) | C16—C15—H151 | 106.8 |

| | | | |
|---------------|-------------|----------------|-------------|
| C3—N4—C5 | 123.70 (17) | C17—C15—H151 | 108.1 |
| C3—N4—C6 | 113.29 (17) | C15—C16—H162 | 108.5 |
| C5—N4—C6 | 122.47 (19) | C15—C16—H163 | 112.6 |
| N4—C5—H51 | 109.8 | H162—C16—H163 | 108.7 |
| N4—C5—H53 | 110.8 | C15—C16—H161 | 108.4 |
| H51—C5—H53 | 106.1 | H162—C16—H161 | 110.3 |
| N4—C5—H52 | 110.4 | H163—C16—H161 | 108.3 |
| H51—C5—H52 | 109.2 | C15—C17—H171 | 107.4 |
| H53—C5—H52 | 110.4 | C15—C17—H173 | 110.8 |
| N4—C6—C7 | 107.39 (18) | H171—C17—H173 | 109.1 |
| N4—C6—O13 | 126.4 (2) | C15—C17—H172 | 112.4 |
| C7—C6—O13 | 126.2 (2) | H171—C17—H172 | 110.5 |
| C6—C7—N2 | 105.48 (16) | H173—C17—H172 | 106.5 |
| C6—C7—C8 | 116.15 (19) | C14—C18—C19 | 113.0 (2) |
| N2—C7—C8 | 106.89 (17) | C14—C18—H182 | 106.7 |
| C6—C7—H71 | 108.3 | C19—C18—H182 | 110.9 |
| N2—C7—H71 | 108.7 | C14—C18—H181 | 107.8 |
| C8—C7—H71 | 111.0 | C19—C18—H181 | 109.5 |
| C7—C8—C9 | 101.88 (16) | H182—C18—H181 | 108.9 |
| C7—C8—C12 | 114.2 (2) | C18—C19—C20 | 110.74 (19) |
| C9—C8—C12 | 102.57 (18) | C18—C19—H192 | 108.4 |
| C7—C8—H81 | 109.3 | C20—C19—H192 | 107.8 |
| C9—C8—H81 | 114.4 | C18—C19—H191 | 110.1 |
| C12—C8—H81 | 113.9 | C20—C19—H191 | 110.4 |
| C8—C9—O1 | 105.88 (16) | H192—C19—H191 | 109.2 |
| C8—C9—C10 | 104.18 (18) | C19—C20—C21 | 109.3 (2) |
| O1—C9—C10 | 114.75 (19) | C19—C20—C22 | 111.8 (2) |
| C8—C9—H91 | 111.5 | C21—C20—C22 | 110.0 (2) |
| O1—C9—H91 | 109.5 | C19—C20—H201 | 109.0 |
| C10—C9—H91 | 110.8 | C21—C20—H201 | 108.1 |
| C9—C10—O11 | 106.84 (18) | C22—C20—H201 | 108.7 |
| C9—C10—H102 | 110.7 | C3—C21—C20 | 113.52 (19) |
| O11—C10—H102 | 110.6 | C3—C21—H211 | 107.3 |
| C9—C10—H101 | 111.5 | C20—C21—H211 | 107.5 |
| O11—C10—H101 | 108.3 | C3—C21—H212 | 108.2 |
| H102—C10—H101 | 108.9 | C20—C21—H212 | 110.3 |
| C10—O11—C12 | 105.74 (19) | H211—C21—H212 | 109.9 |
| C8—C12—O11 | 104.57 (19) | C20—C22—H222 | 111.5 |
| C8—C12—H122 | 111.7 | C20—C22—H221 | 109.4 |
| O11—C12—H122 | 112.4 | H222—C22—H221 | 108.8 |
| C8—C12—H121 | 111.6 | C20—C22—H223 | 111.6 |
| O11—C12—H121 | 109.7 | H222—C22—H223 | 106.8 |
| H122—C12—H121 | 107.0 | H221—C22—H223 | 108.7 |
| C3—C14—C15 | 115.54 (18) | | |
| C9—O1—N2—C3 | 156.12 (16) | C21—C3—C14—C18 | 53.8 (2) |
| C9—O1—N2—C7 | 45.58 (18) | N2—C3—C21—C20 | 64.1 (2) |
| N2—O1—C9—C8 | -40.8 (2) | N4—C3—C21—C20 | 179.91 (18) |

| | | | |
|----------------|--------------|-----------------|--------------|
| N2—O1—C9—C10 | 73.5 (2) | C14—C3—C21—C20 | -57.7 (2) |
| C12—O11—C10—C9 | -30.4 (2) | O13—C6—C7—N2 | -169.1 (2) |
| C10—O11—C12—C8 | 41.4 (2) | O13—C6—C7—C8 | -51.0 (3) |
| O1—N2—C3—N4 | -88.72 (18) | N4—C6—C7—N2 | 13.2 (2) |
| O1—N2—C3—C14 | 152.15 (15) | N4—C6—C7—C8 | 131.3 (2) |
| O1—N2—C3—C21 | 30.7 (2) | N2—C7—C8—C9 | 9.3 (2) |
| C7—N2—C3—N4 | 18.2 (2) | N2—C7—C8—C12 | -100.5 (2) |
| C7—N2—C3—C14 | -100.94 (18) | C6—C7—C8—C9 | -108.1 (2) |
| C7—N2—C3—C21 | 137.64 (18) | C6—C7—C8—C12 | 142.1 (2) |
| O1—N2—C7—C6 | 91.23 (18) | C7—C8—C9—O1 | 19.2 (2) |
| O1—N2—C7—C8 | -32.94 (19) | C7—C8—C9—C10 | -102.2 (2) |
| C3—N2—C7—C6 | -19.4 (2) | C12—C8—C9—O1 | 137.68 (18) |
| C3—N2—C7—C8 | -143.56 (17) | C12—C8—C9—C10 | 16.3 (2) |
| C5—N4—C3—N2 | 177.7 (2) | C7—C8—C12—O11 | 74.2 (2) |
| C5—N4—C3—C14 | -64.6 (3) | C9—C8—C12—O11 | -35.1 (2) |
| C5—N4—C3—C21 | 56.2 (3) | O1—C9—C10—O11 | -107.8 (2) |
| C6—N4—C3—N2 | -10.6 (3) | C8—C9—C10—O11 | 7.6 (2) |
| C6—N4—C3—C14 | 107.2 (2) | C3—C14—C15—C16 | 146.3 (2) |
| C6—N4—C3—C21 | -132.0 (2) | C3—C14—C15—C17 | -90.4 (2) |
| C3—N4—C6—O13 | -179.3 (2) | C18—C14—C15—C16 | -84.9 (2) |
| C3—N4—C6—C7 | -1.6 (3) | C18—C14—C15—C17 | 38.4 (3) |
| C5—N4—C6—O13 | -7.4 (4) | C3—C14—C18—C19 | -54.8 (3) |
| C5—N4—C6—C7 | 170.3 (2) | C15—C14—C18—C19 | 174.06 (19) |
| N2—C3—C14—C15 | 59.5 (2) | C14—C18—C19—C20 | 55.5 (3) |
| N2—C3—C14—C18 | -70.2 (2) | C18—C19—C20—C21 | -55.6 (3) |
| N4—C3—C14—C15 | -54.8 (2) | C18—C19—C20—C22 | -177.5 (2) |
| N4—C3—C14—C18 | 175.48 (17) | C19—C20—C21—C3 | 58.7 (2) |
| C21—C3—C14—C15 | -176.47 (18) | C22—C20—C21—C3 | -178.24 (19) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|----------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| C5—H52 \cdots O13 ⁱ | 0.97 | 2.57 | 3.536 (3) | 172 |

Symmetry code: (i) $-x+1, y-1/2, -z+1/2$.