

DOI: 10.1002/open.201300033

Metallocyclodextrins: Combining Cavitands with Metal Centres

Rafael Gramage-Doria^{*[a]}

Awarding Institution: Université de Strasbourg (France)

Date Awarded: January 3, 2012

Supervisors: Dr. Dominique Matt and Prof. Dr. Dominique Armspach, Université de Strasbourg (France)

In the last decades, combining the properties of a transition-metal ion with those of a covalently linked, molecular cavity has attracted a great deal of attention. Numerous studies focused on the design and synthesis of systems exploiting the binding properties of a receptor moiety attached to an active metal site, with the aim of mimicking nature's metal-containing catalysts: metalloenzymes. Other potential applications involve (1) the study of metal-promoted reactions occurring in a confined environment in order to find out whether this strategy leads to selective transformations, and (2) molecular recognition via supramolecular interactions in conjunction with photo- or electrochemical applications. In this respect, cyclodextrins (CDs) and their chemically modified derivatives occupy a position of choice. Their rigid, conical structure, inherent chirality, as well as the presence of hydroxy groups that can be substituted in a regioselective manner, offer many possibilities in terms of cavitand (cavity-shaped ligand) design. So far, most studies have dealt with the smallest member of the family, namely α -CD, because of the ease with which it can be functionalised, β -CD derivatives remaining largely unexplored, even if their bigger inner space seems more appropriate for taking full advantage of the cavity. The present thesis is mainly concerned with the preparation of new *introverted* phosphanes (**WIDEPHOS** and **L**, Figure 1) derived from multifunctionalised methylated β -CDs and the study of their coordination and catalytic properties. The last part of the manuscript focuses on the use of




Figure 1. Cavtands designed and studied in this work.

[a] Dr. R. Gramage-Doria
Van't Hoff Institute for Molecular Sciences, University of Amsterdam
Science Park 904, 1098 XH, Amsterdam (The Netherlands)
E-mail: r.gramagedoria@uva.nl



The full thesis is available as Supporting Information on the WWW under <http://dx.doi.org/10.1002/open.201300033>.

© 2013 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

a smaller α -CD analogue of the aforementioned phosphanes, namely the *trans*-chelating diphosphane **TRANSDIP** (Figure 1) in palladium-promoted carbon–carbon bond-forming reactions.

The first part of this thesis focuses on reviewing transition-metal-based cavitands, for which the first and second metal coordination spheres are controlled by their cavity-shaped ligand. In the next part, four different regioselective double capping reactions were performed on α - and β -CD scaffolds. More precisely, the use of a rigid and bulky dialkylating reagent, containing two trityl-like moieties, gave access to an A,B,D,E-tetrafunctionalised β -CD regioisomer **1**, which is the key compound for preparing different types of cavitands bearing various functionalities, such as, disulfide **2**, diphosphane **WIDEPHOS** and disulfate **3** (Figure 2). A disulfate based on smaller α -CD (**4**, Figure 2) and introverted mono-phosphanes derived from β -CD were also reported. It is worth noting that disulfate-capped species **3** and **4** might be considered as potential precursors for the preparation of hetero-trifunctionalised cavitands. The capping of adjacent glucose units in CD-derived compounds turned out to be an extremely powerful strategy to straightforwardly prepare a library of cavitands.

The following chapters are devoted to study the coordination and catalytic properties of two new phosphane ligands built on a large β -CD scaffold: **WIDEPHOS** and **L. WIDEPHOS** bears two phosphorus lone pairs pointing to the cavity interior without being aligned because of the unsymmetrical nature of the β -CD scaffold. This geometrical feature, combined with the large distance separating the two phosphorus atoms (P...P distance of 6.91 Å in its corresponding phosphine oxide), promote the formation of “imperfect” *trans*-chelate complexes in which the metal centre swings about the ligand according to detailed NMR studies. This unprecedented molecular movement, named *oschelation*, allows each phosphorus atom to form an optimal bond in turn with the coordinated d^8 (Pd^{II}, Pt^{II}, Rh^I)




Figure 2. Different cavitands prepared from key compound **1**, showing the X-ray structure of **4** and adduct **WIDEPHOS·(BH₃)₂**.





Figure 3. Balance-wheel movement of **WIDEPHOS** (top), reactivity of an *oschelate* palladium complex (middle) and rare dinuclear species derived from **WIDEPHOS** (bottom).

or d¹⁰ (Au^I) transition metal ions (Figure 3). This type of motion contrasts with the well-studied ligand hemilability, since no metal-phosphorus dissociation is observed in this case. Evidence for the weakness of one of the M–P bonds came from the facile preparation of unique palladium species **5** and **6** (Figure 3), in which one of the M–P bonds have been broken and non-conventional monophosphanes and P,O-chelating complexes form. Oschelation can be stopped by addition of a second metal centre to the cavitand. Indeed, further studies on **WIDEPHOS** proved that it is better suited for coordinating dinuclear fragments within the confinement of the large β-CD cavity. However, severe steric constraints on the metal first sphere of coordination result in the formation of unprecedented, single μ-chlorido-bridged dinuclear species; additionally stabilized by Cl...H hydrogen bonding between the intracavity-located chlorido ligands and inwardly oriented CD protons called H-5. In this new type of square-planar complexes, non-optimal orbital overlapping measured by the so-called tilt angle was also found to take place for one of the phosphorus atoms together with an oschelation movement involving non-identical donor atoms, namely a phosphorus and an oxygen atom (Figure 3). Static gold(I) dinuclear complexes displaying similar imperfect orbital overlapping for one of the phosphorus atom were also prepared (Figure 3). **WIDEPHOS** was active in some palladium-catalysed reactions, although its performance was found to be poorer than **L** (Figure 1).



Scheme 1. Following a C–C bond forming reaction on **TRANSDIP**.

The last part of the manuscript concerns the unexpected C–C coupling reaction (ethane formation) mediated by a palladium *trans*-chelate complex derived from the previously reported diphospane **TRANSDIP**. Several key intermediates involved in the mechanism of this C–C bond formation were isolated and identified (Scheme 1), notably the unprecedented $\kappa^1\text{-P}$ -[PdMe₃(**TRANSDIP**)]Li complex (**9**). The confinement imposed by **TRANSDIP** on the palladium centre enabled direct visualisation of three types of highly reactive intermediates, in particular, of the first pentacoordinated alkyl complex containing the *trans*-spanning diphospane **7**. In contrast to analogues based on other *trans*-chelating diphosphanes, the intermediate [PdMeCl-(**TRANSDIP**)] (**8**) undergoes dissociation of one of its phosphorus atoms upon nucleophilic attack of MeLi, re-coordination occurring after the reductive elimination step. Remarkably, the protecting role of the cavity towards M–Cl bonds is essential for stabilising a zerovalent palladium atom in one of the complexes (**10**). Finally, the perfect match between a [Pd($\eta^2\text{-O}_2$)] moiety and the cyclodextrin core makes the [Pd($\eta^2\text{-O}_2$)(**TRANSDIP**)] complex (**11**) unusually stable compared with previously reported peroxo complexes.

Overall, the work described here emphasizes the importance of ligand design in building metallo-receptors derived from cyclodextrins. These new cavity-shaped molecules allowed the study of unprecedented dynamic coordination processes occurring in chelate complexes (*oschelation*), which proved to have a profound effect on metal reactivity. **WIDEPHOS**, with its two intra-annular, but wide apart donor atoms, constitutes a very peculiar cavitand allowing the formation of both mono and dinuclear complexes with extremely distorted coordination geometries. In particular, this study revealed that supramolecular interactions between a metal fragment and the cyclodextrin inner wall play a fundamental role in the stabilisation of otherwise unstable species. Lastly, we have shown the application of introverted phosphanes in homogeneous catalysis and the use of metallocavitands based on an authentic *trans*-spanning ligand (**TRANSDIP**) for unravelling, at that time unknown, mechanistic pathways associated with a palladium-promoted C–C bond formation. At this stage, one can anticipate that metallocavitands will create many opportunities regarding the catalytic outcome of reactions, in particular, in terms of selectivity that may result from the supramolecular confinement of non-conventional and otherwise unstable species and/or the activation of small molecules such as dioxygen, amongst others.

Keywords: cavitands · C–C bond formation · cyclodextrins · oschelation · phosphanes

Publications arising from this work:

- R. Gramage-Doria, D. Armspach, D. Matt, L. Toupet, *Angew. Chem. Int. Ed.* **2011**, *50*, 1554–11559; *Angew. Chem.* **2011**, *123*, 1592–1597.
- R. Gramage-Doria, D. Rodriguez-Lucena, D. Armspach, C. Egloff, M. Jouffroy, D. Matt, L. Toupet, *Chem. Eur. J.* **2011**, *17*, 3911–3921.
- R. Gramage-Doria, D. Armspach, D. Matt, L. Toupet, *Dalton Trans.* **2012**, *41*, 8786–8796.
- R. Gramage-Doria, D. Armspach, D. Matt, L. Toupet, *Chem. Eur. J.* **2012**, *18*, 10813–10816.
- R. Gramage-Doria, D. Armspach, D. Matt, *Coord. Chem. Rev.* **2013**, *257*, 776–816.

Published online on August 28, 2013