

Original research

Open Access

Diagnostic performance and predictive value of rheumatoid factor, anti-cyclic-citrullinated peptide antibodies and HLA-DRB1 locus genes in rheumatoid arthritis

Nihal A Fathi¹, Azza M Ezz-Eldin², Eman Mosad^{*3}, Rania M Bakry³, Hosny B Hamed³, Sahar Ahmed¹, Marwa Mahmoud¹, Hebat-Allah G Rashed² and Fatma Abdullah¹

Address: ¹Rheumatology department, Assiut University Hospital, Assiut, Egypt, ²Clinical pathology department, Assiut University Hospital, Assiut, Egypt and ³HLA Unit, Clinical pathology department, South Egypt Cancer Institute, Assiut University, Assiut, Egypt

Email: Nihal A Fathi - nfathy@yahoo.com; Azza M Ezz-Eldin - azzam80@hotmail.com; Eman Mosad* - eman_mosaad@hotmail.com; Rania M Bakry - rania.bakry@lycos.com; Hosny B Hamed - badrawyh@yahoo.com; Sahar Ahmed - saharsaad68@hotmail.com; Marwa Mahmoud - hha_79@hotmail.com; Hebat-Allah G Rashed - h_h_70@yahoo.com; Fatma Abdullah - fatma_abdallah@rocketmail.com

* Corresponding author

Published: 22 October 2008

Received: 26 August 2008

International Archives of Medicine 2008, 1:20 doi:10.1186/1755-7682-1-20

Accepted: 22 October 2008

This article is available from: <http://www.intarchmed.com/content/1/1/20>

© 2008 Fathi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: We evaluated the significance of the genes, defined as *DRB1*04* or *DRB1*01*, in rheumatoid arthritis (RA) patients. We focused on the role of genetic and serologic markers to predict disease activity and destructive process of joints.

Methods: Sixty patients with RA were examined. Radiographic changes were evaluated by (Larsen score) and disease activity was measured by disease activity score 28 (DAS28). The markers analyzed were: erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), rheumatoid factor (RF), anti-cyclic citrullinated peptides (anti-CCP2) and HLA-*DRB1* alleles typed by PCR.

Results: In this study, anti-CCP antibodies, CRP, RF and AKA were detected in 83.3%, 56.7%, 71.7% and 52% of patients respectively. HLA-*DRB1*01* was found in 45% of patients and 35% of them had one or two HLA-*DRB1*04* alleles. According to *DRB1*04* subtypes, (*DRB1*0405*) was present in of 80% them. For prediction of grade of activity, the independent predictors were anti-CCP (OR 19.6), and *DRB1*04* positive allele (OR 5.1). The combination of *DRB1*04* + anti-CCP antibodies gave increase in the specificity and positive predictive value to 92% and 90 respectively. As regards to the prediction of radiological joint damage, the independent predictors were HLA-*DRB1*04*, HLA-*DRB1*01*, RF, and CRP > 18 (OR 5.5, 4.5, 2.5, 2.0 respectively).

Conclusion: Our findings suggest that anti-CCP2 is superior to RF for the detection of RA and provided predictive information on joint destruction and disease activity. The presence of RA associated antibodies (ACCP or RF) and/or the SE genes are indicative for a poorer radiological outcome and higher grade of activity.

Introduction

Rheumatoid arthritis (RA) is an inflammatory disease of unknown cause. The course of rheumatoid arthritis is ranging from mild to aggressive forms, the latter being very difficult to cope with. It has been shown that early diagnosis and treatment reduce joint destruction, and improve survival [1]. Risk factors have been identified in groups of patients with different outcomes such as baseline radiographic joint changes, presence of rheumatoid factor (RF), specific human leukocyte antigens (HLA); HLA-*DRB1* genotypes, high disease activity, high disability scores, and high levels of acute phase proteins [2]. Antibodies to cyclic citrullinated peptide (anti-CCP) have been documented extensively over recent years as highly specific serological markers for rheumatoid arthritis, with important clinical implications for diagnosis and prognosis [3].

It has been suggested that the etiology of RA is due to an interaction between genetic and environmental factors. Genetic studies have demonstrated multiple HLA-*DRB1* alleles encoding a conserved sequence at amino acid positions 70–74 are associated with susceptibility and severity of RA. This conserved sequence is commonly known as the shared epitope (SE). The role of the SE in the evolution of articular destruction has yielded conflicting results [4,5].

The present study is a cross-sectional analysis aimed to evaluate the significance of the presence of SE genes, defined as *DRB1**01 or *DRB1**04, in relation to anti-CCP antibodies, antikeratin antibody (AKA) and RF in individuals who developed RA. We focused on disease activity and joint damage, evaluated on radiographs, as outcome variables.

Methods

This study was carried out on 60 outpatients who fulfilled the American college of rheumatology criteria for RA. A written consent was obtained from the patients according to the Declaration of Helsinki prior to enrollment in the study. The study was approved by the Assiut University local ethical committee. Patients were subsequently treated with disease modifying antirheumatic drugs (usually methotrexate, sulphasalazine, or a combination of both). The patients were evaluated for age; sex; body mass index; disease duration; duration of morning stiffness; patients' assessment of pain (on a visual analogue scale); number of swollen and tender joints, disease activity score; presence or absence of nodules and extra-articular manifestations. Disease activity by the disease activity score (DAS28) [6]. Global health and pain and visual analogue scale were assessed. Functional disability was evaluated using health assessment questionnaire [7]. Hand,

wrist, and foot radiographs were obtained, evaluated and scored using Larsen method [8].

Sample handling and investigations

Blood samples were collected; for complete blood count, erythrocyte sedimentation rate (ESR), C-reactive protein (CRP) and DNA isolation. One hundred samples were taken from healthy blood donors as a control for HLA typing; twenty of them were used for serologic markers.

Determination of antibodies

RF was detected by the kit supplied Biotec Laboratories Lot No. RF-460 based on agglutination test using particles sensitized with human IgG. ANA was detected by the Fluro-kit supplies by supplied by DiaSorine Catalog No 1740 based on indirect immunofluorescent for the screening and titration of antinuclear antibodies (ANA). Anti-CCP2 antibodies were detected by enzyme-linked immunosorbent assay Kit supplies by INOVA Diagnostica Cat. NO 570139. Serum samples with a test result 25 U/ml were considered positive, and designated as the "standard" cut off. AKA was detected by kit supplied by IMMECO Diagnostic, Lot No. 2120-8, based on indirect immunofluorescence antibodies on rat esophagus substrate.

HLA-*DRB1* genotyping

HLA typing was performed using sequence specific oligonucleotide technique, reversed dot blot hybridization technique (DynaL RELI HLA-*DRB1*, product No.802.01) [9]. DNA for HLA-*DRB1* typing of patients was isolated from peripheral blood mononuclear cells by using the DNA Isolation Kit (QIAamp minikit, Qiagen, Hilden, Germany). The biotinylated primers anneal to their target, Tag DNA polymerase extend the annealed primers along the target templates to produce a biotinylated DNA sequence (amplicon). After polymerase chain reaction amplification, the amplicons are chemically denatured, these are added to nylon membrane that contains an array of immobilized sequence specific oligonucleotide probes that contain a complementary target sequence and thus are captured onto the membrane strip. The amplicon-probe complex is visualized using a colorimetric reaction. The resulting probe signals are captured to the control probe intensity and the samples recoded for interpretation [10]. The specificity of the amplification can be locus specific (eg, HLA-A, HLA-B, HLA-*DRB1*) group specific (eg, *DRB1**01, *DRB1**02), or allele-specific (eg, *DRB1*-0404, *DRB1*-0405). According to HLA-*DRB1**04 subtyping the shared epitopes encoding alleles *0409, *0413, *0416 and *0421 were classified as *0401 alleles, *0408, *0410 and *0419 were classified as *0404. The *DRB1**0405 alleles were analyzed as a separate entity. All other *DRB1**04 alleles were classified as *DRB1**04 non-SE alleles [4].

Statistical analysis

Statistical analyses were done using SPSS 15 statistical software (SPSS Inc, Chicago, Illinois, USA). Continuous variables were transformed into categorical variables using the median value as the cut off point. Comparison of categorical variables between groups, as well as univariate analysis of the relation between all baseline values and the outcome measure was undertaken using the χ^2 test, with Fisher's exact correction when appropriate. Sensitivity and specificity, and negative (NPV) and positive (PPV) predictive values of the possible predictors were calculated. Significant univariate variables were included in a multivariate stepwise logistic regression model to find relevant independent prognostic variables. P value < 0.05 is considered significant.

Results

Clinical and laboratory investigations were illustrated in Table 1. The median age of the patients was 50 years and 90% were women. According the positivity of anti-CCP antibodies, 50 (83.3%) patients were anti-CCP antibodies positive. CRP, RF and AKA were detected in 34 (56.7%), 43 (71.7%) and 31 (52%) of patients respectively.

Comparing anti-CCP positive with anti-CCP negative patients (Table 1); anti-CCP positive patients had a significantly higher frequency of high ESR, RF, ANA and AKA ($P = 0.04, 0.02, 0.03$ and 0.05) respectively.

The frequencies of HLA-DRB1 alleles in patients and control are shown in Table 2. There was a statistically significant increase in the frequency in DRB1*01, DRB1*02, DRB1*04 compared to control ($P = 0.001, 0.05$ and 0.05 respectively). In this study 45% had HLA-DRB1*01 and 35% of patients had one or two HLA-DRB1*04 alleles.

It was found that RF associated allele (DRB1*0405) was present in of 80% of DRB1*04 positive patients. "Double dose" of DRB1*04 (homozygous) was found in 8/60 (13%) of our patients versus 3/100 (3%) in the control $P = 0.05$ (Data is not shown in the table).

Predictors of the disease activity grade

Univariate predictors of the grade of disease activity were summarized in Table 3. Anti-CCP had the highest sensitivity; 97%, and NPV 90%. On multivariate analysis, the independent predictors of disease activity were anti-CCP (OR 19.6), and DRB1*04 positive allele (OR 5.1). (Table

Table 1: Clinical and laboratory characteristics of patients

		All patients	Anti-CCP		P value
			Negative	Positive	
Age	<50*	45 (75%)	8 (17.8%)	37 (82.2%)	NS
	≥50	15 (25%)	2 (13.3%)	13 (86.7%)	
Sex	female	54 (90%)	9 (16.7%)	45 (83.3%)	NS
	male	6 (10%)	1 (16.7%)	5 (83.3%)	
Duration	<6m	3 (5%)	1 (33.3%)	2 (66.7%)	NS
	≥6m	57 (95%)	9 (15.8%)	48 (84.2%)	
Smoking	yes	5 (8.3%)	1 (20%)	4 (80%)	NS
	no	55 (91.7%)	9 (16.4%)	46 (83.6%)	
CRP	<18 *	26 (43.3%)	7 (26.9%)	19 (73.1%)	NS
	≥18	34 (56.7%)	3 (8.8%)	31 (91.2%)	
RF	+ve	43 (71.7%)	4 (9.3%)	39 (90.7%)	0.02
	-ve		6 (35%)	11 (65%)	
ESR-I	<32 *	14 (23.3%)	5 (35.7%)	9 (64.3%)	0.04
	≥32	46 (76.7%)	5 (10.9%)	41 (89.1%)	
ANA	+ve	22 (36.7%)	0 (0%)	22 (100%)	0.03
	-ve	38 (63.3%)	10 (40.8%)	28 (59.2%)	
AKA	+ve	31 (52%)	3 (10%)	28 (90%)	0.05
	-ve	29 (48%)	17 (59%)	12 (41%)	
Grade of activity (DAS)	- Mild or moderate	25 (41.7%)	9 (31.6%)	16 (68.4%)	0.001
	- Severe	35 (58.3%)	1 (9.8%)	34 (90.2%)	
Larsen score	<35*	19 (31.7%)	6 (10%)	13 (21.7%)	0.05
	≥35	41 (68.3%)	4 (6.7%)	37 (61.7%)	
HLA-DRB1*04	+ve	21 (35%)	2 (9.5%)	19 (90.5%)	NS
	-ve	39 (65%)	8 (20.5%)	31 (79.5%)	
HLA-RB1*01	+ve	27 (45%)	5 (19%)	22 (81%)	NS
	-ve	33 (55%)	5 (15%)	28 (85%)	

Values are expressed as n (%). AKA: antikeratin antibody; anti-CCP: anti-cyclic citrullinated peptides antibodies; ANA: antinuclear antibodies; DAS: disease activity score; CRP: C-reactive protein; RF: rheumatoid factor; HLA: human leukocyte antigen. * Cut-off values are medians.

Table 2: Frequencies of HLA-DRB1 alleles in patients and control

HLA-DRB1 allele	Patients	Control	P value
HLA-DRB1-01	27(45%)	5(5%)	0.001
HLA-DRB1-02	11(18%)	0(0%)	0.05
HLA-DRB1-03	10(16%)	50(50%)	0.03
HLA-DRB1-04*	21(35%)	10(10%)	0.05
- HLA-DRB1-0401	3(15%)	3(13%)	
- HLA-DRB1-0404	1(5%)	9(9%)	
- HLA-DRB1-0405	17(80%)	77(77%)	
DRB1*04 non-SE alleles	0 (0%)	1(1%)	
HLA-DRB1-07	4(6%)	25(25%)	0.02
HLA-DRB1-08	6(10%)	5(5%)	NS
HLA-DRB1-09	1(1.7%)	1(1%)	NS
HLA-DRB1-10	0(0%)	15(15%)	0.002
HLA-DRB1-11	5(8.3%)	20(20%)	NS
HLA-DRB1-12	10(16%)	16(16%)	NS
HLA-DRB1-13	16(26.7%)	30(30%)	NS
HLA-DRB1-14	1(1.7%)	4(4%)	NS
HLA-DRB1-15	0(0%)	16(16%)	0.002

Values are expressed as n (%). HLA: human leukocyte antigen. *According to HLA-DRB1*04 subtyping the shared epitopes encoding alleles *0409, *0413, *0416 and *0421 were classified as *0401 alleles, *0408, *0410 and *0419 were classified as *0404. The DRB1*0405 alleles were analyzed as a separate entity. All other DRB1*04 alleles were classified as DRB1*04 non-SE alleles.⁽⁴⁾

4) The combination of DRB1*04 + anti-CCP antibodies gave increase in the specificity and PPV to 92% and 90%, with a sensitivity and NPV of 42% and 41% respectively. The presence of single and double gene carriage of the HLA-DRB1*04 resulted in a significantly higher grade of disease activity ($P = 0.009$) compared with its absence. With regard to a possible gene dosage effect, no statistically significant difference between double and single gene carriage of the DRB1*04 was observed (data is not shown in tables).

Table 3: Univariate predictors of disease activity grade

	Sensitivity	Specificity	PPV	NPV	P value
Anti-CCP present	97%	36%	68%	90%	0.001
RF present	77%	36%	63%	53%	NS
CRP > 18	37%	72%	100%	51%	0.002
ESRI > 32	89%	40%	67%	71%	0.014
ANA	20%	68%	51%	45%	NS
AKA	65%	68%	74%	59%	0.01
Age > 50	29%	80%	67%	44%	NS
Duration > 6m	97%	8%	60%	67%	NS
Smoking	9%	92%	60%	42%	NS
Larsen score ≥35	80%	48%	68%	63%	0.02
HLA-DRB1*01	40%	48%	51%	36%	NS
HLA-DRB1*04	48%	84%	81%	54%	0.009

AKA: antikeratin antibody; anti-CCP: anti-cyclic citrullinated peptides antibodies; ANA: antinuclear antibodies; CRP: C-reactive protein; HLA: human leukocyte antigen; NPV: negative predictive value; PPV: positive predictive value; RF: rheumatoid factor

Table 4: Multivariate predictors of disease activity grade

	OR (95%CI)	P value
Anti-CCP present	19.6 (2.1–48.5)	0.009
HLA-DRB1*04	5.1 (1.2–21.1)	0.03

Anti-CCP: anti-cyclic citrullinated peptides antibodies; HLA: human leukocyte antigen, OR: Odds ratio; CI: confidence interval.

Predictors of radiological joint damage

ESR first hour > 32 had the highest sensitivity 98%, and NPV 93% (Table 5). In multivariate analysis, the independent predictors were HLA-DRB1*04, HLA-DRB1*01, CRP > 18, and RF (Table 6).

Discussion

Anti-CCP-antibody is a synthetic cyclic peptides containing citrulline. It is incorporated into newly proposed diagnostic criteria for rheumatoid arthritis, with higher sensitivity. The high predictive value of anti-CCP antibodies is indicating that citrullination and production of anti-CCP antibodies and RF are early processes in the development of RA [11]. The present work revealed strong complementarities of RF to anti-CCP as 65% patients lacking RF found to be positive for anti-CCP. Some reports suggest that RF and anti-CCP should be combined to reach optimum diagnostic properties [12]. Discrepancy can be explained by the fact that in the literature stratification for specificity was not performed, and therefore, the results of different studies are not comparable [13]. The frequencies of ANA and AKA were 36%, 52% respectively which indicate that testing for this spectrum of auto antibodies carries little advantage over RF or anti-CCP in diagnosing RA. A significant association of AKA with Anti-CCP may relate to that citrulline (profilaggrin) appears to be an essential

Table 5: Univariate predictors of radiological joint damage (Larsen score)

	Sensitivity	Specificity	PPV	NPV	P value
Anti-CCP present	90%	32%	74%	60%	0.05
RF present	81%	47%	77%	53%	0.03
CRP > 18	66%	63%	79%	46%	0.05
ESRI > 32	98%	68%	87%	93%	0.0001
ANA	21%	74%	73%	37%	NS
AKA	56%	58%	74%	38%	NS
Age > 50	32%	90%	86%	38%	NS
Duration > 6m	98%	11%	70%	67%	NS
Risk factor	7%	90%	60%	31%	NS
HLA-DRB1*01	53%	74%	81%	42%	0.05
HLA-DRB1*04	44%	84%	86%	54%	0.04

AKA: antikeratin antibody; anti-CCP: anti-cyclic citrullinated peptides antibodies; ANA: antinuclear antibodies; CRP: C-reactive protein; HLA: human leukocyte antigen; NPV: negative predictive value; PPV: positive predictive value; RF: rheumatoid factor.

Table 6: Multivariate predictors of radiological joint damage (Larsen score)

	OR (95%CI)	P value
RF present	2.5 (1.1–5.9)	0.01
CRP > 18	2.0 (0.2–5.3)	0.01
HLA-DRB1*01	4.5 (0.4–7.8)	0.002
HLA-DRB1*04	5.5 (0.4–8.7)	0.003

CRP: C-reactive protein; RF: rheumatoid factor; HLA: human leukocyte antigen; OR: Odds ratio; CI: confidence interval.

constituent of the antigenic determinants recognized by AKA [14].

In the present work, the sensitivity of anti-CCP was significantly higher than that of the RFs for prediction of grade of disease activity. This findings were supported by the data previously published [1,15].

It was reported that anti-CCP is an important predictor of radiological outcome. Preventing and diminishing joint damage is an important treatment goal in early rheumatoid arthritis [1].

According the prediction models reported by, Combe et al [2], for instance, variables reflecting disease activity also contribute to the prediction of joint damage. In this study, anti-CCP gave statistically significant differences between the positive and negative group with radiological damage. These data are comparable with the findings of other groups using a similar approach [15-17]. In our study serum anti-CCP added some prognostic information to ESR, CRP and RF, and it was the most sensitive predictor of joint damage, following ESR. These findings were in agreement with earlier reports [4], but other studies mentioned that anti-CCP was the only significant predictor, together with C reactive protein [18]. Vencovsky et al reported that the combination RF, CRP and anti-CCP2 antibodies revealed 10 times higher radiological damage [17] and early erosions [19]. However it was needed to be emphasized that the results based on dichotomization not useful in everyday clinical practice. So, we investigated whether appropriate cut off could be determined to identify those patients with the most aggressive radiological course.

Genetic studies have demonstrated that HLA alleles have been implicated in a number of chronic inflammatory diseases. RA has been associated with the SE of HLA-DRB1, which includes DRB1*04 and DRB1*01 alleles [19]. In the present study we found a discrepant relationship between RF and anti-CCP2 with the presence of the SE gene. Also our results do not support the opinion that there is a direct association between SE gene carriage and the occurrence of antibodies directed to CCP (or RFs)

leading to the development of RA, but rather suggest that there is a synergistic interaction between these factors. In contrast Berglin et al reported that patients with early RA, had significant association between anti-CCP antibodies and expression of DRB1*0401/0101 [5]. The conversion of arginine to citrulline in HLA-DRB1*0401 transgenic mice has been demonstrated to significantly increase activation of CD4+ T cells [20], so that individuals carrying the SE genes may have more sustained T- and B-cell responses to citrullinated antigens than non carriers [13]. The presence of citrulline within the HLA binding peptide enhances the peptide-MHC affinity and leads to the activation of CD4+ T cells during presentation of citrullinated antigens [20]. This specific T-cell dependent immune response to citrullinated peptides may contribute to the occurrence of RA. A quite contradictory suggestion is that HLA antigens do not predispose to the autoimmune disease per se but rather fail to provide protection. Abnormal T-cell regulation associated with certain HLA haplotypes leads to the loss of self-tolerance followed by polyclonal activation of T and B cells and the subsequent production of auto antibodies [21].

In the present work there was a statistically significant difference observed between carriage of the DRB1*04 and DRB1*01 alleles with radiological damage and DRB1*04 with disease activity, these findings were in agreement with those previously reported [13,17]. Recent genome wide scanning studies using micro satellite loci have confirmed that there is strong linkage between this region with susceptibility and severity of RA [22,23]. Moreover Petersson et al [4] reported that DRB1*04 SE double gene dose is associated with disease severity in RA. In contrast Eberhardt et al [24] did not find the genotypes to be strongly associated with disease severity after two and five years.

The role of the SE in the evolution of articular destruction has yielded conflicting results; however, in most studies the presence of the SE is associated with increased joint destruction [14,22]. Others revealed an association between HLA-DRB1*04 and erosive disease [25], Goronzy et al [25], and Petersson et al [4] considered homozygosity for HLA-DRB1*04 as a major predictor of the development of erosions. In Caucasians of Northern European origin, the double dose of DRB1*04 SE alleles (0401/0401 genotype) was associated with radiographic signs of progressive joint damage [26]. A strong concordance was revealed between our study and that in East Asian populations, in which DRB1*0401 is rare and DRB1*0405 is the most frequent RA associated HLA-DRB1 genotype. The latter allele has also been reported to be associated with an increased risk for extra-articular manifestations of RA [27]. In contrast, others reported that the presence of shared epitope in single or double dose did not provide

any predictive information on the development of joint damage and was not related to functional outcome [28]. In another study, by Matthey et al [16], reported that a predictive value of shared epitopes for radiological changes was restricted to RF negative patients. These discrepancies may reflect variability in the relative frequencies of HLA-DRB1*04 in different populations. Possibly, further studies on larger numbers of patients might solve this issue.

In the Present work the combination between anti-CCP, and the HLA-DR-B1*04 gave more identification of the grade of disease activity than a single test alone. The presence of RF, the SE, and anti-CCP2 antibodies yielded a 10 times higher average expectancy rate for a high radiological progression rate compared with the absence of the three parameters [15,16].

Conclusion

Anti-CCP2 is superior to RF for the detection of RA and provided predictive information on joint destruction and disease activity. The presence of RA associated antibodies and/or the SE genes are indicative for a poorer radiological outcome and higher grade of activity. Thus, we shall need new strategies, both in research and in clinical practice, where we may now possess a new means for analyzing the risk of RA in individuals who will have different needs to acquire such information.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

NAF participated in study design, data acquisition and critical manuscript revision. AME-E participated in study design, performing laboratory investigations other than HLA, and critical manuscript revision. EM participated in study design, HLA analysis, analysis and interpretation of data, performed the statistical analysis, wrote the manuscript, and performing manuscript drafting and submission. RMB participated in study design, HLA analysis, analysis and interpretation of data, and critical manuscript revision. HBH participated in HLA analysis and manuscript revision. SA participated in study design, data acquisition, and critical manuscript revision. MM participated in study design and data acquisition. H-AGR participated in laboratory investigations other than HLA and manuscript revision. FA participated in study design, data acquisition, and critical manuscript revision. All authors read and approved the final manuscript.

References

- Forslind K, Ahlmen M, Eberhardt K, Hafstrom I, Svensson B: **BARFOT Study Group. Prediction of radiological outcome in early rheumatoid arthritis in clinical practice: role of antibodies to citrullinated peptides (anti-CCP).** *Ann Rheum Dis* 2004, **63**:1090-1095.
- Combe B, Dougados M, Goupille P, Cantagrel A, Eliaou JF, Sibilia J, Meyer O, Sany J, Daures JP, Dubois A: **Prognostic factors for radiographic damage in early rheumatoid arthritis: a multiparameter prospective study.** *Arthritis Rheum* 2001, **44**:1736-1743.
- Lindqvist E, Eberhardt K, Bendtzen K, Heinegard D, Saxne T: **Prognostic laboratory markers of joint damage in rheumatoid arthritis. Prognostic laboratory markers of joint damage in rheumatoid arthritis.** *Ann Rheum Dis* 2005, **64**:196-201.
- Petersson IF, Sturfelt G, Nyhall-Wahlin BM, Truedsson L, Dechant SA, Matteson EL: **The impact of HLA-DRB1 genes on extra-articular disease manifestations in rheumatoid arthritis.** *Arthritis Res Ther* 2006, **8**:103.
- Berglin E, Padyukov L, Sundin U, Hallmans G, Stenlund H, Van Venrooij WJ, Klareskog L, Dahlqvist SR: **A combination of autoantibodies to cyclic citrullinated peptide (CCP) and HLA-DRB1 locus antigens is strongly associated with future onset of rheumatoid arthritis.** *Arthritis Res Ther* 2004, **6**:R303-308.
- DAS-Score.NL** [<http://www.das-score.nl/www.das-score.nl/index.html>]
- Kdahl C, Eberhardt K, Andersson SI, Svensson B: **Assessing disability in patients with rheumatoid arthritis. Use of a Swedish version of the Stanford Health Assessment Questionnaire** *Scand J Rheumatol* 1988, **17**:263-271.
- Larsen A, Dale K, Eek M: **Radiographic evaluation of rheumatoid arthritis and related conditions by standard reference.** *Acta Radiol Diagn (Stockh)* 1977, **18**(4):481-491.
- Saiki RK, Walsh PS, Levenson CH, Erlich HA: **Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes.** *Proc Natl Acad Sci USA* 1989, **86**:6230-6234.
- Schreuder GM, Hurley CK, Marsh SG, Lau M, Maiers M, Kollman C, Noreen H: **The HLA dictionary 1999: a summary of HLA-A, -B, -C, -DRB1/3/4/5, -DQB1 alleles and their association with serologically defined HLA-A, -B, -C, -DR and -DQ antigens.** *Tissue Antigens* 1999, **54**:409-437.
- Rantapaa-Dahlqvist S, Boman K, Tarkowski A, Hallmans G: **Up regulation of monocyte chemoattractant protein-1 expression in anti-citrulline antibody and immunoglobulin M rheumatoid factor positive subjects precedes onset of inflammatory response and development of overt rheumatoid arthritis.** *Ann Rheum Dis* 2007, **66**:121-123.
- Union A, Meheus L, Humbel RL, Conrad K, Steiner G, Moereels H, Pottel H, Serre G, De Keyser F: **Identification of citrullinated rheumatoid arthritis-specific epitopes in natural filaggrin relevant for anti-filaggrin autoantibody detection by line immunoassay.** *Arthritis Rheum* 2002, **46**:1185-1195.
- Goldbach-Mansky R, Lee J, McCoy A, Hoxworth J, Yarburo C, Smolen JS, Steiner G, Rosen A, Zhang C, Menard HA, Zhou ZJ, Palosuo T, Van Venrooij WJ, Wilder RL, Klippel JH, Schumacher HR Jr, El-Gabalawy HS: **Rheumatoid arthritis associated autoantibodies in patients with synovitis of recent onset.** *Arthritis Res* 2000, **2**:236-243.
- Suzuki K, Sawada T, Murakami A, Matsui T, Tohma S, Nakazono K, Takemura M, Takasaki Y, Mimori T, Yamamoto K: **High diagnostic performance of ELISA detection of antibodies to citrullinated antigens in rheumatoid arthritis** *Scand J Rheumatol* 2003, **32**(4):197-204.
- Matthey DL, Hassell AB, Dawes PT, Cheung NT, Poulton KV, Thomson W, Hajeer AH, Ollier WE: **Independent association of rheumatoid factor and the HLA-DRB1 shared epitope with radiographic outcome in rheumatoid arthritis.** *Arthritis Rheum* 2001, **44**:1529-1533.
- Matthey DL, Hassell AB, Dawes PT, Cheung NT, Poulton KV, Thomson W, Hajeer AH, Ollier WE: **Independent association of rheumatoid factor and the HLA-DRB1 shared epitope with radiographic outcome in rheumatoid arthritis.** *Arthritis Rheum* 2001, **4**:1529-1533.
- Vencovsky J, Machacek S, Sedova L, Kafkova J, Gatterova J, Pesakova V, Ruzickova S: **Autoantibodies can be prognostic markers of an erosive disease in early rheumatoid arthritis.** *Ann Rheum Dis* 2003, **62**:427-430.
- De Rycke L, Peene I, Hoffman IE, Kruihof E, Union A, Meheus L, Lebeer K, Wyns B, Vincent C, Mielants H, Boullart L, Serre G, Veys EM, De Keyser F: **Rheumatoid factor and anticitrullinated protein antibodies in rheumatoid arthritis: diagnostic value, associations with radiological progression rate, and extra-articular manifestations.** *Ann Rheum Dis* 2004, **63**:1587-1593.

19. Gregersen PK, Silver J, Winchester RJ: **The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis.** *Arthritis Rheum* 1987, **30**:1205-1213.
20. Hill JA, Southwood S, Sette A, Jevnikar AM, Bell DA, Cairns E: **Cutting edge: the conversion of arginine to citrulline allows for a high-affinity peptide interaction with the rheumatoid arthritis-associated HLA-DRB1*0401 MHC class II molecule.** *J Immunol* 2003, **171**:538-541.
21. Zanelli E, Breedveld FC, de Vries RR: **HLA association with autoimmune disease: a failure to protect.** *Rheumatology* 2000, **39**:1060-1066.
22. Conrad K, Sack U, Shoenfeld Y: **From proteomics to molecular epidemiology: relevance of autoantibodies.** *Autoimmun Rev* 2003, **2**:165-169.
23. Jawaheer D, Seldin MF, Amos CI, Chen WV, Shigeta R, Monteiro J, Kern M, Criswell LA, Albani S, Nelson JL, Clegg DO, Pope R, Schroeder HW Jr, Bridges SL Jr, Pisetsky DS, Ward R, Kastner DL, Wilder RL, Pincus T, Callahan LF, Flemming D, Wener MH, Gregersen PK: **Screening the genome for rheumatoid arthritis susceptibility genes: a replication study and combined analysis of 512 multicase families.** *Arthritis Rheum* 2003, **48**:906-916.
24. Eberhardt K, Fex E, Johnson U, Wollheim FA: **Associations of HLA-DRB and -DQB genes with two and five year outcome in rheumatoid arthritis.** *Ann Rheum Dis* 1996, **55**:34-39.
25. Goronzy JJ, Matteson EL, Fulbright JW, Warrington KJ, Chang-Miller A, Hunder GG, Mason TG, Nelson AM, Valente RM, Crowson CS, Erlich HA, Reynolds RL, Swae RG, O'Fallon WM, Weyand CM: **Prognostic markers of radiographic progression in early rheumatoid arthritis.** *Arthritis Rheum* 2004, **50**:43-54.
26. Turesson C, Jacobsson L, Bergstrom U, Truedsson L, Sturfelt G: **Predictors of extra-articular manifestations in rheumatoid arthritis.** *Scand J Rheumatol* 2000, **29**:358-364.
27. Koh WH, Chan SH, Lin YN, Boey ML: **Association of HLA-DRB1*0405 with extra-articular manifestations and erosions in Singaporean Chinese with rheumatoid arthritis.** *J Rheumatol* 1997, **24**:629-632.
28. Lindqvist E, Saxne T, Geborek P, Eberhardt K: **Ten year outcome in a cohort of patients with early rheumatoid arthritis: health status, disease process, and damage.** *Ann Rheum Dis* 2002, **61**:1055-1059.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

