

δ -Opioid Receptor Activation Modified MicroRNA Expression in the Rat Kidney under Prolonged Hypoxia

Xiaozhou He¹, Yilin Yang¹, Feng Zhi¹, Meredith L. Moore², Xuezhi Kang³, Dongman Chao^{2,4}, Rong Wang¹, Gianfranco Balboni⁵, Severo Salvadori⁶, Dong H Kim², Ying Xia^{2,4*}

1 Research Institute of Modern Medicine, The Third Clinical College of Soochow University, Changzhou, Jiangsu, People's Republic of China, **2** Department of Neurosurgery, The University of Texas Medical School at Houston, Houston, Texas, United States of America, **3** Laboratory of Molecular Neurology, Shanghai Research Center for Acupuncture and Meridians, Shanghai, People's Republic of China, **4** Department of Pediatrics, Yale University School of Medicine, New Haven, Connecticut, United States of America, **5** Department of Life and Environment Sciences, University of Cagliari, Cagliari, Italy, **6** Department of Pharmaceutical Sciences, University of Ferrara, Ferrara, Italy

Abstract

Hypoxic/ischemic injury to kidney is a frequently encountered clinical problem with limited therapeutic options. Since microRNAs are differentially involved in hypoxic/ischemic events and δ -opioid receptor (DOR) activation is known to protect against hypoxic/ischemic injury, we speculated on the involvement of DOR activation in altering the microRNA (miRNA) expression in kidney under hypoxic condition. We selected 31 miRNAs based on microarray data for quantitative PCR analysis. Among them, 14 miRNAs were significantly altered after prolonged hypoxia, DOR activation or a combination of both. We found that 1) DOR activation alters miRNA expression profiles in normoxic conditions; 2) hypoxia differentially alters miRNA expression depending on the duration of hypoxia; and 3) DOR activation can modify hypoxia-induced changes in miRNA expression. For example, 10-day hypoxia reduced the level of miR-212 by over 70%, while DOR activation could mimic such reduction even in normoxic kidney. In contrast, the same stress increased miR-29a by >100%, which was reversed following DOR activation. These first data suggest that hypoxia comprehensively modifies the miRNA profile within the kidney, which can be mimicked or modified by DOR activation. Ascertaining the targeted pathways that regulate the diverse cellular and molecular functions of miRNA may provide new insights into potential therapies for hypoxic/ischemic injury of the kidney.

Citation: He X, Yang Y, Zhi F, Moore ML, Kang X, et al. (2013) δ -Opioid Receptor Activation Modified MicroRNA Expression in the Rat Kidney under Prolonged Hypoxia. PLoS ONE 8(4): e61080. doi:10.1371/journal.pone.0061080

Editor: Cesar V. Borlongan, University of South Florida, United States of America

Received: November 20, 2012; **Accepted:** March 5, 2013; **Published:** April 15, 2013

Copyright: © 2013 He et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Supported by National Institutes of Health (HD-034852 and AT004422), National Science Foundation of China (31071046), Changzhou Science Development Project (CS20102010), Changzhou Health Bureau (ZD201007), and National Basic Research Project (09CB522901; 12CB518502). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Ying.Xia@UTH.TMC.edu

Introduction

Hypoxic/ischemic injury to kidney is a frequently encountered problem in vascular and urologic clinic given the sensitivity of kidney to changes in oxygen/blood delivery. Although blood flow to the kidney accounts for 20% of the cardiac output, the oxygen diffusion between arterial and venous vessels running parallel and in close proximity that keeps oxygen tension in the renal tissue comparatively low [1]. This high sensitivity to changes in oxygen tension makes the kidney prone to hypoxic/ischemic injury. Many clinical conditions, including chronic respiratory insufficiency, prolonged cardiovascular dysfunction, acute renal failure, chronic glomerular, tubulointerstitial or renovascular disease, diabetes, hypertension, aging, renal hypertrophy, anemia and obstructive uropathy, can reduce renal oxygenation and result in kidney dysfunction.

Hypoxia plays a significant role in the development of nephrotoxic acute kidney injury, radiocontrast nephropathy, and acute glomerulonephritis [1,2]. In patients with chronic respiratory insufficiency, the arterial blood hypoxemia impairs the renal clearance of salt ions, urea and creatinine. The correlations between the renal clearances of these substances and the pO₂ in

arterial blood has a greater significance compared with the effects of pCO₂ or [H⁺] levels, which in turn suggests the existence of a causal association between hypoxemia and renal dysfunction [3]. Severe energy depletion and subsequent activation of a number of critical alterations in metabolism and gene expression occur under hypoxic conditions. In fact, chronic hypoxia may be the final common pathway in end-stage kidney failure in kidney diseases. Hypoxic injury may have a pivotal role in both the development and progression of acute and chronic kidney diseases. Although hypoxic/ischemic insults are known to be associated with free radical induced injuries, the precise mechanisms underlying hypoxia-induced kidney injury remains poorly understood.

Recent research suggests that hypoxia, especially a prolonged stress, may influence the kidney through alterations in the RNAome. Over the past few decades, miRNAs have rapidly emerged as a new frontier in across different medical fields including kidney research [4,5]. The kidney exhibits a unique miRNA expression profile [5]. Therefore, miRNA expression is likely important for renal function in normoxic condition and for the genesis of hypoxic pathology. Based on our previous studies [6–8], we hypothesized that hypoxia could influence the expres-

sion of miRNA in the kidneys. However, current literature on research studies involving this topic is scant.

There is no promising therapy available for the prevention and treatment of hypoxic kidney injury. We have recently demonstrated that the δ-opioid receptor (DOR) is protective against anoxic/ischemic injury in the brain and the underlying mechanisms involve the regulation of ionic homeostasis and antioxidative capacity [6,7,9–17]. In addition, DOR activation is also protective against hypoxic/ischemic injury in the heart [18]. In fact, studies show that DOR agonists can prolong survival of peripheral organs, such as lung, heart, liver, and kidney preserved en bloc or as a single preparation [19]. A growing body of evidence indicates that miRNAs play several regulatory roles in opioid pharmacology [20–22]. In view of this, we examined if DOR activation protects the kidney against hypoxic injury through a miRNA-mediated mechanism in the present study.

Our present study is a stepping stone in understanding hypoxic effect on the miRNAs in the kidney and its response to DOR activation in normoxia vs. hypoxia since there is no previously published data available on the subjects. Specifically, this work was conducted in adult rat kidney to determine: 1) if DOR activation alters miRNA expression profiles in the normoxic condition; 2) if short-term and prolonged hypoxia differentially alter miRNA expression; and 3) if DOR activation modifies hypoxia-induced changes in miRNA expression. The outcome of this study on renal miRNAs under hypoxic stress can result in discovery of new diagnostic biomarkers for kidney diseases. This in turn will provide novel insights into newer mechanisms of modulating the complex regulatory networks in the hypoxic kidney.

Results

Hypoxia-induced Changes in Body and Kidney Weights

Changes in body and kidney weight were monitored in all animals. Hypoxic rats demonstrated a reduced ability to gain weight that was significant after 24 hours of hypoxic treatment and persisted for the remainder of the experiment (Figure 1a). Individual kidney weights were also significantly lower in the hypoxic animals (data not shown). The addition of UFP-512, a DOR agonist, did not affect the rate of increase in body weight neither in the control nor the hypoxic groups. The ratios of kidney/body weight were significantly lower following chronic hypoxia indicating that the kidneys exhibited a greater sensitivity to oxygen deprivation (Figure 1b). This reduction in kidney/body weight ratio was recorded earlier in the UFP-512 treated animals and was statistically significant as early as 24 hours after hypoxia (Figure 1b).

MicroRNA Microarray Analysis

To identify the molecular changes induced by hypoxia, comparative miRNA microarray analysis was performed on RNA isolated from the kidneys of rats exposed to 10-day hypoxia and normoxia. Analysis revealed that 22 miRNAs were significantly and differentially expressed between hypoxic and control samples (Figure 2). Of these 22 subtypes of miRNAs known to be expressed in the rat kidney, 12 were significantly downregulated, while 10 were significantly upregulated. All these changes in 22 miRNAs were verified by RT-PCR. In addition to the miRNAs selected following microarray analysis, we also investigated supplementary miRNAs which were proven to be correlated with hypoxia or ischemia-responsiveness in other organs. In total, 31 miRNAs were selected for further quantitative PCR analysis, which include let-7d, let-7f, miR-25*, miR-101b, miR-186, miR-187, miR-212, miR-291a-5p, miR-292-5p, miR-298, miR-324-3p,




Figure 1. (A) Body weight measurement. Data points represent the mean \pm SE with a minimum $n=9$. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. ** $p<0.01$ (C vs. H; C+DOR vs. H+DOR) for all time points encompassed by the bar. **(B) Kidney weight/body weight ratios** for each study group. * $p<0.05$, ** $p<0.01$ with student's t-test.

doi:10.1371/journal.pone.0061080.g001

miR-347, miR-351, miR-363*, miR-365, miR-370, miR-431, miR-466b, miR-487b, miR-511, miR-615-5p, miR-743a, miR-20b-3p, miR-20b-5p, miR-21, miR-29a, miR-29b, miR-31, miR-135a, miR-199a-3p and miR-199a-5p [23–28].

Hypoxia and DOR Activation Shifted miRNA Expression Profiles of Rat Kidney

Quantitative RT-PCR analysis confirmed the direction and amplitude of all miRNA changes with the exception of let-7d, miR-25*, miR-187, miR-291a-5p, miR-292-5p, miR-365, miR-431, miR-487b, miR-615-5p, miR-743a, miR-20b-3p, miR-199a-3p which remained unaltered or showed no statistical significance. A subset of miRNAs was influenced only by long-term exposure to hypoxia or DOR agonist. Chronic hypoxia for 10 days depressed miR-363* expression greater than 50%. DOR treatment alone did not significantly alter basal miR-363* levels in the normoxic kidney, but prevented hypoxia-induced miR-363* down-regulation (Figure 3a). In contrast, the same hypoxic exposure did not induce any significant reduction in let-7f despite the tendency to decrease let-7f under hypoxia (Figure 3b). However, reduced let-7f levels were noted in the kidneys following long-term DOR agonist exposure (Figure 3b). As with let-7f, miR-370 did not alter in response to hypoxia alone. Conversely, UFP-512 tended to increase miR-370 in normoxic condition but tended to decrease it under hypoxia (Figure 3c), suggesting a differential regulation of miR-370 in response to DOR signals in different environments.

The miRNA-466b levels were significantly repressed by either long-term hypoxia or UFP-512 treatment. Interestingly, the




Figure 2. Dendrogram of aberrant miRNA expression in the kidney after hypoxia. Unsupervised hierarchical cluster analysis of 22 miRNAs differentially expressed in the kidney after hypoxia separates hypoxic kidney from control samples based on miRNA profiling. Upregulated miRNAs are labeled red and downregulated miRNAs are colored green.
doi:10.1371/journal.pone.0061080.g002




Figure 3. Relative miRNA expression levels of miR-363*, miR-let-7f, and miR-370 in the kidney following 10 days of hypoxia as determined by quantitative RT-PCR. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. **p<0.01.
doi:10.1371/journal.pone.0061080.g003

combination of long-term hypoxia and UFP-512 treatment returned miR-466b expression to control levels (Figure 4a). The miR-511 levels were also down-regulated by chronic hypoxia. The addition of UFP-512 further downregulated miR-511 levels in the




Figure 4. Relative miRNA expression levels of miR-466b and miR-511 in the kidney following 10 days of hypoxia as determined by quantitative RT-PCR. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. doi:10.1371/journal.pone.0061080.g004

hypoxic kidney when compared to DOR treatment with UFP-512 alone (Figure 4b).

Longitudinal miRNA Changes in Response to Hypoxia

Several miRNAs were altered at two given time points in the least. In the presence of UFP-512, miR-298 showed a hypoxia-induced downregulation that was lost after 10 days of hypoxia. Hypoxia alone did significantly upregulate miR-298 expression by 50% at 10 days while DOR treatment returned the levels to baseline. The miRNA 324-3p was consistently upregulated by hypoxia with or without DOR signaling following 5-day treatment. After 10 days of treatment, the upregulated miR324-3p expression seen in 5-day hypoxia disappeared and was further downregulated by DOR treatment in hypoxia. A similar DOR-induced downregulation in the presence of hypoxia was also found in miR-20b-5p at 5-day time point, which was disappeared at 10 days of hypoxia but replaced by a 3-fold stimulation of miR-20b-5p expression with UFP-512 treatment (Figure 5).

The miR-347 and -212 expression showed no appreciable effect following exposure to hypoxia for 1-day and/or DOR activation, whereas a longer duration (5 or 10 days) of treatment demonstrated marked alterations in their levels of renal expression (Figure 6). While hypoxia alone had no effect on miR-347 expression after 5 days, UFP-512 treatment produced a 60% reduction in normoxic condition, and induced a similar change under hypoxia. The DOR-triggered depression was further increased to 25% of the control level after 10 days. DOR activation did not alter these hypoxic changes significantly (Figure 6a). In 5-day samples, miR-212 was largely reduced by UFP-512 in both the control and hypoxic conditions. Hypoxia alone did not significantly affect miR-212 expression until the 10-day time point. With this chronic hypoxia, miR-212 levels were repressed to less than 40% of the control in all combinations of hypoxia and UFP-512 treatment tested (Figure 6b).

Four miRNAs investigated were altered by hypoxia and/or DOR activation at all three time points. In the case of miR-351, 10-day hypoxia induced a 30% decrease though no significant reduction was seen after 1- or 5-day hypoxia. The miR-351 expression after UFP-512 treatment showed an initial slight yet significant upregulation following 1-day treatment, followed by a substantial downregulation by 45% after 5 days. Chronic UFP-512 treatment did not alter miR-351 levels under hypoxia. Hypoxia alone downregulated the miR-351 expression after 10 days of exposure (Figure 7). Although UFP-512 treatment alone did not alter miR-29a levels, the addition of UFP-512 to the hypoxic kidney significantly downregulated the miR-29a expression to 40–60% of the control levels after both 1 and 5 days of hypoxia. Following chronic exposure to hypoxia, miR-29a was upregulated by more than 100% as opposed to hypoxia alone. The addition of UFP-512 appeared to reduce the hypoxia-increased level of miR-29a (Figure 7).

Hypoxia for 1 day reduced miR-21 expression by 40% and the addition of UFP-512 further reduced its expression to 25% of the control levels (Figure 8). After 5 days the hypoxia-induced reduction disappeared, but the combination of hypoxia and UFP-512 produced a dramatic loss of miR-21. The presence of UFP-512 alone suppressed miR-29b levels following 1 day of hypoxia exposure (Figure 8). Either hypoxia or UFP-512 alone had no effect of miR-29b at 5 days, but the combination significantly stunted miRNA levels similar to miR-21. After 10 days, DOR stimulation significantly depressed miR-29b levels only in the presence of hypoxia.

Discussion

Current research on describing the renal miRNA profiles focusses on neoplastic lesions or cultured fibroblasts. Here we present the first data that identify 14 miRNAs in the intact kidney altered in response to chronic hypoxia and DOR activation. Our comprehensive data demonstrate that 1) DOR activation shifts miRNA expression profiles in normoxic conditions; 2) hypoxia differentially alters the miRNA expression profiles depending on hypoxic duration; and 3) DOR activation modifies hypoxia-induced changes in miRNA expression, suggesting significant hypoxia- and DOR-induced alterations in the miRNA profiles of the kidney.

There is accumulating evidence on the diagnostic and therapeutic potentials of a new class of small regulatory RNA molecules, termed microRNAs. These miRNAs are a class of endogenous non-coding RNAs comprised of 19–24 nucleotides that regulate the post-transcriptional silencing of protein-coding genes by base-pairing to complementary sites on their target









Figure 5. Relative miRNA expression levels of miR-298, miR-324-3p, and miR-20b-5p in the kidney following either 5 or 10 days of hypoxia as determined by quantitative RT-PCR. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. doi:10.1371/journal.pone.0061080.g005

mRNAs, thus suppressing protein synthesis, promoting protein degradation or translational blockade. Several miRNAs (e.g., miR-29) that showed a dynamic change with hypoxic durations in this work regulate the process of oxidative stress and inflammation and are involved in regulation of apoptotic and survival signal pathways. The miR-29 family directly target at least 16 extracellular matrix genes and are relevant to renal and cardiovascular injury [29–31]. Recent research suggests that miR-29 inhibits cell proliferation and induces cell cycle arrest [32], modulates oxidative injury [33] and promote apoptosis through a mitochondrial pathway that involves Mcl-1 and Bcl-2 [28]. We observed that 1-day hypoxia down-regulated miR-29a expression, which is probably an adaptive compensation of the kidney for overcoming the hypoxic stress. However, the kidney loses this ability to compensate after a prolonged period (5 days) of hypoxia. Furthermore, hypoxic exposure for an even longer duration, i.e., 10 days, significantly increased both miR-29a and miR-29b levels that can explain induction of renal injury under prolonged hypoxia. Interestingly, DOR activation decreases the level of miR-29a in all these scenarios, suggesting a potential for therapeutic utilization of DOR activation in hypoxic/ischemic renal injury.

A few of the miRNAs showed changes only after 10 days of hypoxia. For example, chronic hypoxia largely reduced expression

of miR-363*. DOR activation prevents the hypoxic reduction of miR-363*, suggesting that DOR signaling may regulate miR-363*, thus reducing hypoxic/ischemic injury. In contrast, DOR activation demonstrated an opposite effect on other miRNAs. For example, when hypoxia exposure decreased miRNA let-7f, DOR activation failed to increase the level of let-7f. Instead, it further potentiated the let-7f reduction under hypoxia, which suggests a differential regulation of various miRNAs through DOR activated pathways in the hypoxic kidney.

Our longitudinal study identified three DOR-sensitive miRNAs (miR-347, -212 and -351) that were downregulated after 5 days way ahead in time when the same change was induced by hypoxia after 10 days of exposure. These three miRNAs are predicted to target the scaffolding protein Bassoon, involved in vesicle clustering and regulate trans-Golgi trafficking in the hypoxic kidney. In a study on alcoholic liver disease in which alcohol-induced gut leakiness is a key factor, ethanol increased miR-212 expression, decreased the level of Zonula occludens 1 protein, disrupted tight junctions, and increased the permeability of monolayers of Caco-2 cells [34]. This study suggests an adverse effect of increased miR-212 on tissues in pathophysiological conditions. Therefore, the hypoxic reduction of miR-212 may be an adaptive strategy of the kidney in response to hypoxia. The notion that DOR action could mimic the hypoxic change in miR-212 provides additional evidence since DOR activation is protective against hypoxic injury [6–8,16]. Also, these miRNAs are predicted to target the HIF-regulator MAF and prolong the expression of this hypoxia-specific factor. Accumulated HIF binds to the hypoxia-responsive element within the erythropoietin promoter and activates transcription of erythropoietin to increase erythropoietin production [1]. Erythropoietin has been proven as a hypoxia-responsive cytokine to provide protective effects in the damaged brain during hypoxic/ischemic events and neurodegenerative diseases [35,36]. Therefore, the above-mentioned changes may potentially induce a protective effect on the hypoxic kidney.

The downregulation of miR-21 and miR-29b produced by either hypoxia or DOR activation in 1-day group appeared to be reversed by day 5 unless the two treatments (hypoxia plus DOR activation) were given simultaneously. Addition of UFP-512 suppresses these miRNAs and enhances the expression of a subset of mRNAs targeted by miR-21 and miR-29b. Expression of these genes is important for the structure of the basement membrane and kidney development, cell signaling as well as the matrix and junctional proteins matrilin and vinculin. Our data suggest that their expression could be modified during hypoxia with the addition of DOR agonists and thus affect the survival of the kidney in hypoxic environment. DOR is known to prevent hypoxic disruption of Na^+ and K^+ homeostasis in the brain through a PKC-dependent pathway [12–15,17]. DOR activation may regulate ionic homeostasis partially via miR-21 since PKCε and several K^+ channels are predicted targets of miR-21. Recent evidence also suggests the miR-21 inhibition of the PKC substrate MARCKS [37]. On the other hand, DOR activation to prolong miR-29b suppression could lengthen the expression of HIF3a as well as its transcriptional regulators MAFG and MAFB.

Proteinuria, resulting from a disrupted slit diaphragm, is an indication of kidney injury. Recently, the presynaptic adhesion molecule, neurexin-1, was reported to be present in the podocytes of the glomerular epithelial cells forming this capillary wall barrier




Figure 6. Relative miRNA expression levels of miR-347 and miR-212 in the kidney following either 1, 5, or 10 days of hypoxia as determined by quantitative RT-PCR. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. doi:10.1371/journal.pone.0061080.g006




Figure 7. Relative miRNA expression levels of miR-351 and miR-29a in the kidney following either 1, 5, or 10 days of hypoxia as determined by quantitative RT-PCR. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. * $p < 0.05$, ** $p < 0.01$. doi:10.1371/journal.pone.0061080.g007




Figure 8. Relative miRNA expression levels of miR-21 and miR-29b in the kidney following either 1, 5, or 10 days of hypoxia as determined by quantitative RT-PCR. C, control; C+DOR, control+UFP-512; H, hypoxia; H+DOR, hypoxia+UFP-512. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. doi:10.1371/journal.pone.0061080.g008

Table 1. Primer sequences used for quantitative RT-PCR.

miRNA	Forward Primer	Reverse Primer
rno-let-7d	ACACTCCAGCTGGGAGAGGTAGTAGTTGC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAACTATGC
rno-let-7f	ACACTCCAGCTGGGTGAGGTAGTAGATTGT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAACTATAC
rno-mir-101b	ACACTCCAGCTGGGTACAGTACTGTGATA	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTTCAGCTA
rno-mir-135a	ACACTCCAGCTGGGTATGGCTTTTTATTCTT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTACATAG
rno-mir-186	ACACTCCAGCTGGGCAAAGAATTCTCCTTT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAGCCAAA
rno-mir-187	ACACTCCAGCTGGGTCTGTCTTGTGTGTC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCCGGCTGC
rno-mir-199a-3p	ACACTCCAGCTGGGACAGTAGTCTGCACAT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTAACCAAT
rno-mir-199a-5p	ACACTCCAGCTGGGCCAGTGTTCAGACTAC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGGAACAGGT
rno-mir-20b-3p	ACACTCCAGCTGGGACTGCAGTGTGAGCAC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCCAGAAGT
rno-mir-20b-5p	ACACTCCAGCTGGGCAAAGTGCTCATAGT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGACCTGCAC
rno-mir-21	ACACTCCAGCTGGGTAGCTTATCAGACTGA	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTCAACATC
rno-mir-212	ACACTCCAGCTGGGTAACAGTCTCCAGTCA	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTGGCCGTG
rno-mir-25*	ACACTCCAGCTGGGAGGCGGAGACACGGGC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGGCAATTGC
rno-mir-291a-5p	ACACTCCAGCTGGGCATCAAAGTGGAGGCC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAGAGAGGG
rno-mir-292-5p	ACACTCCAGCTGGGACTCAAAGTGGGGCT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCAAAGAG
rno-mir-298	ACACTCCAGCTGGGGCAGAGGAGGGCTGTT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGGGGAAGAA
rno-mir-29a	ACACTCCAGCTGGGTAGCACCATTGAAAT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTAACCGAT
rno-mir-29b	ACACTCCAGCTGGGTAGCACCATTGAAATC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAACACTGA
rno-mir-31	ACACTCCAGCTGGGAGGCAAGATGTGGCA	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCAGCTATG
rno-mir-324-3p	ACACTCCAGCTGGGCCACTGCCAGGTGC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCCAGCAGC
rno-mir-347	ACACTCCAGCTGGGTGTCCCTCTGGGT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTGGGCGAC
rno-mir-351	ACACTCCAGCTGGGTCCCTGAGGAGCCCTTTGA	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTCAAGGCTC
rno-mir-363*	ACACTCCAGCTGGGCGGGTGGATCAGGATG	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAAATTGCA
rno-mir-365	ACACTCCAGCTGGGTAATGCCCTAAAAAT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGATAAGGAT
rno-mir-370	ACACTCCAGCTGGGGCTGCTGGGTGGAAC	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAACCAGGT
rno-mir-431	ACACTCCAGCTGGGTGTCTTGACAGGCCGT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTGCATGAC
rno-mir-466b	ACACTCCAGCTGGGTATGTGTGTGTATG	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCATGGACA
rno-mir-487b	ACACTCCAGCTGGGAATCGTACAGGTCA	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAGTGGATG
rno-mir-511	ACACTCCAGCTGGGATGCCTTTTGTCTG	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTGAAGTCA
rno-mir-615-5p	ACACTCCAGCTGGGGGGTCCCGGTGCT	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGGATCCGAG
rno-mir-743a	ACACTCCAGCTGGGAAAGACGCCAAACTG	CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGTCTACCCA
U6	CTCGCTTCGGCAGCACA	AACGCTTCACGAATTTGCGT

doi:10.1371/journal.pone.0061080.t001

in the kidney [38]. Models of kidney injury demonstrate a loss of neurexin and propose a role for the transmembrane protein in the formation and maintenance of the kidney slit diaphragm [39]. Neurexophilin, a ligand for neurexin, is one of the predicted targets of the hypoxia-upregulated miR-298. Prolonged hypoxia (10 days) significantly increased the level of miR-298, suggesting a downregulation of Neurexophilin. This alteration could potentially reduce the functionality of the neurexin signaling pathway, which may mimic the proteinuria/injury seen in other models of kidney disease and contribute, at least partially, to kidney damage in prolonged hypoxia. The application of DOR agonist attenuated these changes in miR-298 expression. In addition, the downregulation of miR-511 seen at 10 days could contribute to injury recovery by stimulating the translation of other neurexin isoforms.

The disruption of the neuropilin receptor system could upset the balance of semaphorin and VEGF signaling necessary to maintain the glomerular filtration barrier [40]. Plexin A1 is an important component of the neuropilin receptor complex and a known target of the three altered miRNAs studied in the present

work (20b-5p, 347 and 466b). No clear pattern was observed in the expressions of these miRNAs. As for instance, miR-20b-5p was upregulated by DOR agonist in the 10-day group, while miRs-347 and 466b were suppressed under the same conditions. In contrast, DOR activation significantly upregulated miR-466b expression in the hypoxic kidney. Further investigation is required to determine a role for miRNA regulation of semaphoring signaling in hypoxic/ischemic kidney.

Hypoxic rats were unable to maintain the body and kidney weight gains as opposed to the control animals, concurrent to our previous observations [8,41–43]. This reflects a mismatch between energy production and consumption under hypoxic conditions. Interestingly, the reductions in kidney/body weight ratio appeared earlier in the DOR-treated kidneys when compared to controls. Since opioid agents may affect feeding and appetite [44,45], it is likely that UFP-512 influences the animal’s feeding through a DOR signaling pathway.

In summary, hypoxia comprehensively alters miRNA expression in the kidney with differential changes in various miRNAs

depending on the duration of hypoxia. The altered miRNAs are known to target the antioxidant, apoptotic, survival and may other pathways. DOR activation has a significant effect on the miRNAome in the kidney under both normoxic and hypoxic conditions, leading to an improved condition for renal survival and functioning, though it does not improve the rate of body weight gain under hypoxia. MicroRNAs mediate mRNA suppression by recruiting miRNA-induced silencing complex (miRISC). The core of miRISC contains miRNA-loaded Argonature proteins (AGOs) and GW182. Since DOR activation greatly influences the miRNA profiling under hypoxic condition, the activity of AGOs and GW182 could be changed in response to hypoxia and DOR activation and needs to be further investigated. Moreover, it is important to explore if DOR inactivation with its antagonists influences the hypoxia and/or DOR activation induced changes in miRNAs. Further illumination of the targeted pathways associated with hypoxic kidney injury and the mechanism of DOR activation and the underlying mechanism may provide new insights into the potential therapeutics for hypoxic/ischemic injury of the kidney.

Materials and Methods

Animals

The experiments were approved by the Review Board and the Animal Care and Use Committee of Shanghai Research Center for Acupuncture and Meridians, and were performed in accordance with its guidelines. Male Sprague Dawley Rats were purchased from Shanghai Experimental Animal Center of Chinese Academy of Sciences. Twenty-one day old rats were randomly divided into 4 groups: (A) control, (B) DOR agonist (UFP-512), (C) chronic anoxia, and (D) chronic anoxia+DOR agonist (UFP-512). Groups A and B were raised in normal air and Groups C and D under hypoxic conditions.

Induction of Chronic Hypoxia

Chronic hypoxia was induced as previously described [8,41,43,46]. In brief, the hypoxic facility consisted of two parts: a plexiglass box (1.1 m×0.7 m×0.6 m) and an O₂/CO₂ analyzer to maintain an O₂ level at 9.5%–10% in the box. The rats of Groups C and D were kept in the hypoxic box for 1, 5 or 10 days. The box was rapidly cleaned daily when animals were removed to record body weights.

DOR Activation with UFP-512

The rats of Groups B and D were subjected to intraperitoneal injection of UFP-512 (H-Dmt-Tic-NH-CH[CH₂-COOH]-Bid), a specific and potent DOR agonist synthesized by our team [13]. The injections (1 mg/kg in <1 ml) were performed on day 0 (immediately before the beginning of the hypoxia), day 4 and day 8. As a control, Groups A and C received the same amount of saline.

Tissue Collection

After 1, 5 or 10 days of hypoxia, the rats were decapitated after deep anesthesia. Their kidneys were rapidly removed, weighed, frozen in liquid nitrogen and stored at –80°C until use.

RNA Extraction and Microarray Experiments

The expression profile of kidney miRNAs after chronic hypoxia has not been previously described. Therefore, microarrays were used to examine the expression of miRNAs in the kidney after 10 days of hypoxia. Tissue samples were removed from liquid

nitrogen and total RNA was extracted using Trizol Reagent (Invitrogen, Carlsbad, CA, USA). RNA labeling and hybridization on miRNA microarray chips were performed as previously described [47,48]. In brief, 50 µg of total RNA were purified using the mirVANA miRNA isolation kit (Ambion, Carlsbad, CA, USA) to enrich for the small RNA fraction. The purified RNA was labeled with Cy3 and hybridization was carried out on a miRNA microarray chip (CapitalBio Corp., Beijing, China) containing 509 probes in triplicate, corresponding to 435 human miRNA genes. Hybridization signals were detected and the scanned images were used for quantification as described previously [47,48]. Data analysis employed a two class-paired analysis within SAM, a statistical method that calculates a score for each gene and therefore identifies genes that are significantly associated with an outcome variable, such as hypoxia exposure. For this analysis, a false discovery rate of <5% was selected. MicroRNAs were considered significantly altered only when they fulfilled three criteria: (1) mean fold change >2 or <0.5; (2) *q*-value = 0; and (3) SAM score >2 or <–2 [49,50]. After SAM analysis selection, the data of 22 miRNAs were assessed using unsupervised hierarchical clustering. Predicted targets for altered miRNAs were identified by cross-referencing the online data-bases: microCosm, TargetScan and Pictar.

Quantitative RT-PCR

Total RNA isolated as above was treated with RNase-free DNase using a standardized protocol. Assays to quantify the mature miRNA were conducted as described previously [51,52]. Briefly, 1 µg of total RNA was reverse-transcribed to cDNA using AMV reverse transcriptase and looped antisense primers. The mixture was incubated at 16° for 15 min, 42° for 60 min and 85° for 5 min to generate a library of miRNA cDNAs. SYBR Green real-time PCR was then performed using a standardized protocol with a sequence detection system (Model 7500, Applied Biosystems, Foster City, CA, USA). The primers are listed in Table 1. In each assay, 1 µl of cDNA (1:50 dilution) was used for amplification. The reactions were incubated in a 96-well optical plate at 95° for 5 min, followed by 40 cycles consisting of a 15 s interval at 95° and a 1-min interval at 60°. All reactions were performed in triplicate. After the experiments were completed, the threshold cycle (Ct) values were determined using the default threshold settings and for analyzing the RT-PCR data. The relative amount of each gene to internal control was calculated by using the equation $2^{-\Delta Ct}$, in which $\Delta Ct = Ct_{miRNA} - Ct_{U6}$. To exclude extreme outliers, miRNAs with expression lower than a threshold ($Ct_{miRNA} - Ct_{U6} < 15$, mean fold change >2 or <0.5) were eliminated. The retained Ct data were normalized, mean-centered and log₂-transformed.

Statistical Analysis

Data are presented as mean ± SE with a minimum n = 3 for each group. The mean value of Group A for each time point is set equal to one. Statistical significance was determined using either a student's t-test or a one-way ANOVA followed by Tukey's Multiple Comparison Test on paired columns as appropriate.

Author Contributions

Conceived and designed the experiments: YX XH. Performed the experiments: FZ XK RW. Analyzed the data: FZ XH YY XK MLM DC. Contributed reagents/materials/analysis tools: GB SS. Wrote the paper: YX XH MLM DC DHK.

References

1. Nangaku M, Eckardt KU (2007) Hypoxia and the HIF system in kidney disease. *J Mol Med* 85: 1325–1330.
2. Heyman SN, Khamaisi M, Rosen S, Rosenberger C (2008) Renal Parenchymal Hypoxia, Hypoxia Response and the Progression of Chronic Kidney Disease. *Am J Nephrol* 28: 998–1006.
3. Martínez-Ballarín E, Pié J, Martínez-Berganza A (1986) Effects of chronic hypoxia on kidney function. *Rev Esp Fisiol* 42: 319–322.
4. White NMA, Yousef GM (2010) MicroRNAs: exploring a new dimension in the pathogenesis of kidney cancer. *BMC Med* 8: 65.
5. Liang M, Liu Y, Mladinov D, Cowley AW, Trivedi H Jr et al. (2009) MicroRNA: a new frontier in kidney and blood pressure research. *Am J Physiol Renal Physiol* 297: F553–F558.
6. Ma MC, Qian H, Ghassemi F, Zhao P, Xia Y (2005) Oxygen sensitive delta-opioid receptor-regulated survival and death signals: Novel insights into neuronal preconditioning and protection. *J Biol Chem* 280: 16208–16218.
7. Zhang J, Qian H, Zhao P, Hong SS, Xia Y (2006) Rapid hypoxia preconditioning protects cortical neurons from glutamate toxicity through delta-Opioid Receptor. *Stroke* 37: 1094–1099.
8. Feng Y, He X, Yang Y, Chen J, Yin K, et al. (2011) Effect of delta-opioid receptor over-expression on cortical expression of GABAA receptor $\alpha 1$ -subunit in hypoxia. *Chin J Physiol* 54: 118–123.
9. Zhang JH, Haddad GG, Xia Y (2000) δ -, but not μ - and κ -, opioid receptor activation protects neocortical neurons from glutamate-induced excitotoxic injury. *Brain Res* 885: 143–153.
10. Kang X, Chao D, Gu Q, Ding G, Wang Y, et al. (2009) Delta-opioid receptor activation decreases anoxic Na^+ influx and K^+ leakage via sodium channel regulation. *Cell Mol Life Sci* 66: 3505–3516.
11. Yang Y, Xia X, Zhang Y, Wang Q, Li L, et al. (2009) Delta-opioid receptor activation attenuates oxidative injury in the ischemic rat brain. *BMC Biol* 7: 55.
12. Chao D, Donnelly DF, Feng Y, Bazyz-Asaad A, Xia Y (2007) Cortical delta-opioid receptors potentiate K^+ homeostasis during anoxia and oxygen-glucose deprivation. *J Cereb Blood Flow Metab* 27: 356–368.
13. Chao D, Bazyz-Asaad A, Gianfranco B, Xia Y (2007) Delta-, but not μ -, opioid receptor stabilizes K^+ homeostasis by reducing Ca^{2+} influx in the cortex during acute hypoxia. *J Cell Physiol* 212: 60–67.
14. Chao D, Bazyz-Asaad A, Balboni G, Salvadori S, Xia Y (2008) Activation of DOR attenuates anoxic K^+ derangement via inhibition of Na^+ entry in mouse cortex. *Cereb Cortex* 18: 2217–2227.
15. Chao D, Balboni G, Lazarus LH, Salvadori S, Xia Y (2009) Na^+ mechanism of delta-opioid receptor induced protection from anoxic K^+ leakage in the cortex. *Cell Mol Life Sci* 66: 1105–1115.
16. Chao D, Xia Y (2010) Ionic storm in hypoxic/ischemic stress: Can opioid receptors subside it? *Prog Neurobiol* 90: 439–470.
17. Chao D, He X, Yang Y, Bazyz-Asaad A, Lazarus LH, et al. (2012) DOR activation inhibits anoxic/ischemic Na^+ influx through Na^+ channels via PKC mechanisms in the cortex. *Exp Neurol* 236: 228–239.
18. Patel HH, Head BP, Petersen HN, Niesman IR, Huang D, et al. (2006) Protection of adult rat cardiac myocytes from ischemic cell death: role of caveolar microdomains and delta-opioid receptors. *Am J Physiol Heart Circ Physiol* 291: H344–H350.
19. Borlongan CV, Wang Y, Su TP (2004) Delta opioid peptide (D-Ala 2, D-Leu 5) enkephalin: linking hibernation and neuroprotection. *Front Biosci* 9: 3392–3398.
20. Wu Q, Hwang CK, Zheng H, Wagley Y, Lin HY, et al. (2013) MicroRNA 339 down-regulates μ -opioid receptor at the post-transcriptional level in response to opioid treatment. *FASEB J* 27: 522–535.
21. He Y, Yang C, Kirkmire CM, Wang ZJ (2010) Regulation of opioid tolerance by let-7 family microRNA targeting the μ opioid receptor. *J Neurosci* 30: 10251–10258.
22. Zheng H, Chu J, Zeng Y, Loh HH, Law PY (2010) Yin Yang 1 phosphorylation contributes to the differential effects of μ -opioid receptor agonists on microRNA-190 expression. *J Biol Chem* 285: 21994–22002.
23. Cascio S, D'Andrea A, Ferla R, Surmacz E, Gulotta E, et al. (2010) miR-20b modulates VEGF expression by targeting HIF-1 α and STAT3 in MCF-7 breast cancer cells. *J Cell Physiol* 224: 242–249.
24. Lei Z, Li B, Yang Z, Fang H, Zhang GM, et al. (2009) Regulation of HIF-1 α and VEGF by miR-20b tunes tumor cells to adapt to the alteration of oxygen concentration. *PLoS One* 4: e7629.
25. Gonsalves CS, Kalra VK (2010) Hypoxia-mediated expression of 5-lipoxygenase-activating protein involves HIF-1 α and NF- κ B and microRNAs 135a and 199a-5p. *J Immunol* 184: 3878–3888.
26. Olaru AV, Selaru FM, Mori Y, Vazquez C, David S, et al. (2011) Dynamic changes in the expression of MicroRNA-31 during inflammatory bowel disease-associated neoplastic transformation. *Inflamm Bowel Dis* 17: 221–231.
27. Polytarchou C, Iliopoulos D, HatziaPOSTOLOU M, Kottakis F, Maroulakou I, et al. (2011) Akt2 regulates all Akt isoforms and promotes resistance to hypoxia through induction of miR-21 upon oxygen deprivation. *Cancer Res* 71: 4720–4731.
28. Xiong Y, Fang JH, Yun JP, Yang J, Zhang Y, et al. (2010) Effects of microRNA-29 on apoptosis, tumorigenicity, and prognosis of hepatocellular carcinoma. *Hepatology* 51: 836–845.
29. Luna C, Li G, Qiu J, Epstein DL, Gonzalez P (2009) Role of miR-29b on the regulation of the extracellular matrix in human trabecular meshwork cells under chronic oxidative stress. *Mol Vis* 15: 2488–2497.
30. Hulsmans M, De Keyzer D, Holvoet P (2011) MicroRNAs regulating oxidative stress and inflammation in relation to obesity and atherosclerosis. *FASEB J* 25: 2515–2527.
31. Kriegl AJ, Liu Y, Fang Y, Ding X, Liang M (2012) The miR-29 family: genomics, cell biology, and relevance to renal and cardiovascular injury. *Physiol Genomics* 44: 237–244.
32. Cui Y, Su W-Y, Xing J, Wang Y-C, Wang P, et al. (2011) MiR-29a inhibits cell proliferation and induces cell cycle arrest through the downregulation of p42.3 in human gastric cancer. *PLoS One* 6: e25872.
33. Ye Y, Perez-Polo JR, Qian J, Birnbaum Y (2011) The role of microRNA in modulating myocardial ischemia-reperfusion injury. *Physiol Genomics* 43: 534–542.
34. Tang Y, Banan A, Forsyth CB, Fields JZ, Lau CK, et al. (2008) Effect of alcohol on miR-212 expression in intestinal epithelial cells and its potential role in alcoholic liver disease. *Alcohol Clin Exp Res* 32: 355–364.
35. Hempel C, Combes V, Hunt NH, Kurtzhals JA, Grau GE (2011) CNS hypoxia is more pronounced in murine cerebral than noncerebral malaria and is reversed by erythropoietin. *Am J Pathol* 179: 1939–1950.
36. Kato S, Aoyama M, Kakita H, Hida H, Kato I, et al. (2011) Endogenous erythropoietin from astrocyte protects the oligodendrocyte precursor cell against hypoxic and reoxygenation injury. *J Neurosci Res* 89: 1566–1574.
37. Li J, Lu Y, Zhang J, Kang H, Qin Z, et al. (2010) PI4KIIalpha is a novel regulator of tumor growth by its action on angiogenesis and HIF-1 α regulation. *Oncogene* 29: 2550–2559.
38. Saito A, Miyachi N, Hashimoto T, Karasawa T, Han GD, et al. (2011) Neurexin-1, a presynaptic adhesion molecule, localizes at the slit diaphragm of the glomerular podocytes in kidneys. *Am J Physiol Regul Integr Comp Physiol* 300: R340–R348.
39. Saito A, Miyachi N, Hashimoto T, Karasawa T, Han GD, et al. (2011) Neurexin-1, a presynaptic adhesion molecule, localizes at the slit diaphragm of the glomerular podocytes in kidneys. *Am J Physiol Regul Integr Comp Physiol* 300: R340–R348.
40. Bondeva T, Ruster C, Franke S, Hammerschmid E, Klagsbrun M, et al. (2009) Advanced glycation end-products suppress neuropilin-1 expression in podocytes. *Kidney Int* 75: 605–616.
41. Xia Y, Eisenman D, Haddad GG (1993) Sulfonylurea receptor expression in rat brain: effect of chronic hypoxia during development. *Pediatric Res* 34: 634–641.
42. Xia Y, Warshaw JB, Haddad GG (1995) Chronic hypoxia causes opposite effects on glucose transporter 1 mRNA in mature versus immature rat brain. *Brain Res* 675: 224–230.
43. Xia Y, Warshaw JB, Haddad GG (1997) Effect of chronic hypoxia on glucose transporters in heart and skeletal muscle of immature and adult rats. *Am J Physiol* 273: R1734–R1741.
44. Bodnar RJ (1998) Recent advances in the understanding of the effects of opioid agents on feeding and appetite. *Expert Opin Investig Drugs* 7: 485–497.
45. Israel Y, Kandov Y, Khaimova E, Kest A, Lewis SR, et al. (2005) NPY-induced feeding: pharmacological characterization using selective opioid antagonists and antisense probes in rats. *Peptides* 26: 1167–1175.
46. Xia Y, Haddad GG (1994) Effect of prolonged O₂ deprivation on Na^+ channels: differential regulation in adult versus fetal rat brain. *Neuroscience* 94: 1231–1243.
47. Liu CG, Calin GA, Meloon B, Gamlie N, Sevignani C, et al. (2004) An oligonucleotide microchip for genome-wide microRNA profiling in human and mouse tissues. *Proc Natl Acad Sci U S A* 101: 9740–9744.
48. Thomson JM, Parker J, Perou CM, Hammond SM (2004) A custom microarray platform for analysis of microRNA gene expression. *Nat Methods* 1: 47–53.
49. Tusher VG, Tibshirani R, Chu G (2001) Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci USA* 98: 5116–5121.
50. Calin GA, Croce CM (2006) MicroRNA signatures in human cancers. *Nat Rev Cancer* 6: 857–866.
51. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, et al. (2005) Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res* 33: e179.
52. Tang F, Hajkova P, Barton SC, Lao K, Surani MA (2006) MicroRNA expression profiling of single whole embryonic stem cells. *Nucleic Acids Res* 34: e9.