

Electrochemical Synthesis of Unnatural Amino Acids Embedding 5- and 6-Membered Heteroaromatics

Elena Bombonato, Valerio Fasano, Daniel Pecorari, Luca Fornasari, Francesco Castagnini, Massimo Marcaccio, and Paolo Ronchi*



Cite This: *ACS Omega* 2024, 9, 13081–13085




Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information




ABSTRACT: Using a commercially available potentiostat, the electrochemical synthesis of unnatural amino acids bearing heteroaromatics on the lateral chain has been accomplished. This strategy exploits the side-chain decarboxylative arylation of aspartic/glutamic acid, a reaction that becomes challenging with electron-rich coupling partners such as 5- and 6-membered heteroaromatics. These rings are underrepresented in unnatural amino acids, therefore allowing a wider exploration of the chemical space, given the abundance of the aryl bromides employable in this reaction.

1. INTRODUCTION


The past decade has witnessed spectacular progress in the field of unnatural amino acids (UAAs), molecular entities widely employed in proteomics, protein engineering, and peptidomimetics.^{1–4} Many methodologies have been devised for the *de novo* synthesis of UAAs.^{5–7} An alternative approach is instead based on the lateral chain modification of natural amino acids, a useful strategy to retain the α -stereocenter. This is particularly true for aspartic acid (Asp) and glutamic acid (Glu), whose side-chain carboxylic group offers a strategic handle for derivatizations, especially those based on radical decarboxylations of the corresponding redox-active esters (RAEs).^{8–11} However, side-chain decarboxylation of these amino acids is usually more challenging than decarboxylation of α -CO₂H since only the latter provides a stabilized intermediate, which is an α -amino radical (Scheme 1).¹² While many radical partners (e.g., alkenes, alkynes, CO₂) have been engaged in lateral-chain decarboxylations, decarboxylative arylation of these amino acid residues remains elusive, with only a few examples reported in the literature.^{13–16} Indeed, decarboxylative arylations are limited by H atom abstraction or dimerization of the transient radicals as well as by unproductive N–O bond heterolysis. This limitation is further hampered by the fact that electron-rich heteroaromatics are less suitable as coupling partners in these arylations.¹⁶ Not surprisingly, side-chain decarboxylative arylations of Asp/Glu have not been reported using N-, S-, or O-containing 5-

Scheme 1. State-Of-The-Art Decarboxylation of Asp and Glu for the Synthesis of β/γ -Amino Acids (A) or Unnatural Amino Acids (B). NPHI = *N*-Hydroxyphthalimide


A) α -decarboxylative arylation (well-established)



B) Side-chain decarboxylative arylation (less explored)



Challenging e-rich aromatics in side-chain decarboxylative arylations



Received: November 29, 2023

Revised: February 7, 2024

Accepted: February 21, 2024

Published: March 4, 2024

membered aromatics. Given that these heterocycles are ubiquitous in drugs,^{17–19} herein we report our efforts to develop lateral-chain decarboxylative heteroarylations of aspartic acid and glutamic acids by means of electrochemistry.


This enabling technology is underused in pharmaceutical companies; thus, this collaborative project, based on the design of novel scaffolds in medicinal chemistry,²⁰ is also aimed at the transfer of knowledge of electrochemistry from academia to industry. By exploiting the new opportunities offered by the renaissance of radical chemistry, our approach allows the one-pot creation of novel unnatural glutamic acid by means of C(sp²)–C(sp³) linkages with 5- and 6-membered aromatic heterocycles.

2. RESULTS AND DISCUSSION

Our work takes inspiration from the decarboxylative arylation of RAEs and halogenated aromatics recently reported by Baran and co-workers.¹⁶ The coupling of these reagents was achieved using substoichiometric amounts (20 mol %) of NiCl₂·6H₂O and 2,2'-bipyridine (2,2'-bpy) in the presence of the additive AgNO₃ (0.5 equiv) in dimethylformamide (DMF) under constant current electrolysis with a reticulated vitreous carbon cathode and a sacrificial magnesium anode. This methodology was successfully applied to the decarboxylation arylation of many carboxylic acids, including a couple of examples employing glutamic acid. Using 4-iodophenol, this reaction allowed the multigram-scale synthesis of homotyrosine in a 43% yield, a classic UAA usually synthesized via multistep syntheses, but no other examples with electron-rich halides (especially 5-membered aromatics) were further investigated. Building on this precedence, we set off to develop an electrochemical method that could allow the synthesis of UAAs containing the challenging 5- and 6-membered heteroaromatics coupling partners. This method should also be cheap and transferable to flow chemistry to increase its appeal to pharmaceutical companies. Our investigation thus started with the search for alternative electrochemical conditions for the coupling of RAE **1** (derived from Asp) and 4-bromobenzaldehyde, an electron-poor aromatic whose coupling should be easier compared to 5-membered heterocycles (Table 1).

Initially, toxic NiCl₂·6H₂O was replaced by safer NiBr₂·diglyme, while *N*-methyl-2-pyrrolidone (NMP) was preferred over DMF as the solvent. Under these conditions, UAA **2** was observed by quantitative UPLC in a 42% yield (entry 1), a value in agreement with the yield reported for homotyrosine by Baran and co-workers (54%).¹⁶ As reported also in their work, dimerization of the unstable primary radical accounts for the rest of the mass balance. Isolation of **2** required two purifications by flash chromatography, thus causing a reduced isolated yield (28%). Replacing the aryl bromide with 4-iodobenzaldehyde furnished **2** in a lower yield (entry 2), a result in apparent disagreement with the work of Baran and co-workers.¹⁶ We believe that the sluggish oxidative addition of these substrates and the different solubilities of AgBr vs AgCl could favor the aryl bromide over the aryl iodide depending on the substrate under investigation. Performing the reaction in the absence of the nickel catalyst or without AgNO₃ resulted in no productive coupling (entry 3 and 4, respectively), whereas replacing AgNO₃ with Fe(OAc)₂ gave **2** in a 20% UPLC yield (entry 5). This is particularly important since it highlights that silver can act both as *in situ* Ag-functionalized electrodes (as previously determined)²¹ and as a mediator, thus making it

Table 1. Optimization of Conditions for Direct Side-Chain Decarboxylative Arylation of RAE **1**




entry	modification	yield ^a
1	none ^b	42% (28% iso)
2	Arl instead of ArBr	28%
3	no NiBr ₂ ·diglyme	0%
4	no AgNO ₃	0%
5	Fe(OAc) ₂ instead of AgNO ₃	20%
6	(+)GC instead of (+)Mg	0%
7	(+)Al instead of (+)mg	40%
8	(–)Pt _{ceramic} , (–)Pt _{foil} , or (–)graphite instead of (–) RCV	0%
9	(–)Ni _{foam} instead of (–)RCV	20%
10	using (–)Ni _{foam} and (+)Al	39%

^aYield determined by UPLC with 1-chloro-trifluoromethylbenzene as an internal standard. ^bStandard conditions: 4-Br-C₆H₄-CHO (1.5 equiv), NiBr₂·diglyme (20 mol %), 2,2'-bpy (20 mol %), AgNO₃ (0.5 equiv), NMP (0.1 M).

replaceable by iron which does not form nanoparticles at the electrode. Attempts to replace the sacrificial magnesium anode with a glassy carbon (GC) electrode gave no reaction (entry 6), supporting the role of magnesium as a source of electrons. Notably, the magnesium sacrificial anode could be replaced with aluminum, which increases the sustainability of the process (entry 7). Attempts to replace the counter electrode with platinum- or graphite-based ones were unsuccessful (entry 8), therefore pointing out how a large surface area is key in these decarboxylative arylations. Nevertheless, the use of nickel foam as the counter electrode showed reactivity to some extent (entry 9). This, in combination with aluminum as the sacrificial anode, gave product **2** in a 39% yield (entry 10), thus providing a cheaper and safer alternative to previously reported working conditions. Based on the optimization study and in analogy with previous results,^{16,22} a nickel-based catalytic mechanism system is believed to be responsible for the C(sp²)–C(sp³) coupling, with the deposition of silver nanoparticles on the electrode having a key role in the reactivity. However, in addition to this established mechanism, a concomitant pathway involving silver or other additives as a mediator could occur too, as observed when performing the reaction with FeOAc and using brand-new electrodes.

With the optimized conditions in hand, the scope of UAAs was explored, performing the reactions under air and at room temperature using technical-grade solvents and a commercially available potentiostat (Scheme 2). (+)Mg/(–)RCV electrodes were chosen since they were slightly more performing than (+)Al/(–)Ni_{foam}, although these still represent a valid option for economical reasons. Moreover, aryl bromides were employed as coupling partners since their greater availability (relative to aryl iodides) would allow a wider exploration of the chemical space. First, other protective groups on the starting aspartic acid were evaluated. To our delight, Boc and Cbz could both be tolerated under the reaction conditions, thus providing isolated UAA **3** and **4** in yields of 57 and 36%, respectively. Despite the higher yield obtained with the Boc group, the exploration of the scope was continued using Fmoc-

Scheme 2. Synthesis of UAAs by Means of Electrochemical Lateral-Chain Decarboxylative Arylations^a

^aIn brackets isolated yield after two purifications by flash chromatography.

or Cbz-protected starting materials since this would give access to orthogonally protected UAAs. Second, electron-poor (hetero)aromatics other than benzaldehyde were also amenable to this chemistry, including pharmaceutically relevant diazines (products 5–9). Notably, the stereocenter remains intact during this decarboxylative arylation, as demonstrated by the chiral UPLC of compound 7 (see Supporting Information). This is quite relevant since alternative approaches starting from dehydroalanine would furnish a racemic product.²³ Moving toward electron-rich benzenes and 5-membered heterocycles, a slightly lower reactivity was observed, in agreement with the reduced ability of these aromatics to engage in decarboxylative arylations. Yet, a wide range of heterocycles could be obtained in one-pot using the methodology described. For instance, thiophene-containing UAAs 11 and 12, potentially useful as fluorescent chemosensors,²⁴ were obtained in up to 30% yields. Thiazole, another ubiquitous 5-membered ring in medicinal chemistry, could be embedded too, as reported for UAAs 13–16. Interestingly, the relative position of the nitrogen atom within the structure can dramatically impact the outcome, as observed when comparing

regioisomeric UAAs 13 and 14. This could be attributed to the different stereoelectronic properties of the starting methylthiazole bromide, which affect the catalytic cycle (e.g., irreversible coordination to nickel).²⁵ Other 5-membered heterocycles incorporated for the first time in a UAA were *N*-methyl-pyrazole (unsuccessful in previous studies),¹⁶ thiadiazole, and oxadiazole (entries 17–20), with the last three representing examples of an aromatic ring containing three heteroatoms. Isothiazole- and triazole-based coupling partners were instead unsuccessful in the decarboxylative arylation, with the mass balance being given by the dimerization product of the unstable primary radical. The decarboxylative arylation could also be extended to glutamic acid, thus obtaining UAAs 21–24, bearing in the side-chain benzaldehyde (33%), 2-methylthiazole (50%), *N*-methyl-pyrazole (27%), and thiadiazole (27%).

Using an RAE derived from a dipeptide, UAA 25 was isolated in a 36% yield, thus opening possibilities for late-stage functionalization of small proteins.²⁶ Single-crystal X-ray crystallography of 25 confirmed its structure, hence highlighting that the developed method is a useful strategy to retain

the α -stereocenter. Finally, using the six-reactor carousel, this methodology can be scaled up to 3 g of starting material in one batch, as demonstrated with compound **13** isolated in 1.2 g (49% yield).

3. CONCLUSIONS

In conclusion, the synthesis of UAAs by means of side-chain decarboxylation of aspartic/glutamic acid has been presented. This simple protocol, requiring short reaction times and working at room temperature, allowed the incorporation of challenging electron-rich heterocycles on the lateral chain of novel UAAs, thus providing a more intuitive retrosynthesis of high-value targets. These aromatics, especially 5-membered ones, are underrepresented in previously reported UAAs, so novel building blocks are now available for use in proteomics. Finally, other relevant features of this methodology are the use of readily available aryl bromides as well as the possibility to use cheaper variants based on (+)Al/(-)Ni_{foam} electrodes.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsomega.3c09357>.

Procedures including preparation of substrates, optimization of reaction conditions, characterization data, and NMR spectra (PDF) (CIF)

■ AUTHOR INFORMATION

Corresponding Author

Paolo Ronchi – Medicinal Chemistry and Drug Design Technologies Department, Global Research and Preclinical Development, Chiesi Farmaceutici S.p.A, Parma 43122, Italy; orcid.org/0000-0001-5496-5068; Email: p.ronchi@chiesi.com

Authors

Elena Bombonato – Department of Chemistry “Giacomo Ciamician”, Università di Bologna, Bologna 40126, Italy
Valerio Fasano – Department of Chemistry, Università degli Studi di Milano, Milano 20133, Italy; orcid.org/0000-0003-1819-4483
Daniel Pecorari – Analytics and Early Formulations Department, Global Research and Preclinical Development, Chiesi Farmaceutici S.p.A, Parma 43122, Italy
Luca Fornasari – Analytics and Early Formulations Department, Global Research and Preclinical Development, Chiesi Farmaceutici S.p.A, Parma 43122, Italy
Francesco Castagnini – Department of Food and Drug Sciences, University of Parma, Parma 43124, Italy
Massimo Marcaccio – Department of Chemistry “Giacomo Ciamician”, Università di Bologna, Bologna 40126, Italy; orcid.org/0000-0001-9032-0742

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acsomega.3c09357>

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Chiesi Group for financial support. Dr Fabio Rancati, Dr Maurizio Delcanale, and Dr Daniele Fiorito are acknowledged for support and useful discussion.

■ ABBREVIATIONS

UAA	unnatural amino acid
Asp	aspartic acid
Glu	glutamic acid
RAE	redox-active ester
DMF	dimethylformamide
RCV	reticulated vitreous carbon
NHPI	<i>N</i> -hydroxyphthalimide
NMP	<i>N</i> -methyl-2-pyrrolidone
UPLC	ultraperformance liquid chromatography
GC	glassy carbon
Ni _{foam}	Nickel foam

■ REFERENCES

- (1) Adhikari, A.; Bhattarai, B. R.; Aryal, A.; Thapa, N.; Kc, P.; Adhikari, A.; Maharjan, S.; Chanda, P. B.; Regmi, B. P.; Parajuli, N. Reprogramming natural proteins using unnatural amino acids. *RSC Adv.* **2021**, *11*, 38126–38145.
- (2) Drienovská, I.; Roelfes, G. Expanding the enzyme universe with genetically encoded unnatural amino acids. *Nat. Catal.* **2020**, *3*, 193–202.
- (3) Wang, X.; Yang, X.; Wang, Q.; Meng, D. Unnatural amino acids: promising implications for the development of new antimicrobial peptides. *Crit. Rev. Microbiol.* **2023**, *49*, 231–255.
- (4) Lugtenburg, T.; Gran-Scheuch, A.; Drienovská, I. Non-canonical amino acids as a tool for the thermal stabilization of enzymes. *Protein Eng. Des. Sel.* **2023**, *36*, 1–10.
- (5) Narancic, T.; Almaboub, S. A.; O'Connor, K. E. Unnatural amino acids: production and biotechnological potential. *World J. Microbiol. Biotechnol.* **2019**, *35*, 67.
- (6) Hickey, J. L.; Sindhikara, D.; Zultanski, S. L.; Schultz, D. M. Beyond 20 in the 21st Century: Prospects and Challenges of Non-canonical Amino Acids in Peptide Drug Discovery. *ACS Med. Chem. Lett.* **2023**, *14*, 557–565.
- (7) Yousif, A. M.; Colarusso, S.; Bianchi, E. Katritzky Salts for the Synthesis of Unnatural Amino Acids and Late-Stage Functionalization of Peptides. *Eur. J. Org. Chem.* **2023**, *26*, No. e202201274.
- (8) Aguilar Troyano, F. J.; Merckens, K.; Anwar, K.; Gómez-Suárez, A. Radical-Based Synthesis and Modification of Amino Acids. *Angew. Chem., Int. Ed.* **2021**, *60*, 1098–1115.
- (9) Liu, Y. A.; Liao, X.; Chen, H. Recent Progress in Radical Decarboxylative Functionalizations Enabled by Transition-Metal (Ni, Cu, Fe, Co or Cr) Catalysis. *Synthesis* **2021**, *53*, 1–29.
- (10) Wadamoto, M. Method for preparing aromatic amino acid derivative. Chugai Seiyaku Kabushiki Kaisha. Tokyo. WO 2020/189540, 2020.
- (11) Laudadio, G.; Neigenfind, P.; Péter, Á.; Emmanuel, M. A.; Palkowitz, M. D.; Oderinde, M. S.; Sloane, J. L.; Gillman, K. W.; Ridge, D.; Bolduc, P.; Nicastrì, M.; Zhang, B.; Clementson, S.; Nasser Petersen, N.; Martín-Gago, P.; Mykhailiuk, P.; Schmidt, M. A.; Mandler, M. D.; Baran, P. S. Ni-Electrocatalytic Decarboxylative Arylation to Access Quaternary Centers. *ChemRxiv* **2023**.
- (12) King, T. A.; Mandrup Kandemir, J.; Walsh, S. J.; Spring, D. R. Photocatalytic methods for amino acid modification. *Chem. Soc. Rev.* **2021**, *50*, 39–57.
- (13) Toriyama, F.; Cornella, J.; Wimmer, L.; Chen, T. G.; Dixon, D. D.; Creech, G.; Baran, P. S. Redox-Active Esters in Fe-Catalyzed C–C Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 11132–11135.
- (14) Huihui, K. M.; Caputo, J. A.; Melchor, Z.; Olivares, A. M.; Spiewak, A. M.; Johnson, K. A.; DiBenedetto, T. A.; Kim, S.; Ackerman, L. K.; Weix, D. J. Decarboxylative Cross-Electrophile

Coupling of N-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc.* **2016**, *138*, 5016–5019.

(15) Wang, J.; Qin, T.; Chen, T. G.; Wimmer, L.; Edwards, J. T.; Cornella, J.; Vokits, B.; Shaw, S. A.; Baran, P. S. Nickel-Catalyzed Cross-Coupling of Redox-Active Esters with Boronic Acids. *Angew. Chem., Int. Ed.* **2016**, *55*, 9676–9679.

(16) Palkowitz, M. D.; Laudadio, G.; Kolb, S.; Choi, J.; Oderinde, M. S.; Ewing, T. E.; Bolduc, P. N.; Chen, T.; Zhang, H.; Cheng, P. T. W.; Zhang, B.; Mandler, M. D.; Blaszczak, V. D.; Richter, J. M.; Collins, M. R.; Schioldager, R. L.; Bravo, M.; Dhar, T. G. M.; Vokits, B.; Zhu, Y.; Echeverria, P. G.; Poss, M. A.; Shaw, S. A.; Clementson, S.; Petersen, N. N.; Mykhailiuk, P. K.; Baran, P. S. Overcoming Limitations in Decarboxylative Arylation via Ag-Ni Electrocatalysis. *J. Am. Chem. Soc.* **2022**, *144*, 17709–17720.

(17) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the structural diversity, substitution patterns, and frequency of nitrogen heterocycles among U.S. FDA approved pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274.

(18) Baumann, M.; Baxendale, I. R.; Ley, S. V.; Nikbin, N. An overview of the key routes to the best selling 5-membered ring heterocyclic pharmaceuticals. *Beilstein J. Org. Chem.* **2011**, *7*, 442–495.

(19) Sperry, J. D.; Wright, D. L. Furans, thiophenes and related heterocycles in drug discovery. *Curr. Opin. Drug Discov. Dev.* **2005**, *8*, 723–740.

(20) Failla, M.; Lombardo, G. W.; Orlando, P.; Fiorito, D.; Bombonato, E.; Ronchi, P.; Passarella, D.; Fasano, V. Late-Stage Functionalisation of Pyridine-Containing Bioactive Molecules: Recent Strategies and Perspectives. *Eur. J. Org. Chem.* **2023**, *26*, No. e202300074.

(21) Harwood, S. J.; Palkowitz, M. D.; Gannett, C. N.; Perez, P.; Yao, Z.; Sun, L.; Abruña, H. D.; Anderson, S. L.; Baran, P. S. Modular terpene synthesis enabled by mild electrochemical couplings. *Science* **2022**, *375*, 745–752.

(22) Lin, Q.; Diao, T. Mechanism of Ni-Catalyzed Reductive 1,2-Dicarbofunctionalization of Alkenes. *J. Am. Chem. Soc.* **2019**, *141*, 17937–17948.

(23) Peng, X.; Xu, K.; Zhang, Q.; Liu, L.; Tan, J. Dehydroalanine modification sees the light: a photochemical conjugate addition strategy. *Trends Chem.* **2022**, *4*, 643–657.

(24) Esteves, C. I. C.; Raposo, M. M. M.; Costa, S. P. G. Non-canonical amino acids bearing thiophene and bithiophene: synthesis by an Ugi multicomponent reaction and studies on ion recognition ability. *Amino Acids* **2017**, *49*, 921–930.

(25) Milner, P. J.; Yang, Y.; Buchwald, S. L. In-Depth Assessment of the Palladium-Catalyzed Fluorination of Five-Membered Heteroaryl Bromides. *Organometallics* **2015**, *34*, 4775–4780.

(26) Malins, L. R. Decarboxylative couplings as versatile tools for late-stage peptide modifications. *Pept. Sci.* **2018**, *110*, No. e24049.

NOTE ADDED AFTER ASAP PUBLICATION

This paper originally published ASAP on March 4, 2024. The TOC graphic was modified, and a new version reposted on March 5, 2024.