



Published in final edited form as:

Gene Ther. 2012 September ; 19(9): 877–885. doi:10.1038/gt.2011.175.

Hydrodynamic delivery of plasmid DNA encoding human Fc γ R-Ig dimers blocks immune-complex mediated inflammation in mice

Rangaiah Shashidharamurthy, Ph.D, Deepa Machiah, Ph.D, Erica N. Bozeman, B.E, Sanjay Srivatsan, B.S, Jaina Patel, B.S, Alice Cho, B.S, Joshy Jacob, Ph.D, and Periasamy Selvaraj, Ph.D*

Department of Pathology and Laboratory Medicine, Emory University School of Medicine, Atlanta, GA 30322. USA.

Abstract

Therapeutic use and function of recombinant molecules can be studied by the expression of foreign genes in mice. In this study, we have expressed human Fc γ receptor –Ig fusion molecules (Fc γ R-Igs) in mice by administering Fc γ R-Ig plasmid DNAs hydrodynamically and compared their effectiveness to purified molecules in blocking immune-complex (IC) mediated inflammation in mice. The concentration of hydrodynamically expressed Fc γ R-Igs (CD16A^F-Ig, CD32A^R-Ig and CD32A^H-Ig) reached a maximum of 130 μ g/ml of blood within 24 h after plasmid DNA administration. The *in vivo* half-life of Fc γ R-Igs was found to be 9-16 days and Western blot analysis showed that the Fc γ R-Igs were expressed as a homodimer. The hydrodynamically expressed Fc γ R-Igs blocked 50-80% of IC-mediated inflammation up to 3 days in a reverse passive Arthus reaction model. Comparative analysis with purified molecules showed that hydrodynamically expressed Fc γ R-Igs are more efficient than purified molecules in blocking IC-mediated inflammation and had a higher half-life. In summary, these results suggest that the administration of a plasmid vector with a Fc γ R-Ig gene can be used to study the consequences of blocking IC-binding to Fc γ Rs during the development of inflammatory diseases. This approach may have potential therapeutic value in treating IC-mediated inflammatory autoimmune diseases such as lupus, arthritis and autoimmune vasculitis.

Keywords

Hydrodynamic delivery; Plasmid DNA; Fc γ receptors; Inflammation

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

*Address for Correspondence: Periasamy Selvaraj, PhD. Department of Pathology Emory University School of Medicine 7309 Woodruff Memorial Building 101 Woodruff Circle Atlanta, GA 30322 Phone: 404-727-5929 pselvar@emory.edu.

Conflict of interest: The authors declare no competing financial interests.

INTRODUCTION

The *in vivo* expression of a foreign gene in animal models is of great interest because it not only provides an opportunity to study the structure/function of a protein, but also has therapeutic efficacy in treating various pathological disorders. *In vivo* transgene expression requires efficient delivery of specific genes into the cells. Currently, viral and non-viral vectors are the two predominant gene delivery systems being used. Recently, it has been shown that a significant amount of protein can be expressed *in vivo* by rapidly injecting plasmid DNA in a large volume through the tail vein by a process called hydrodynamic-based gene delivery¹⁻⁴. Though the mechanism of this hydrodynamic-based *in vivo* gene expression is not clearly understood, it has been suggested that the rapid injection of a large volume of plasmid DNA solution causes shearing forces on the hepatocytes. These forces induce transient pore formation in the plasma membrane facilitating the direct entry of the plasmid DNA into the hepatocytes' cytosol resulting in a high level of transient *in vivo* gene expression^{1,2,5-7}. Since the discovery of this hydrodynamic gene delivery technique in the 1990s, it has been reported as an effective method of gene delivery in experimental animal models. The introduction of exogenous genes by the hydrodynamic method using plasmid DNA has many advantages such as ease of preparation of large quantities of DNA in a short period of time and stability. Recently, several studies have shown the hydrodynamic-based *in vivo* expression of several protein molecules (CTLA4-Ig, IL22-Ig, IL10-Ig, CD40-Ig, fetal liver kinase-1, DNA cancer vaccine, hFlex-TRAIL) and their role in various disease conditions such as experimental autoimmune myocarditis, allergic encephalomyelitis, systemic lupus erythematosus, collagen-induced arthritis, nephritis, and cancer⁸⁻¹⁷.

In this report, we have investigated whether hydrodynamic-based delivery of Fc γ R genes results in a functional product that can block immune-complex (IC)-mediated inflammation. The receptors (Fc γ R) for the Fc domain of IgG molecules play a vital role in IC-mediated autoimmune diseases. Inflammatory cells, such as neutrophils, monocytes, and NK cells, express three types of Fc γ R¹⁸⁻²². Fc γ RI (CD64) is a high affinity receptor for monomeric IgG whereas Fc γ RII (CD32) and Fc γ RIII (CD16) are low affinity receptors for monomeric IgG; however, all three bind stably to ICs. Both *in vivo* and *in vitro* studies from various laboratories have shown that interaction of Fc γ Rs expressed on inflammatory cells with antibody-coated target cells/tissues is a key event in the destruction of antibody coated tissues through antibody dependent cellular cytotoxicity (ADCC) and phagocytosis²³⁻³¹, which leads to the development of various autoimmune diseases^{28,32-37}. During the development of autoimmune diseases such as arthritis, systemic lupus erythematosus and autoimmune vasculitis, autoantibodies bind to the antigen expressed on cells and form ICs. These ICs bind to inflammatory cells through Fc γ Rs leading to chronic inflammation and destruction of the target cells. Therefore, blocking the interaction of pathogenic ICs with the cell surface Fc γ Rs expressed on inflammatory cells using recombinant Fc γ R-Igs could be a potential therapeutic approach. We have previously demonstrated that the administration of a purified dimeric form of a low affinity Fc γ R (CD16A-Ig) can be successfully employed to treat IC-mediated acute inflammation in mice³⁸. In this report, we have hydrodynamically expressed human low affinity Fc γ R-Igs *in vivo* and studied their effectiveness in blocking IC-mediated inflammation in a murine model. We show that *in vivo* expressed recombinant

Fc γ R-Ig molecules are secreted in high concentrations and are sustained for longer periods of time in circulation compared to the administered purified Fc γ R-Ig. These molecules are also as effective as purified Fc γ R-Igs in blocking the interaction of ICs with inflammatory cells, thus preventing inflammation *in vivo* for a relatively longer period of time.

RESULTS

Purified human Fc γ R-Igs block IC-binding to mouse Fc γ Rs *in vitro* and antibody-mediated inflammation *in vivo* in mice

Previously, we have shown that the purified human CD16A-Ig molecule, when administered in mice, is capable of blocking IC-induced inflammation in a reverse passive Arthus (RPA) model³⁸. In this study we have made additional human Fc γ R-Ig molecules. Prior to testing the anti-inflammatory effect of human Fc γ R-Ig molecules in mice, we assessed the ability of purified human Fc γ R-Ig molecules (CD16A^V-Ig, CD16A^F-Ig, CD32A^R-Ig and CD32A^H-Ig) to compete and cross-block IC-binding to mouse Fc γ Rs *in vitro*. We carried out a soluble IC (sIC)-binding assay using the mouse macrophage cell line (P388D1) as described earlier³⁸. P388D1 cells express all four types of mouse Fc γ Rs: CD16A, CD32B, CD64 and Fc γ RIV. As shown in Figure 1a, all of the hFc γ R-Ig molecules, except hCD32A^R-Ig, inhibited sIC binding to mouse Fc γ Rs expressed on P388D1 cells in a dose dependent fashion. More than 70% inhibition was achieved when P388D1 cells were co-incubated with 25 μ g/ml of hCD16A^F-Ig, hCD16A^V-Ig and hCD32A^H-Ig, whereas, hCD32A^R-Ig was not able to compete with mouse Fc γ Rs at the same concentration. The results with purified hCD16A^F-Ig, and hCD32A^R-Ig are consistent with our previous studies³⁸.

The *in vivo* efficacy of purified human Fc γ R-Ig molecules was analyzed using RPA an acute antibody-mediated inflammation in a murine model³⁸. In this model, inflammation is initiated by the formation of antigen-antibody complexes at the antibody-injected site. To induce inflammation, rabbit anti-chicken ovalbumin (anti-Ova) IgG was injected intradermally on the dorsal side of the mouse skin and chicken ovalbumin (Ova) along with 1% Evan's blue (blue dye) was injected intravenously. Extravasation of inflammatory cells and vascular permeability during the inflammation at the antibody-injected site can be visualized upon leakage of the blue dye^{38,39}. Before the induction of RPA, hFc γ R-Ig molecules (50 μ g/ml of mouse blood) were administered intravenously into a separate group of mice (n=3) to study the effect of purified hFc γ R-Ig molecules. As shown in Figure 1b, hCD16A^F-Ig, hCD16A^V-Ig and hCD32A^H-Ig inhibited more than 70% inflammation as measured by the intensity of the Evan's blue extravasation (Fig. 1c); whereas under similar conditions, hCD32A^R-Ig did not inhibit inflammation (Fig. 1b). The lack of inhibition by purified hCD32A^R-Ig may be due to the difference in affinity of human Fc γ R-Igs towards rabbit IgG binding or because hCD32A^R-Ig might have partially denatured during the purification process. As specificity controls, mice treated with 2.4G2 antibody, a mAb to mouse Fc γ Rs (CD16/32), did not cause RPA (Fig. 1b).

***In vivo* expression of human Fc γ R-Ig molecules in mice by the hydrodynamic-based method has a longer half-life**

Next, we determined the level and kinetics of expression of human Fc γ R-Ig molecules *in vivo*. As shown in Figure 2, all of the hydrodynamically expressed hFc γ R-Ig molecules in mice reached maximum expression by 24h. The concentration of hFc γ R-Ig in circulation was found to be around 130 μ g/ml of blood. The *in vivo* half-life was 9 days for hCD16A^F-Ig; whereas for the hCD32A-Ig molecules, the half-life was estimated to be 16 days. hCD16A^F-Ig was detected up to 17 days in circulation, while, hCD32A-Ig molecules were detected until day 24. We reported previously that the *in vivo* half-life of the hFc γ R-Ig molecules is 5 days when administered as purified molecules³⁸. These results suggest that the *in vivo* expression of a foreign gene in mice by the hydrodynamic-based method increases the *in vivo* half-life of the foreign protein when compared to administration of purified proteins. At present the reason for increased half-life of hydrodynamically expressed Fc γ R-Igs is not clear. However it is possible that the purification process may have made the molecules more susceptible to *in vivo* degradation.

Hydrodynamically expressed human Fc γ R-Ig molecules block antibody-mediated inflammation in mice

We then determined whether hydrodynamically expressed hFc γ R-Ig can block antibody-mediated inflammation in a murine RPA model. In these experiments, a group of mice (n=12 for each molecule) were injected with plasmid DNA of hFc γ R-Ig as described under Materials and Methods. DNA was injected on day 0 and at different time intervals, and a group of mice (n=3) were used to conduct the RPA for each molecule. We have chosen different time points for each molecule to conduct the RPA since the hydrodynamically expressed hCD16A-Ig and hCD32A-Ig molecules differ in their *in vivo* half-life. For hCD16A^F-Ig, the RPA was carried out on days 1, 3, 7 and 13. As shown in Figure 3a, *in vivo* expressed hCD16A^F-Ig was able to significantly (P=0.001) reduce antibody-mediated inflammation up to 3 days. As measured by the intensity of the Evan's blue extravasation, hCD16A^F-Ig was able to block inflammation by 83% and 77% on days 1 and 3, respectively, while on day 7 it was only able to block 20% (Fig. 3a; left panel). These results suggest that *in vivo* expressed hCD16A^F-Ig can block antibody-mediated inflammation up to 3 days and becomes ineffective by day 7 (Fig. 3a; right panel bar graph). For hCD32A^R-Ig and hCD32A^H-Ig DNA injected animals, the RPA was conducted on days 1, 3, 9 and 16. Interestingly, as shown in Figure 3b, the *in vivo* expressed hCD32A^R-Ig was able to block antibody mediated inflammation completely on day 1, but the blocking efficiency as measured by the intensity of the blue dye began to decline by day 3 and was completely ineffective by day 9 (Fig. 3b; right panel). Most notably, hydrodynamically expressed hCD32A^R-Ig blocks IC-mediated inflammation in mice, while the purified molecule cannot (compare Fig. 1 and Fig. 3). The *in vivo* expressed hCD32A^H-Ig was able to block antibody-mediated inflammation completely up to 9 days (Fig. 3c). About 50% of RPA was blocked by hCD32A^H-Ig even on day 9 and became ineffective on day 16 as indicated by the decrease in the intensity of the blue dye (Fig. 3c; right panel). As a specificity control, mice were treated with 2.4G2 mAb, which completely blocked RPA on day 3. The untreated group of mice (n=3) injected with rabbit anti-Ova antibody alone served as the positive

control. These results suggest that hydrodynamically expressed hFc γ R-Ig molecules are capable of competing with mFc γ Rs and can effectively block IC binding.

Functionality of hydrodynamically expressed hFc γ R-Ig molecules

To determine whether the level of expression correlates with the ability to block RPA, plasma was collected from the mice used for RPA experiments before induction of RPA, and the concentration of human Fc γ R-Igs in the blood of the mice was determined using a sandwich ELISA. As shown in Figure 4 (upper panel), hCD16A^F-Ig was expressed at about 120 μ g/ml of blood up to day 3. About 50% of the hCD16A^F-Ig was cleared from circulation by day 7 and was not detectable in circulation on day 13. hCD32A^R-Ig and hCD32A^H-Ig were expressed at 160 and 120 μ g/ml of blood, respectively, within 24 h of hydrodynamic injection, (Fig. 4, middle and lower panel). The same concentration was maintained until day 9, and about 50% was cleared from circulation by day 16. Although the concentration of Fc γ R-Ig molecules present within the mice should be sufficient to block RPA, CD16A^F-Ig and CD32A^R-Ig were not able to block RPA after three and nine days of expression, respectively. This suggests that the level expression of hFc γ R-Ig does not correlate with the ability to block RPA when the molecules are in circulation more than 3 days. Therefore, we determined whether the hFc γ R-Ig molecules present in circulation are capable of binding IgG in *in vitro* assays.

ELISA plates were coated with 10 μ g/ml of rabbit IgG, and then 100 μ l of 1:100 diluted plasma was added to the wells. The plates were washed, and the bound hFc γ R-Ig was detected using HRP-conjugated anti-human Fc specific antibody. The mAb coated wells (CLB-Fcgran-1 for to human CD16A and IV.3 for CD32A) and the plasma from naïve mice served as specificity controls. As shown in Figure 5 (upper panel), *in vivo* expressed hCD16A^F-Ig bound to rabbit IgG efficiently on day 1 of its expression. On day 3, its binding efficiency to rabbit IgG was reduced by 50% and by day 7 it was no longer able to bind to rabbit IgG. The *in vivo* expressed hCD32A^R-Ig (Fig. 5: middle panel) bound to rabbit IgG poorly on day 1 of expression, and its binding efficiency was reduced by 50% on day 3. On day 9 it was not able to bind to rabbit IgG. The *in vivo* expressed hCD32A^H-Ig (Fig. 5: lower panel) bound efficiently to rabbit IgG on day 1 and retained up to 50% of its binding efficiency until day 16. Correlation of the level of expression (Fig. 4) and functional activity (Fig. 5 and Fig. 3) shows that hCD32A^H-Ig and hCD16A^F-Ig bind more efficiently to rabbit IgG when compared to hCD32A^R-Ig. These data suggest that dimeric Fc γ R-Ig molecules may become inactive or degraded partially after being present in circulation for several days, and the lack of functionality may vary from molecule to molecule. At present, the reason for the ineffectiveness of CD16A^F-Ig and CD32A^R-Ig after being in circulation for a few days is not clear.

Hydrodynamically expressed hFc γ R-Ig molecules are secreted as a homodimer

To determine the molecular nature and kinetics of degradation of circulating hFc γ R-Ig, we analyzed the plasma by Western blot. Previously we have reported that recombinant hFc γ R-Igs secreted by CHO-K1 cell transfectants are disulfide-linked homodimers^{38,40,41}. As shown in Figure 6a, on day 1, hCD16A^F-Ig is secreted as a disulfide-linked homodimer as evidenced by the presence of 120 kDa band under non-reducing conditions and the 70 kDa

band under reducing conditions. However, we also observed a band around 200 kDa in CD16A-Ig blots, while on day 3 this band had mostly disappeared. The reason for the secretion of this high molecular weight band is not clear; however it may be due to the aggregation of hCD16A^F-Ig. We did not observe this high molecular weight band when we purified the hCD16A^F-Ig molecule *in vitro*⁴⁰. On day 3, the molecules began to degrade, as suggested by the appearance of low molecular weight bands, and was completely degraded by day 13. These results are consistent with the detectable rabbit IgG binding activity (Fig. 5). The hCD32A^R-Ig (Fig. 6b) and hCD32A^H-Ig (Fig. 6c) molecules are also secreted as disulfide linked homodimers as seen by the 110 kDa band under non-reducing conditions and a 60 kDa band under reducing conditions. The intensity of the bands under reducing conditions was lower which may be due to the decreased reactivity of the antibodies with denatured molecules. Interestingly, hCD32A-Ig molecules are more stable *in vivo* than hCD16A-Ig (compare Fig. 6a, b and c) as suggested by its dimeric nature up to 16 days. In addition to the major bands, we observed minor low molecular weight bands, which may be due to different glycosylation or degradation products of the hCD32A-Ig molecules. These results suggest that the progressive loss of ability to block inflammation in RPA with time by the *in vivo* expressed Fc γ R-Ig molecules correlates with the *in vitro* rabbit IgG binding activity and is partly due to the degradation of the molecules in circulation.

Hepatocytes and antigen presenting cells (APCs) such as macrophages in the spleen are responsible for the hydrodynamic expression of hFc γ R-Ig

To investigate the specific organ and cell type involved in the high expression of hFc γ R-Ig molecules, a group of mice (n=3) were hydrodynamically injected with hCD32A^H-Ig plasmid DNA. Various organs were then harvested at different time points post DNA injection and subjected to immunohistochemistry. The expression of hCD32A^H-Ig was detected using the F(ab)[']2 fragment of goat anti-human Fc specific antibody conjugated with Cy-5. Consistent with previous reports^{1,42}, our results show that hepatocytes are the major cell type responsible for the expression of hFc γ R-Igs and maintained expression for up to 48h after plasmid DNA administration (Fig 7A). Interestingly, APCs such as macrophages in the red pulp region of the spleen also expressed the hCD32A^H-Ig molecules, but not the B- or T-cell zones in the white pulp region (Fig 7B). Maximum expression was reached after 24h and declined thereafter (Fig 7B). In contrast, expression of hCD32A^H-Ig was not found in other organs tested such as kidney, lung or heart (figure not shown). These results suggest that in addition to hepatocytes in the liver, APCs such as macrophages present in the red pulp area of the spleen are also responsible for the expression of hFc γ R-Ig molecules following the hydrodynamic delivery of plasmid DNA.

DISCUSSION

In the recent past, the significance of low affinity Fc γ Rs in autoimmune/inflammatory disease conditions has been well-documented⁴³⁻⁴⁷. The interaction of ICs with Fc γ Rs on inflammatory cells is a key event in the progression of IC-mediated diseases. Previously, we have shown that inflammation caused by the interaction of ICs with inflammatory cells can be blocked by administering purified Fc γ R-Ig molecules^{38,48,49}. In this report, we demonstrate that the hydrodynamic administration of plasmid DNAs encoding human low-

affinity Fc γ R-Igs is capable of blocking IC-mediated inflammation in a murine RPA model. The hFc γ R-Ig molecules were secreted in high concentrations in circulation and sustained for longer periods of time after hydrodynamic administration when compared to the administered purified molecules. Recently, we have reported that the purified hCD32A^R-Ig molecules, despite being active *in vitro*⁴¹, failed to block inflammation induced by RPA in mice³⁸. The reason for the ineffectiveness of the purified hCD32A^R-Ig molecule *in vivo* is not clear. However, it is possible that the purified molecules may not be able to compete with mouse Fc γ Rs for IC binding³⁸ or may have partially denatured during the purification process, thus leading to its ineffectiveness. Interestingly, in this report we observed that hydrodynamically expressed hCD32A^R-Ig molecules are effective in blocking antibody-induced inflammation in mice, suggesting that the hCD32A^R-Ig molecules may have been partially denatured during the purification process. These results also suggest that *in vivo* expressed molecules are more active when compared to purified molecules, indicating that the *in vivo* expression of a transgene by the hydrodynamic-based method is a more desirable method to study the function of a particular gene product *in vivo*. It has been shown that after hydrodynamic injection of plasmid DNA, a high expression of recombinant luciferase was found in the liver whereas a very low level of expression was observed in the spleen, kidney, lung and heart¹. In agreement with this, we observed that the hepatocytes are the major cell types responsible for the high level of expression of recombinant molecules. Interestingly, we also found that APCs present in the red pulp region of the spleen are also responsible for the expression of recombinant molecules and not T or B-cells present in the white pulp region (fig 7). In contrast, we did not observe the expression of recombinant molecules in the other organs tested such as kidney, lung and heart (data not shown). Taken together, these results suggest that the hepatocytes in the liver and APCs in the spleen are responsible for the expression of recombinant molecules when plasmid DNA is hydrodynamically injected in mice.

Recently, hydrodynamic-based *in vivo* expression of several Ig fusion protein molecules such as interleukins and cell adhesion molecules (CTLA4-Ig, IL22-Ig, IL10-Ig, Flex/TRIAL-Ig) were studied, and their role in various disease conditions was established^{8-10,15-17}. For instance, hydrodynamically expressed CTLA4-Ig dramatically reduced experimental autoimmune myocarditis by blocking the costimulatory signal from the T cells. The *in vivo* expression of CTLA4-Ig was approximately 2 μ g/ml of blood and was detectable up to 16 days¹⁰. In this report, for the first time, we have expressed hFc γ R-Ig molecules by injecting naked plasmid DNA in mice and showed that hydrodynamically expressed hFc γ R-Ig molecules have a longer half life compared to the purified molecules, remain as dimers in circulation, and are effective in blocking IC-mediated inflammation in a mouse model for an extended period of time. It has been shown that the *in vivo* half-life of purified monomeric hCD32A was only 30 min in mice²⁵. Whereas, recently we have shown that the *in vivo* half-life of purified, dimeric forms of Fc fusion hCD16A^F-Ig and hCD32A^R-Ig was 120h when administered intravenously to mice³⁸. In this report we observed that the hydrodynamically expressed Fc γ R-Ig molecules exhibited a longer half-life when compared to purified Fc γ R-Ig molecules. The reason for this prolonged half-life is not clear, however, this may be due to continuous production of the Fc γ R-Ig molecules from hepatocytes and macrophages present in the spleen, which took up the DNA during hydrodynamic injection.

Moreover it has been shown that the *in vivo* half-life of Fc fusion proteins is due to the presence of the Fc domain. The Fc domain binds to neonatal Fc receptors (FcRn) and protects the IgG from degradation by the endocytic machinery of cells⁵⁰. The Fc domain used in the present investigation is mutated to abolish the Fc γ R binding activity⁵¹⁻⁵³. Since both Fc γ Rs and FcRn bind to different sites of the Fc domain^{54,55}, the mutation that we introduced in the Fc domain of the Fc γ R-Ig molecules will not affect the FcRn binding activity, therefore potentially aiding in extending the *in vivo* half-life of hFc γ R-Ig molecules. It has been shown that hydrodynamically injected plasmid DNA can stay in the hepatocytes for upto 3 days after administration^{15,42}. Alternatively, hydrodynamically expressed Fc γ R-Igs may be in a less denatured state in circulation due to lack of exposure to harsh acidic conditions that are used for affinity purification of Fc γ R-Igs.

In conclusion, we have demonstrated that the hydrodynamic delivery of plasmid DNA encoding hFc γ R-Igs inhibited antibody mediated inflammation in mice. The hFc γ R-Ig molecules are expressed at high levels, intact and dimeric in nature, and sustained in circulation for an extended period of time. The data presented herein will be useful in studying the consequences of uncoupling IC binding to Fc γ Rs expressed on inflammatory cells during IC-mediated inflammatory/autoimmune diseases such as arthritis, systemic lupus erythematosus, and autoimmune vasculitis. This approach also obviates the laborious purification process of recombinant molecules to delineate the specific role of a particular gene of interest *in vivo* within a short period of time and suggests that Fc γ R-Ig molecules can be expressed *in vivo*. Further, these hydrodynamically expressed molecules are useful in evaluating the roles and therapeutic use of Fc γ Rs during the course of various inflammatory and autoimmune diseases.

MATERIALS AND METHODS

Reagents

HRP-conjugated anti-human Fc antibody and ovalbumin (OVA) were purchased from Sigma (St. Louis, MO). The Ovalbumin-FITC, rabbit anti-Ovalbumin IgG and rabbit IgG were from Roche Molecular Biochemicals (Indianapolis, IN). HRP-substrate and SDS-PAGE gels were from BioRad (Hercules, CA). DNA preparation kits were from QIAGEN (Valencia, CA). DH5 α bacteria, CHO serum free media and lipofectamin were from Invitrogen (Carlsbad, CA). The Micro BCA-protein assay kit was from Pierce (Rockford, IL). Quick-change II site-directed mutagenesis kit was from Stratagene (La Jolla, CA). Cell culture reagents were from Life Technologies (Gaithersburg, MD). Site directed mutagenesis kit was from Stratagene (La Jolla, CA). 2.4G2 (anti- mouse mAb CD16/32) mAb was from BD biosciences (San Jose, CA). Anti-human CD16 and CD32A mAb were referenced previously^{40,41}. Human Fc specific Cy5 conjugated F(ab)'2 fragment of goat antibody was from Jackson ImmunoResearch Laboratories (West grove, PA). Mouse macrophage specific FITC conjugated rat monoclonal (clone#BM8) antibody F4/80, rabbit polyclonal mouse liver mitochondrial marker antibody (CSP1) and FITC conjugated, affinity purified goat polyclonal antibody against rabbit IgG were purchased from Abcam (San Francisco, CA). C57BL/6 (8-10 weeks old) female mice were from Jackson laboratory

(Bar Harbor, Maine). The animal experiments were conducted according to the Emory University IACUC protocol.

Construction of human Fc γ R-Ig plasmids

The construction of the dimeric form of hCD32A^R-Ig and hCD16A^F-Ig was carried out by ligating the extracellular domain of hCD32A^R or CD16A^F to the mutated Fc domain of the human IgG1 heavy chain as described earlier^{38,40,41}. The mutations in the Fc domain were shown to abolish the binding of Fc γ Rs⁵¹⁻⁵³. The hCD32A^H-Ig and hCD16A^V-Ig alleles were constructed by Quick-change II site directed mutagenesis kit using hCD32A^R-Ig and hCD16A^F-Ig DNA as a template. cDNA encoding the Fc γ R-Ig were then subcloned into the pcDNA3.1 expression vector with neomycin selection marker^{40,41}. The plasmid DNAs of hFc γ R-Ig were transfected into CHO-K1 cells and culture supernatant was collected and used to purify the Fc γ R-Ig molecules using a Protein-G column as previously described^{40,41}.

Soluble IC binding assay

Soluble immune complex (sIC) was prepared as described^{38,41}. Briefly FITC-Ova was mixed with rabbit anti-Ova IgG (1:1 molar ratio) and incubated for 4 h at 4°C. The complex was centrifuged using a microcentrifuge at 15,000 rpm for 30 min at 4°C and the supernatant was used for the FITC-IC binding assay. The P388D1 mouse macrophage cells (50 μ l of 5×10^6) were preincubated with recombinant receptors or mAbs for 30 min at 4°C and then incubated with FITC-IC (20 μ g/ml) in binding buffer (PBS/EDTA with 1% BSA, pH 7.4) for 1 h at 4°C. The cells were then washed and analyzed by flow cytometry.

Hydrodynamic-based injection of Fc γ R-Ig in mice

Hydrodynamic-based injection of plasmid DNA was carried out as described^{1,56}. Briefly, 10 μ g of pcDNA3.1 plasmid vector containing human Fc γ R-Ig cDNAs (hCD16A^F-Ig, hCD16A^V-Ig, hCD32A^R-Ig or hCD32A^H-Ig) was diluted in 1.6 ml of sterile saline (0.9% NaCl) and injected into each group of mice (n=5 per group) through the tail vein within 5-7 sec, using a 27.5-gauge needle. Subsequently, 5 μ l blood samples were collected at several time points and diluted to 500 μ l in PBS/5 mM EDTA, and the plasma was separated by centrifugation. The plasma samples were frozen immediately at -80°C for future analysis.

Estimation of human Fc γ R-Ig levels in mice blood

Plasma collected at different time points was used to detect Fc γ R-Ig dimers by sandwich ELISA. ELISA plates were coated with 100 μ l of 10 μ g/ml anti-hCD16A (CLBFCgran-1) and anti-hCD32A (IV.3) mAbs overnight at 4°C. The wells were then blocked with PBS/5 mM EDTA /1% BSA. After washing, 100 μ l of the plasma samples were added into the wells and incubated for 1 h. The wells were washed and then incubated for another 1 h after adding 100 μ l of HRP-conjugated goat anti-human IgG Fc specific antibody. HRP substrate was added to the washed wells and read at 450 nm. The plasma from mice injected with PBS served as a specificity control. To quantify the level of Fc γ R-Ig dimers in the blood, purified hCD16A^F-Ig and hCD32A^R-Ig were used as standards, while BSA coated wells served as negative controls.

Reverse passive Arthus reaction (RPA)

RPA was carried out as described³⁸. Briefly, the mice (n=12) were hydrodynamically injected with 10 µg of human FcγR-Ig plasmids in 1.6 ml of saline. Then RPA was carried out at different time points. For each time point, a group of mice (n=3) was used. Before the initiation of RPA, blood was collected (5 µl), diluted to 500 µl with PBS/EDTA, and centrifuged to collect plasma. The diluted plasma was used to analyze the hFcγR-Ig molecules in the circulation. At each time point, a group of mice (n=3) was injected with 12.5 or 25 µg per site of anti-Ova intradermally in a total volume of 25 µl of PBS at different sites. After 5 min following the injection of anti-Ova antibody, RPA was initiated by injecting 100 µl PBS containing 500 µg of ovalbumin and 1% Evan's blue through the tail vein. A group of mice (n=3) injected with PBS served as a control at each time point. Both the control and experimental mice were sacrificed after 3 h and the injection sites on the reverse side of the skin were examined for extravasation of the blue dye. Photographs were taken immediately and used to quantitate the degree of inflammation elicited by RPA reaction. The intensity of each dermal lesion, seen blue in the photographs, was quantified using ImageJ (National Institute of Health, Bethesda) and KaleidaGraph (Synergy Software, Reading, PA).

Western blot analysis

The plasma samples collected at various time points after the hydrodynamic injection of FcγR-Ig plasmid DNA were subjected to Western blot analysis. Briefly, 20 µl of plasma was mixed with SDS-PAGE sample buffer of both reducing and nonreducing conditions. Samples were boiled for 5 min and subjected to SDS-PAGE under reducing or non-reducing conditions. After blotting the protein onto a PVDF membrane, the proteins were detected using a HRP-conjugated goat anti-human Fc specific antibody.

Analyses of ligand binding ability of circulating FcγR-Ig by ELISA

The plasma sample collected at various time points were used to detect the ligand binding activity of hydrodynamically expressed human FcγR-Igs. Briefly, ELISA plates were coated overnight with rabbit IgG (100 µl of 10 µg/ml) and blocked for 1 h with a binding buffer. The plasma samples from mice injected with human FcγR-Ig (100 µl of 1:100 diluted) were then added to the plates, and the incubation continued for 1h at 4°C. The FcγR-Ig bound to the rabbit IgG was detected as described in a previous section. The hFcγR-Igs bound to BSA-coated wells were taken as non-specific binding. Experiments are representative of three individual experiments. Data are mean ± SD of triplicates.

Immunohistochemistry

Expression of FcγRs-Ig molecules in different organs was determined by immunohistochemistry. Briefly, 10 µg of pcDNA3.1 plasmid vector containing human hCD32A^H-Ig was injected hydrodynamically into a group of mice (n=3 per group) as described above. Subsequently, different organs (liver, spleen, kidney, lung and heart) were collected at several time points and snap frozen immediately using liquid nitrogen. 5µm cryo sections were taken and stained for the hCD32A^H-Ig, hepatocytes and macrophages using antibodies. The specific antibodies are: F4/80: a mouse macrophage specific FITC

conjugated rat monoclonal antibody, CSP1: rabbit polyclonal mouse liver mitochondrial marker antibody, and a Cy5 conjugated F(ab)'₂ fragment of goat antibody specific to human Fc domain.

Statistical analysis

A statistical comparison of the control and treated samples was performed using student T-test; $p < 0.01$ (*) considered as significant and $p < 0.001$ (**) considered as highly significant.

ACKNOWLEDGEMENTS

This study was supported by NIH grants R21HL09126802 and R21AR05682101 to P.S and AHA grant 11SDG5710004 to R.S. We also thank Ms. Archana Boopathy, Ms. Sumi Selvaraj and Ms. Danielle Daniels, for critical reading of the article.

REFERENCES

1. Liu F, Song YK, Liu D. Hydrodynamics-based transfection in animals by systemic administration of plasmid DNA. *Gene Ther.* 1999; 6:1258–1266. [PubMed: 10455434]
2. Zhang G, Budker V, Wolff JA. High levels of foreign gene expression in hepatocytes after tail vein injections of naked plasmid DNA. *Hum Gene Ther.* 1999; 10:1735–1737. [PubMed: 10428218]
3. Suda T, Liu D. Hydrodynamic gene delivery: Its principles and applications. *Mol Ther.* 2007; 15:2063–2069. [PubMed: 17912237]
4. Herweijer H, Wolff JA. Gene therapy progress and prospects: Hydrodynamic gene delivery. *Gene Ther.* 2007; 14:99–107. [PubMed: 17167496]
5. Budker VG, Subbotin VM, Budker T, Sebestyen MG, Zhang G, Wolff JA. Mechanism of plasmid delivery by hydrodynamic tail vein injection. II. Morphological studies. *J Gene Med.* 2006; 8:874–888. [PubMed: 16718734]
6. Andrianaivo F, Lecocq M, Wattiaux-De Coninck S, Wattiaux R, Jadot M. Hydrodynamics-based transfection of the liver: entrance into hepatocytes of DNA that causes expression takes place very early after injection. *J Gene Med.* 2004; 6:877–883. [PubMed: 15293346]
7. Kobayashi N, Kuramoto T, Yamaoka K, Hashida M, Takakura Y. Hepatic uptake and gene expression mechanisms following intravenous administration of plasmid DNA by conventional and hydrodynamics-based procedures. *J Pharm exper ther.* 2001; 297:853–860.
8. Chang HE, Hanawa H, Liu H, Yoshida T, Hayashi M, Watanabe R, Abe S, Toba K, Yoshida K, Elnagger R, Minagawa S, Okura Y, Kato K, Kodama M, Maruyama H, Miyazaki J, Aizawa Y. Hydrodynamic-based delivery of an interleukin-22-Ig fusion gene ameliorates experimental autoimmune myocarditis in rats. *J Immunol.* 2006; 177:3635–3643. [PubMed: 16951323]
9. Chang HE, Hanawa H, Yoshida T, Hayashi M, Liu H, Ding L, Otaki K, Hao K, Yoshida K, Kato K, Toba K, Kodama M, Maruyama H, Miyazaki J, Aizawa Y. Alteration of IL-17 related expressions in experimental autoimmune myocarditis and inhibition of IL-17 by IL-10-Ig fusion gene transfer. *Circ J.* 2008; 72:813–819. [PubMed: 18441464]
10. Abe S, Hanawa H, Hayashi M, Yoshida T, Komura S, Watanabe R, Lie H, Chang H, Kato K, Kodama M, Maruyama H, Nakazawa M, Miyazaki J, Aizawa Y. Prevention of experimental autoimmune myocarditis by hydrodynamics-based naked plasmid DNA encoding CTLA4-Ig gene delivery. *J Card Fail.* 2005; 11:557–564. [PubMed: 16198253]
11. Matsui Y, Inobe M, Okamoto H, Chiba S, Shimizu T, Kitabatake A. Blockade of T cell costimulatory signals using adenovirus vectors prevents both the induction and the progression of experimental autoimmune myocarditis. *J Mol Cell Cardiol.* 2002; 34:279–295. [PubMed: 11945021]
12. Kawaguchi Y. A gene therapy or purified CTLA4IgG treatment of experimental allergic encephalomyelitis. *Igaku Zasshi.* 1999; 74:467–475.

13. Mihara M, Tan I, Chuzhin Y, Reddy B, Budhai L, Holzer A, et al. CTLA4Ig inhibits T cell - dependent B-cell maturation in murine systemic lupus erythematosus. *J Clin Invest*. 2000; 106:91–101. [PubMed: 10880052]
14. Quattrocchi E, Dallman MJ, Feldmann M. Adenovirus-mediated gene transfer of CTLA-4Ig fusion protein in the suppression of experimental autoimmune arthritis. *Arthritis Rheum*. 2000; 43:1688–1697. [PubMed: 10943858]
15. Wu X, He Y, Falo L Jr, Hui K, Huang L. Regression of human mammary adenocarcinoma by systemic administration of a recombinant gene encoding the hFlex-TRAIL Fusion Protein. *Mol Ther*. 2001; 3:368–374. [PubMed: 11273779]
16. Yazawa H, Murakami T, Li H-M, Back T, Kurosaka K, Suzuki Y, Shorts L, Akiyama Y, Maruyama K, Parsoneault E, Wilttrout RH, Watanabe M. Hydrodynamics-based gene delivery of naked DNA encoding fetal liver kinase-1 gene effectively suppresses the growth of pre-existing tumors. *Cancer Gene Therapy*. 2006; 13:993–1001. [PubMed: 16763608]
17. Neal ZC, Bates MK, Albertini MR, Herweijer H. Hydrodynamics limb vein delivery of a xenogeneic DNA cancer vaccine effectively induces antitumor immunity. *Mol Ther*. 2007; 15:422–430. [PubMed: 17235322]
18. Hulett MD, Hogarth PM. Molecular basis of Fc receptor function. *Adv Immunol*. 1994; 57:1–127. [PubMed: 7872156]
19. Kimberly RP, Salmon JE, Edberg JC. Receptors for immunoglobulin G. Molecular diversity and implications for disease. *Arthritis Rheum*. 1995; 38:306–314. [PubMed: 7880183]
20. Ravetch JV. Fc receptors. *Curr Opin Immunol*. 1997; 9:121–125. [PubMed: 9039776]
21. Unkeless JC, Scigliano E, Freedman VH. Structure and function of human and murine receptors for IgG. *Annu Rev Immunol*. 1988; 6:251–281. [PubMed: 2968084]
22. Selvaraj P, Fifadara N, Nagarajan S, Cimino A, Wang G. Functional regulation of human neutrophil Fc gamma receptors. *Immunol Res*. 2004; 29:219–230. [PubMed: 15181284]
23. Anderson CL, Looney RJ. Human leukocyte IgG Fc receptors. *Immunol Today*. 1986; 7:264–266. [PubMed: 25290629]
24. Fanger MW, Shen L, Graziano RF, Guyre PM. Cytotoxicity mediated by human Fc receptors for IgG. *Immunol Today*. 1989; 10:92–99. [PubMed: 2525910]
25. Hogarth PM. Fc receptors are major mediators of antibody-based inflammation in autoimmunity. *Curr Opin Immunol*. 2002; 14:798–802. [PubMed: 12413532]
26. Takai T. Fc receptors and their role in immune regulation and autoimmunity. *J Clin Immunol*. 2005; 25:1–18. [PubMed: 15742153]
27. Schmidt RE, Gessner JE. Fc receptors and their interaction with complement in autoimmunity. *Immunol Lett*. 2005; 100:56–67. [PubMed: 16125792]
28. Takai T, Li M, Sylvestre D, Clynes R, Ravetch JV. FcR gamma chain deletion results in pleiotropic effector cell defects. *Cell*. 1994; 76:519–529. [PubMed: 8313472]
29. Ravetch JV, Clynes RA. Divergent roles for Fc receptors and complement *in vivo*. *Annu Rev Immunol*. 1998; 16:421–432. [PubMed: 9597136]
30. Clynes R, Dumitru C, Ravetch JV. Uncoupling of immune complex formation and kidney damage in autoimmune glomerulonephritis. *Science*. 1998; 279:1052–1054. [PubMed: 9461440]
31. Hazenbos WL, Gessner JE, Hofhuis FM, Kuipers H, Meyer D, Heijnen IA, Schmidt RE, Sandor M, Capel PJ, Daeron M, Van de Winkel JGJ, Verbeek JS. Impaired IgG-dependent anaphylaxis and Arthus reaction in FcγRIII (CD16) deficient mice. *Immunity*. 1996; 5:181–188. [PubMed: 8769481]
32. Clarkson SB, Bussel JB, Kimberly RP, Valinsky JE, Nachman RL, Unkeless JC. Treatment of refractory immune thrombocytopenic purpura with anti-Fcγ receptor antibody. *New Engl J Med*. 1986; 314:1236–1239. [PubMed: 2939345]
33. Ravetch JV, Bolland S. IgG Fc Receptors. *Annu Rev Immunol*. 2001; 19:275–290. [PubMed: 11244038]
34. Takai T, Nakamura A, Akiyama K. Fc receptors as potential targets for the treatment of allergy, autoimmune disease and cancer. *Curr Drug Targets Immune Endocr metabol Disord*. 2003; 3:187–197. [PubMed: 12871025]

35. Cohen-Solal JF, Cassard L, Fridman WH, Sautes-Fridman C. Fc gamma receptors. *Immunol Lett.* 2004; 92:199–205. [PubMed: 15081612]
36. Tarzi RM, CooK HT. Role of Fc gamma receptors in glomerulonephritis. *Nephron Exp Nephrol.* 2003; 95:e7–12. [PubMed: 14520009]
37. Nakamura A, Takai T. A role of FcγRIIB in the development of collagen-induced arthritis. *Biomed Pharmacother.* 2004; 58:292–298. [PubMed: 15194165]
38. Shashidharamurthy R, Hennigar RA, Fuchs S, Palaniswami P, Sherman M, Selvaraj P. Extravasations and emigration of neutrophils to the inflammatory site depend on the interaction of immune-complex with Fcγ receptors and can be effectively blocked by decoy Fcγ receptors. *Blood.* 2008; 111:894–904. [PubMed: 17962513]
39. Arthus M. Injections repetees de serum de cheval chez le lapin. *C R Soc Biol.* 1903; 55:817.
40. Li P, Nagarajan S, Zhu C, Selvaraj P. Recombinant CD16A-Ig forms a homodimer and cross-blocks the ligand binding functions of neutrophil and monocyte Fcγ receptors. *Mol Immunol.* 2002; 38:527–538. [PubMed: 11750654]
41. Shashidharamurthy R, Zhang F, Amano A, Kamat A, Panchanathan R, Ezekwudo D, Zhu C, Selvaraj P. Dynamics of the interaction of human IgG subtypes immune complexes with cells expressing R and H allelic forms of a low affinity Fcγ receptor CD32A. *J Immunol.* 2009; 183:8216–8224. [PubMed: 20007585]
42. Ignatovich AI, Dizhes EB, Palvotskayan AV, Akifiev BN, Burov SV, Orlov SV, Perevozchikov AP. Complexes of plasmid DNA with basic domain 47-57 of the HIV-1 tat protein are transferred to mammalian cells by endocytosis-mediated pathways. *J Biol Chem.* 2003; 278:42625–42636. [PubMed: 12882958]
43. van Sorge NM, van der Pol W-L, Van de Winkel JGJ. FcγR Polymorphisms: Implications for function, disease susceptibility and immunotherapy. *Tis Ant.* 2003; 61:189–202.
44. Dijkstra HM, Bijl M, Fijnheer R, Scheepers RH, Oost WW, Jansen MD, Sluiter WJ, Limburg PC, Derksen RH, van de Winkel JG. Fcγ receptor polymorphisms in systemic lupus erythematosus: association with disease and in vivo clearance of immune complexes. *Arthritis and Rheum.* 2000; 43:2793–2800. [PubMed: 11145038]
45. Ludo van der Pol W, van de Winkel GJ. IgG receptor polymorphism: risk factor for disease. *Immunogen.* 1998; 48:222–232.
46. Tan SY. FcγRIIIa polymorphism in systemic lupus erythematosus. *Kidney Blood Press Res.* 2000; 23:138–142. [PubMed: 10765117]
47. Manger K, Repp R, Spriewald BM, Rascu A, Geiger A, Wassmuth R, Westerdal NA, Wentz B, Manger B, Kalden JR, van de Winkel JG. Fcγ receptor IIa polymorphism in caucasian patients with systemic lupus erythematosus: association with clinical symptoms. *Arthritis Rheum.* 1998; 41:1181–1189. [PubMed: 9663473]
48. Ierino FL, Powell MS, McKenzie IFC, Hogarth PM. Recombinant soluble human Fc-gamma-RII: Production, characterization, and inhibition of the Arthus reaction. *J Exp Med.* 1993; 178:1617–1628. [PubMed: 8228810]
49. Ellsworth JL, Hamacher N, Harder B, Bannink K, Bukowski TR, Byrnes-Blake K, Underwood S, Oliver C, Waggle KS, Noriega C, Hebb L, Rixon MW, Lewis KE. Recombinant soluble human FcγRIA (CD64A) reduces inflammation in murine collagen induced arthritis. *J Immunol.* 2009; 182:7272–7279. [PubMed: 19454724]
50. Junghans RP, Anderson CL. The protection receptor for IgG catabolism is the β2-microglobulin-containing neonatal intestinal transport receptor. *Proc Natl Acad Sci USA.* 1996; 93:5512–5516. [PubMed: 8643606]
51. Chappel MS, Isenman DE, Everett M, Xu YY, Dorrington KJ, Klein MH. Identification of the Fcγ receptor class I binding site in human IgG through the use of recombinant IgG1/IgG2 hybrid and point-mutated antibodies. *Proc Natl Acad Sci USA.* 1991; 88:9036–9040. [PubMed: 1833770]
52. Chappel MS, Isenman DE, Oomen R, Xu YY, Klein MH. Identification of a secondary FcγRI binding site within a genetically engineered human IgG antibody. *J Biol Chem.* 1993; 268:25124–25131. [PubMed: 8227075]

53. Ledbetter JA, Gilliland LK, Hayden MS, Linsley PS, Bajorath J, Fell HP. Expression vectors encoding bispecific fusion proteins and methods of producing biologically active bispecific fusion proteins in mammalian cells. United States Patent. 2000:6132992.
54. Tao MH, Morrison SL. Studies of aglycosylated chimeric mouse-human IgG. Role of carbohydrate in the structure and effector functions mediated by the human IgG constant region. *J Immunol.* 1989; 143:2595–2601. [PubMed: 2507634]
55. Roopenian DC, Akilesh S. FcRn: the neonatal Fc receptor comes of age. *Nat Rev Immunol.* 2007; 7:715–725. [PubMed: 17703228]
56. He Y, Pimenov AA, Nayak JV, Plowey J, Falo LD, Huang L. Intravenous injection of naked DNA encoding secreted flt3 ligand dramatically increases the number of dendritic cells and natural killer cells in vivo. *Hum Gene Ther.* 2000; 11:547. [PubMed: 10724033]




Figure 1. Purified human Fc γ R-Igs compete with cell surface mouse Fc γ Rs and block IC binding both *in vitro* and *in vivo*

(A) Human Fc γ R-Igs blocked the binding of sICs to the mouse macrophage cell line P388D1 in a dose dependent manner. P388D1 cells were incubated with FITC-IC (sIC) in the presence and absence of various concentrations of hFc γ R-Ig dimers, and the cells were analyzed for binding of FITC-IC using flow cytometry. (B) Systemic administration of purified hFc γ R-Ig dimers efficiently blocked the RPA reaction *in vivo* in mice. Groups of mice (n=3/group) were injected with hFc γ R-Ig intravenously (100 μ g/mouse). After 1 h, the mice were injected intradermally with PBS (site 1) or anti-Ova (12.5 μ g in site 2 and 25 μ g in site 3). RPA was initiated by injecting Ova with 1% Evan's blue intravenously through the tail vein. The anti-Fc γ R antibody (2.4G2) treated control mice were injected with 25 μ g/ml of blood (40 μ g/mice) mAb. The PBS injected mice served as the untreated positive control. After 3 h the mice were euthanized, and the dorsal side of the skin was photographed for analysis. The figure shows three representative mice. (C) Quantitative analysis of RPA. The dermal lesion, seen blue in the photographs, was quantified using ImageJ and KaleidaGraph softwares for groups with or without Fc γ R-Ig dimer treatment. Data are presented as the mean \pm SD from three experiments. *p<0.01, **p<0.001.




Figure 2. Hydrodynamic based expression of human FcγR-Ig molecules in mice

Human FcγR-Ig genes in plasmid DNA (10 μg) were diluted in 1.6 ml of sterile saline and injected within 5 seconds into each group of mice through the tail vein. Subsequently, 5 μl blood samples were collected at different time points and diluted in PBS/EDTA. Plasma was separated to detect FcγR-Ig dimers using ELISA as described under Materials and Methods. The plasma from a group of normal mice injected with PBS served as a specificity control. Purified hCD16A-Ig and hCD32A-Ig were used as positive controls, while BSA coated wells served as negative controls. Purified hCD32A-Ig was used as a standard to quantify the level of dimers in the blood. Experiments are representative of three individual experiments. Data are mean ± SD of triplicates.




Figure 3. Hydrodynamically expressed human Fc γ R-Ig molecules block antibody mediated inflammation in mice

Mice (n=12) were injected with 10 μ g of plasmid vector containing human Fc γ R-Ig cDNA (Panel A: hCD16A^F-Ig, Panel B: hCD32A^R-Ig, Panel C: hCD32A^H-Ig) intravenously as described under Material and Methods. RPA was carried out using three mice at each time point. Mice were injected intradermally with PBS (site 1) and 25 μ g (site 2) of anti-Ova per site. RPA was initiated by injecting Ova with 1% Evan's blue intravenously through the tail vein. The antibody (2.4G2) treated control mice were injected with 25 μ g/ml of blood (40 μ g/mouse) mAb. The PBS injected mice served as the untreated positive control. After 3 h the mice were euthanized, and the dorsal side of the skin was photographed for analysis. The figure is representative of three individual mice. Bar graph on the right side represents the quantitative analysis of RPA. The dermal lesion, seen blue in the photographs, was quantified using ImageJ and KaleidaGraph softwares for groups with or without Fc γ R dimer treatment. Data are presented as the mean \pm SD from three experiments. *p<0.01, **p<0.001.




Figure 4. Expression of human FcγR-Ig molecules in mice used to conduct RPA

Before the initiation of RPA (presented in Fig. 3), blood was collected (5 μl) and diluted in 500 μl of PBS/EDTA and plasma was used to detect the expression of human FcγR-Ig molecules by sandwich ELISA as described under the figure 3 legend. Experiments are an average of three individual mice.




Figure 5. Binding of hydrodynamically expressed human Fc γ R-Igs to rabbit IgG

Plasma collected from the mice used to conduct the RPA in figure 3 was used to determine whether the *in vivo* expressed hFc γ R-Igs bind to rabbit IgG using a sandwich ELISA. Rabbit IgG (100 μ l of 10 μ g/ml) was coated on ELISA plates overnight, and the binding of hFc γ R-Ig molecules to rabbit IgG was determined as described under Materials and Methods. The hFc γ R-Ig alleles that bound to BSA-coated wells were taken as non-specific binding. Experiments are an average of three individual mice. Data are mean \pm SD of triplicates.




Figure 6. Western blot analysis of hydrodynamically expressed human Fc γ R-Igs in mice
 To determine the dimeric nature of hydrodynamically expressed hFc γ R-Igs, the plasma samples collected from the mice, used to conduct the RPA in figure 3, were subjected to Western blot analysis. 20 μ l of plasma was mixed with SDS-PAGE sample buffer under both reducing and nonreducing conditions. After electroblotting the protein onto the PVDF membrane, the proteins were detected using the HRP-conjugated goat anti-human Fc specific antibody. (a) hCD16A^F-Ig, (b) hCD32A^R-Ig, (c) hCD32A^H-Ig.




Figure 7. Expression of hFc γ R-Ig in liver and spleen cells after hydrodynamic administration of plasmid DNAs

To determine the specific organ and cell type involved in the expression of hFc γ R-Ig molecules, a group of mice (n=3) were hydrodynamically injected with hCD32A^H-Ig plasmid DNA (10 μ g) and then various organs at different time points were collected and immunohistochemistry was carried out as described under Materials and Methods. **A:** Liver sections were stained with a liver mitochondrial marker specific rabbit antibody to detect hepatocytes followed by FITC conjugated goat anti-rabbit as secondary antibody (green). Liver sections are also counterstained with Cy-5 conjugated goat anti-human Fc specific antibody to detect the presence of hCD32A^H-Ig expression (red). The merger of green and red color (yellow) represents the expression of hCD32A^H-Ig by hepatocytes. Photographs were taken at 40X magnification. **B:** Spleen sections were stained with FITC conjugated macrophage specific rat antibody to detect macrophages in the spleen (green) and counter stained with Cy-5 conjugated goat anti-human Fc specific antibody to detect the presence of hCD32A^H-Ig expression (red). The yellow color, which is a result of merger of green and red color indicates, the expression of hCD32A^H-Ig by macrophages. Photographs were taken at 20X magnification. WRP represents white pulp region where T and B-cells are present. RPR represents red pulp region around WRP where the hCD32A^H-Ig molecules were expressed by macrophages.