ERRATUM Open Access



# Erratum to: Inhibition of myeloperoxidase oxidant production by N-acetyl lysyltyrosylcysteine amide reduces brain damage in a murine model of stroke

Guoliang Yu<sup>1</sup>, Ye Liang<sup>1</sup>, Ziming Huang<sup>1,2</sup>, Deron W. Jones<sup>1</sup>, Kirkwood A. Pritchard Jr<sup>1</sup> and Hao Zhang<sup>1\*</sup>

## **Erratum**

After publication of this work [1], it was noted that there was an error within Fig. 1. Within the graph of Fig. 1a, the labels for PBS and KYC were inadvertently reversed. The open circles should represent KYC, whereas the filled circles should represent PBS. Figure 1 has been corrected in the original article and is also included correctly below.

### **Author details**

<sup>1</sup>Division of Pediatric Surgery, Department of Surgery, Medical College of Wisconsin, 8701 Watertown Plank Rd, Milwaukee 53226, Wl, USA. <sup>2</sup>Department of Breast Surgery, Maternal and Child Health Hospital of Hubei Province, 745 WuLuo Road, Hongshan District, Province 430070, Wuhan City, Hubei. China.

Received: 21 June 2016 Accepted: 21 June 2016 Published online: 27 June 2016

# References

 Yu G, Liang Y, Huang Z, Jones DW, Pritchard KA, Zhang H. Inhibition of myeloperoxidase oxidant production by N-acetyl lysyltyrosylcysteine amide reduces brain damage in a murine model of stroke. J Neuroinflammation. 2016;13:119. doi:10.1186/s12974-016-0583-x.

<sup>1</sup>Division of Pediatric Surgery, Department of Surgery, Medical College of

Wisconsin, 8701 Watertown Plank Rd, Milwaukee 53226, Wl, USA

# Submit your next manuscript to BioMed Central and we will help you at every step:


- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at www.biomedcentral.com/submit





\* Correspondence: hzhang@mcw.edu



**Fig. 1** KYC significantly reduced neurological deficit and brain infarct size in mice subjected to MCAO. **a** Effects of KYC on neurological severity scores during 7 days after MCAO. *Top panel*: Experimental scheme; KYC (10 mg/kg/day) or PBS via i.p. was administrated starting 1 h after reperfusion. *Bottom panel*: Neurological severity scores of mice after MCAO (\*p < 0.05, KYC vs. PBS group at same time interval, nonparametric Mann–Whitney test); **b** Effect of KYC on infarct size in mice 3 days after MCAO. Images of brain coronal sections (20- $\mu$ m thick, 1-mm interval from rostral to caudal) by cresyl violet stain were shown; c Infarct volume (\*p < 0.05, KYC vs. PBS group, n = 5/group, t test)