


Development and Validation of a Patient-Reported Outcome Measure for Fingernail and Toenail Conditions: The NAIL-Q

Anne F Klassen ¹, Charlene Rae ¹, Maureen O'Malley², Trisia Breitkopf¹, Leah Algu¹, Jasmine Mansouri³, Claire R Brown⁴, Yi Wang⁵, Shari R Lipner ⁴

¹Department of Pediatrics, McMaster University, Hamilton, Ontario, Canada; ²Division of Dermatology, Department of Medicine, McMaster University, Hamilton, Ontario, Canada; ³Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada; ⁴Department of Dermatology, Weill Cornell Medicine, New York City, NY, USA; ⁵Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada

Correspondence: Anne F Klassen, McMaster University, 3N27, 1200 Main Street W, Hamilton, ON, L8N 3Z5, Canada, Email aklass@mcmaster.ca

Background: Patient-reported outcome measures (PROMs) are needed to measure outcomes that matter to people with nail conditions, from their perspective.

Objective: To design a comprehensive new PROM (NAIL-Q) to measure outcomes important in toenail and fingernail conditions.

Methods: A mixed methods iterative approach was used. Phase 1 involved concept elicitation interviews that were audio-recorded, transcribed, and coded line-by-line. Concepts were developed into scales and refined through cognitive debriefing interviews with patients and expert input. Data was then collected from an international sample using a crowdsourcing platform. Eligible participants were aged ≥ 18 years with a nail condition for at least 3 months. Rasch Measurement Theory (RMT) analysis was used to examine item and scale performance. Other psychometric tests included test-retest reliability, and convergent and construct validity.

Results: Phase 1 interviews involved 23 patients with 10 nail conditions and input from 11 dermatologists. The analysis led to the development of 84 items for field-testing. In Phase 2, 555 participants completed the survey. Toenail conditions ($n = 441$) were more common than fingernail conditions ($n = 186$). The RMT analysis reduced the number of items tested to 45 in 7 scales measuring nail appearance, health-related quality of life concerns, and treatment outcomes. All items had ordered thresholds and nonsignificant chi-square p values. Reliability statistics with and without extremes for the Person Separation Index were ≥ 0.79 and Cronbach's alpha were ≥ 0.83 , and for intraclass correlation coefficients were ≥ 0.81 . Construct validity was further supported in that most participants agreed that the NAIL-Q was easy to understand, asked relevant and important questions in a respectful way, and that it should be used to inform clinical care.

Conclusion: The NAIL-Q is a rigorously designed and tested PROM that measures nail appearance, health-related quality of life and treatment outcomes. This PROM can be used in clinical practice to inform patient care and to include the patient perspective in research.

Keywords: validity, reliability, psychometrics, health-related quality of life

Introduction

Nail conditions are common in the population, affect people of all ages, and have a broad differential diagnosis.¹⁻⁷ Nail conditions can cause distress due to their cosmetic appearance, symptoms such as pain, reduced function and mobility, reducing one's health-related quality of life (HRQL).⁸⁻¹¹ Patient reported outcome measures (PROMs) are valuable tools that can be used to measure these concepts from the patient perspective.¹² PROMs are increasingly used in research studies to inform clinical care and improve health care quality.¹³⁻¹⁸

There are currently seven published PROMs for nail conditions.¹⁹⁻²⁵ Five PROMs were developed for onychomycosis, including the Onychomycosis-Specific Questionnaire,¹⁹ Health-Related Quality of Life Measure for Onychomycosis,²⁰ Onychomycosis Disease-Specific Questionnaire (ODSQ),²¹ NAILQoL,²² and the OnyCOE-t.²³ Two

PROMs were developed for nail psoriasis, including the Psoriasis Quality of Life Scale (NPQ10)²⁴ and the Nail Assessment in Psoriasis and Psoriatic Arthritis QOL (NAPPA-QOL).²⁵ A review that applied Consensus-based Standards for the selection of health Measurement Instruments (COSMIN) guidelines^{26,27} concluded that these seven PROMs had limited evidence supporting validity, reliability, and responsiveness.²⁸

A new PROM is needed to provide evidence about the impact and effectiveness of treatments for all types of nail conditions from the patient perspective. The aim of our study was to develop the NAIL-Q for use with patients with any type of fingernail or toenail condition using the Rasch Measurement Theory (RMT) approach.^{29,30}

Patients and Methods

Ethics Statement

Ethics board approval was obtained from the Hamilton Integrated Research Ethics Board approvals #7370 and #14622 and from Weill Cornell Medicine (New York, NY, USA).

Approach

We used a mixed methods approach³¹ and adhered to best practices guidelines for PROM development.¹²

Phase 1: Qualitative

In phase 1, Interpretative Description³² was used to generate knowledge relevant to a clinical context. Participants were recruited from a dermatology clinic in Canada and one in the USA and provided with a thank-you gift card for each task they helped with (ie, initial, cognitive, and final review), as described below. Research staff were asked to recruit a purposive sample of English-speaking adults who varied by age, gender, ethnicity, and nail conditions. Informed consent was obtained, and interviews were conducted by phone by a skilled qualitative researcher. A semi-structured interview guide was used and updated as new concepts were elicited ([Table S1](#)).

Interviews were audio-recorded, transcribed verbatim et literatim, and coded using a line-by-line approach. Concepts were labelled with a domain and major/minor theme using constant comparison to refine categories and form a conceptual framework.³³ Interviews continued until redundancy of concepts was reached.³⁴ Codes were used to develop items to form NAIL-Q scales. Scales were assigned instructions, a time frame for answering, and response options.

A series of cognitive debriefing interviews with patients were performed by phone/Zoom by two skilled qualitative interviewers. The think aloud method^{35,36} was used to determine if participants found the instructions, response options, and scale content easy to understand, relevant, and comprehensive.

Interviews were audio-recorded, transcribed verbatim et literatim, and coded. The scales were revised based on patient feedback and included in a REDCap survey³⁷ to which dermatologists with nail expertise were invited by email to review. To establish content validity, experts were asked to identify any content that they deemed would be hard to understand, not relevant to nail conditions, or missing.

To ensure changes suggested by experts were acceptable to patients, a REDCap survey was sent to the cognitive debriefing sample via email. Participants were asked to review the scale and identify any items that were hard to understand, not relevant or missing. Nonrespondents were emailed after one week, followed by a phone call one week later. Feedback was used to finalize the NAIL-Q field-test version.

Phase 2: Quantitative

Data for this phase were collected from Prolific Academic (www.prolific.co), an online crowd working platform. Participants were paid the equivalent of 10.50 sterling per hour. A short pre-screen survey was directed to Prolific participants fluent in English. Instructions asked people to complete the pre-screen if they had a fingernail and/or toenail condition for 3 months or longer for which they had seen a health professional in the past. Six images depicting common nail conditions were included, and people were asked to identify their nail condition from the images, or to indicate “other” or “not sure” ([Figure S1](#)).

Participants with any of the six nail conditions were invited to complete the full survey. This survey included demographics (age, gender, race, marital status, education, daily activity, and country) and clinical questions, including severity, sides and number of nails affected, and how long the condition had lasted. Single items asked about pain,

embarrassment, and activity interference. Participants were asked when they last saw a healthcare professional for their nail condition, and the number of appointments in the past 2 years. For seven treatments (topical medications, oral medications, biologics, laser, surgery, taping, injections, and clipping nails) participants were asked to indicate their status (ie, do not need, need, currently having, have had, had and need more). Those who had treatment were asked when they had treatment, and if the treatment made their condition better, the same, or worse.

For the NAIL-Q, participants whose nail condition affected fingernails and toenails completed the Appearance, Distress and Symptoms scales twice. The Outcome scale was completed by participants who had or were having treatment. The NAIL-Q was followed by six evaluation questions with disagree/agree response options: (1) asked questions in a respectful way, (2) was easy to understand, (3) was thorough, (4) asked questions that are important to me, (5) should be used by doctors treating people with nail conditions, and (6) would help my doctor better understand my nail condition. Finally, participants completed the generic EQ-5D-5L health-related quality of life utility PROM, which was scored using the USA normative values.³⁸ EQ-5D-5L includes 5 items (mobility, self-care, usual activities, pain/discomfort, anxiety/depression) with five response options that range from one (no problem) to five (unable to/extreme problem).

At the end of the survey, participants were asked to participate in a test–retest (TRT) study. Late respondents (less than seven days from the initial survey) and those with missing scale data were not invited. We asked if there had been a change in their nail condition and if treatment had been initiated since the initial survey. We aimed to include 100 participants per scale.

The RMT analysis used RUMM2030 software and the unrestricted Rasch model for polytomous data (RUMM version 2030, RUMM Laboratory Pty Ltd, Duncraig, Western Australia, Australia, 1998–2021). Our goal was to identify the best subset of items to retain in each scale. The RMT analysis is described in detail elsewhere.²⁹ Briefly, we performed a range of statistical and graphical tests to examine if the data fit the Rasch model. We adjusted the sample size to 500 for tests that used a X^2 analysis. We examined: (1) item response categories to determine if item responses were properly ordered; (2) item fit statistics to determine if scale items mapped out an item difficulty continuum; (3) local dependency to determine if item residuals were too closely related to each other and performed subtests if residual correlations were ≥ 0.20 ; (4) targeting to determine how well a scale measured the experience of the sample; (5) Person Separation Index and Cronbach's alpha values³⁹ to determine scale reliability using criteria that values should be ≥ 0.70 ;^{26,40} and (6) Differential Item Functioning (DIF) to determine if items in a scale behaved the same for age (<30 vs ≥ 30 years), gender (man vs woman), country (UK vs USA), education (completed college/trade school or university yes/no), and location (toenail vs fingernail). DIF analysis was performed with random samples to ensure subgroups were equivalent in size and repeated three times. We only performed DIF if there were at least 150 participants in the analysis. Any items with DIF were split on the characteristic, and person locations from the original and split analyses were correlated to inspect the impact of DIF on scale scoring.

In SPSS Version 28, we performed Principal Component Analysis and hypothesized that scale items would load ≥ 0.70 providing evidence that the items were part of the latent variable.⁴¹ Scale scores were transformed from 0 (worst) to 100 (best) based on person locations. For the TRT data, anyone who reported their nail condition had changed, or they had initiated treatment, were excluded. Intra-class correlation coefficients (ICC) were computed with a two-way random effects model; values should be ≥ 0.70 .^{26,40}

For convergent validity, NAIL-Q scales were correlated with the EQ-5D-5L scores. COSMIN criteria stipulate that the correlations should be ≥ 0.50 between scales measuring similar constructs, 0.30 – 0.50 between scales measuring related but dissimilar constructs, and <0.30 between scales measuring unrelated constructs.²⁶ We hypothesized that with EQ-5D-5L, the NAIL-Q's Appearance and Outcome scales represent dissimilar constructs, and the NAIL-Q's HRQL scales represent related but dissimilar constructions. In terms of construct validity, we tested six hypotheses of known group differences (Table 1).

Parametric and nonparametric tests were used based on whether the scale data were normally distributed assessed by Skewness and Kurtosis values.⁴² Categorical variables with small cell sizes were combined for analyses. P-values of <0.05 were considered statistically significant.

Table I Construct Validation Hypotheses and Results

	Nail Appearance	Nail Distress	Nail Symptoms	Strength: Fingernail	Physical: Fingernails	Physical: Toenails	Treatment Outcome
1. As condition severity increases NAIL-Q scores	Decrease**	Decrease**	Decrease**	Decrease*	Decrease**	Decrease**	NA
2. As level of pain increases NAIL-Q scores ...	NA	NA	Decrease**	NA	NA	NA	NA
3. As person reports more embarrassment NAIL-Q scores ...	Decrease**	Decrease**	Decrease**	NA	NA	NA	NA
4. As person reports more interference with activities NAIL-Q scores	NA	Decrease**	Decrease**	Decrease**	Decrease**	Decrease**	NA
5. Both side involvement will have NAIL-Q scores that are...	Lower**	Lower*	NA	NA	NA	NA	NA
6. With improvement with treatment NAIL-Q scores ...	NA	NA	NA	NA	NA	NA	Increase**

Note: *p < 0.05; **p < 0.001. NA= Not applicable

Results

Phase I: Qualitative Study

Interviews with 19 women and four men were conducted between October 2019 and November 2020. [Table S2](#) shows sample characteristics.

Participants described a range of outcomes that mattered to them. Participants were concerned about having weak nails that chip, break, crack, split, peel, loosen, and fall off [“Well they are also splitting and cracking and peeling off”]. Pain was a common symptom [“I will get an ingrown nail on my finger, and it swells up. It gets painful, extremely painful”]. Some participants described trouble using their hands [“I need to remember that, oh, I need to grip that differently or hold that differently”], and feet [“...I started to feel it every time I was walking”]. Nail appearance was typically described negatively [“unattractive”, “ugly”, “not normal”, “bad”]. Specific problems included pits, indents, or nails that were uneven, thin, or discolored (eg, spots or stripes). Participants generally disliked their nails [“It was terribly embarrassing”], and participants felt self-conscious of their nails [“... when I used to take the train, holding onto a pole, I’m conscious then of it...you know, being messed up”]. Many participants described concealment behaviors [“I keep my polish on my nails all the time. If I don’t, I put Band-Aids over them”].

The first draft of NAIL-Q included six scales with 89 items. Changes made in each round are shown in [Table S3](#) and summarized in [Table S4](#). Revisions took place between June 2021 and January 2022. Round 1 included seven cognitive debriefing interviews. Based on this round, 16 items were revised, 20 items dropped, and two items added. NAIL-Q had 71 items after Round 1. In Round 2, 35 clinical experts were invited to review the NAIL-Q and 11 (five men, six women) complied. Experts were from the USA (n = 5), Canada (n = 4), Brazil (n = 1) and Switzerland (n = 1). All experts were dermatologists; five had intermediate and six had advanced expertise in nail conditions. Based on this round, two items were revised, one item dropped, and 13 items added. At the end of Round 2, NAIL-Q had 83 items. In Round 3, the seven cognitive interview participants were invited to review the NAIL-Q and five participated. Based on this round, one item was revised, and one item added. The final field-test version included 84 items.

Phase 2: Quantitative Study

The field-test study took place in June and July 2022. The recruitment process is shown in [Figure 1](#). The screen, full survey, and TRT took participants 3 (SD = 2.4), 12.8 (SD = 6.4) and 8.5 (SD = 5.7) minutes on average to complete, respectively.




Figure 1 Recruitment flow diagram.

Table 2 and 3 show the sample demographic and clinical characteristics, respectively. The 555 participants came from 23 countries and ranged in age from 18 to 80 years (Mean 36.7; SD = 13.9). Toenail conditions were more common (n = 441) than fingernail conditions (n = 186). A small sample (72, 13%) had both fingernail and toenail conditions. Most participants (n = 325, 58.6%) had seen a healthcare professional about their nails 1 or 2 times in the past 2 years.

The RMT analysis reduced the NAIL-Q to 45 items. Figure 2 shows the final conceptual framework. Each aspect of the framework is measured by a NAIL-Q scale. The scales include five or six response options that measure satisfaction (Nail Appearance), frequency (Nail Distress, Physical: Fingernails, Physical: Toenails), concern (Nail Symptoms; Strength: Fingernails), and agreement (Treatment Outcome). Item-level fit statistics and DIF results are shown in Table S5. All 45 items had ordered thresholds and non-significant $X^2 P$ values after Bonferroni adjustment.

Table 2 Sample Demographic Characteristics for 555 Participants

Characteristics		n	%
Country	USA	186	33.5
	UK	170	30.6
	South Africa	53	9.5
	Portugal	36	6.5
	Poland	25	4.5
	Mexico	21	3.8
	Italy	18	3.2
	Spain	14	2.5
	Other	32	5.8
Gender	Woman	299	53.9
	Man	243	43.8
	Other gender	13	2.3
Age	18–29	240	43.3
	30–39	130	23.4
	40–49	77	13.9
	50–59	67	12.1
	60+	41	7.4
Race	White	383	69.0
	Black	69	12.4
	Latin American	35	6.3
	South Asian	18	3.2
	East Asian	8	1.4
	Middle Eastern	7	1.3
	Southeast Asian	5	0.9
	Other	30	5.4

(Continued)

Table 2 (Continued).

Characteristics		n	%
Marital status	Never married	291	52.4
	Separated	13	2.3
	Divorced	24	4.3
	Widowed	4	0.7
	Living common-law	55	9.9
	Married	149	26.8
	Other	11	2.0
	Prefer not to answer	8	1.4
Highest education	Some high school	5	0.9
	Completed high school	71	12.8
	Some college/trade school/university	120	21.6
	Completed college/trade school/university degree	240	43.2
	Some Masters or Doctoral degree	41	7.4
	Completed Masters or Doctoral degree	73	13.2
	Prefer not to answer	5	0.9
Daily activity	Working full time	289	52.1
	Working part-time	92	16.6
	Not working/not looking for work	16	2.9
	Unemployed and looking for work	36	6.5
	Retired	23	4.1
	Disabled/unable to work	10	1.8
	Currently in school	70	12.6
	Other	13	2.3
	Prefer not to answer	6	1.1

Item fit was within +2.5 for 37 items. DIF was identified for 14 items including 2 of 35 tested for country, 4 of 45 tested for age, 4 of 35 tested for gender, and 9 of 23 tested for location. Pearson correlations between person locations for items before and after item split for DIF indicated marginal impact of DIF on scoring, with all correlations ≥ 0.996 .

Scale-level results are shown in Table 4. The data fit the Rasch model for six scales with a slight misfit for Strength: Fingernails ($p = 0.02$). PSI values were ≥ 0.79 , and Cronbach's alpha values were ≥ 0.83 with and without extremes. Three pairs of items in the Appearance scale and one pair in the Outcome scale evidenced local dependency. Subtests to examine their impact on reliability showed a drop in PSI values of ≤ 0.01 . The proportion of patients to score on the scale ranged from 76.5% to 96%. While two scales evidenced some ceiling effect, it should be noted that 80% (Physical: Fingernails) and 83% (Physical: Toenails), of those to score at the ceiling chose "Not at all" when asked: "How much

Table 3 Sample Clinical Characteristics

Characteristics		Fingernails		Toenails	
		n	%	n	%
Condition*	Onychomycosis	18	–	150	–
	Psoriasis	22	–	15	–
	Ingrown	48	–	232	–
	Brittle	98	–	42	–
	Onycholysis	31	–	53	–
	Paronychia	18	–	20	–
	Other	3	–	1	–
Self-reported severity	Mild	72	38.7	156	35.7
	Moderate	101	54.3	238	54.5
	Severe	12	6.5	36	8.2
	Very severe	1	0.5	7	1.6
Sides affected	One side	71	38.2	201	25.9
	Both sides	115	61.8	236	54.0
Number of nails affected	1	43	23.1	135	24.4
	2	28	15.1	156	28.1
	3	16	8.6	44	7.9
	4	16	8.6	35	6.3
	5	11	5.9	17	3.1
	6	6	3.2	12	2.2
	7	9	4.8	4	0.7
	8	9	4.8	5	0.9
	9	1	0.5	4	0.7
	10	46	24.7	25	4.5
Time had condition	≤6 months	56	30.1	144	33.3
	6–12 months	29	15.6	64	14.6
	1–5 years	60	32.3	120	27.5
	>5 years	41	22.0	109	24.9
Treatment status	Do not require	27	14.8	20	4.6
	Need	29	15.9	59	13.6
	Currently having	50	27.5	129	29.7
	Had and need more	32	17.6	127	29.3
	Completed	44	24.2	99	22.8

(Continued)

Table 3 (Continued).

Characteristics		Fingernails		Toenails	
		n	%	n	%
Compared with before treatment	Worse	5	3.0	18	5.0
	About the same	37	29.4	96	26.4
	A little better	59	46.8	163	44.9
	A lot better	25	19.8	86	23.7

Notes: *43 (23.1%) reported more than 1 fingernail condition and 61 (14%) reported more than 1 toenail condition.

does your nail condition interfere with doing your usual daily activities?” Targeting can be seen graphically in the person-item threshold distributions ([Figure S2](#)). The item thresholds distributions (blue histogram) were spread over a reasonable range that matched the sample (pink histograms).

TRT data for toenails (n = 95) and fingernails (n = 97) provided 192 assessments for the Appearance, Distress, and Symptoms scales, 140 for the Outcomes scale, 95 for the Physical: Toenails, and 97 for the Physical: Fingernails and Strength: Fingernails scales. TRT participants who reported a change in their nail condition (n = 12) or started a treatment (n = 18) were excluded. The ICC values for all NAIL-Q scales were ≥ 0.81 , which exceeded the COSMIN criteria (see [Table 4](#)).


**Figure 2** Framework for the NAIL-Q.

Table 4 RMT Scale-Level Statistics and Other Scale Findings

Scale	RMT Analysis										Test-Retest Analysis			Floor		Ceiling	
	Sample n	RMT n	Scored on Scale %	χ^2	DF	p-value	PSI +extr	PSI -extr	α +extr	α -extr	ICC	95% LB	95% UB	n	%	n	%
NAIL APPEARANCE	623	598	96.0	108.1	90	0.09	0.93	0.93	0.95	0.94	0.81	0.74	0.86	22	3.5	1	0.2
NAIL DISTRESS	619	589	95.2	67.8	56	0.13	0.89	0.88	0.91	0.90	0.86	0.81	0.89	26	4.2	4	0.6
NAIL SYMPTOMS	622	516	83.0	53.2	24	0.12	0.87	0.86	0.93	0.90	0.84	0.78	0.88	19	3.1	84	13.5
STRENGTH: FINGERNAILS	186	166	89.2	18.5	8	0.02	0.84	0.80	0.89	0.83	0.84	0.76	0.90	10	5.4	9	4.8
PHYSICAL: FINGERNAILS	183	140	76.5	9.2	12	0.69	0.85	0.85	0.93	0.88	0.82	0.73	0.88	1	0.5	40	21.9
PHYSICAL: TOENAILS	435	296	68.1	23.0	15	0.09	0.79	0.79	0.91	0.84	0.89	0.84	0.93	4	0.9	133	30.6
OUTCOME	427	379	88.8	30.2	35	0.70	0.95	0.94	0.97	0.95	0.84	0.77	0.89	26	6.1	21	4.9

Abbreviations: χ^2 , chi square; DF, degrees of freedom; PSI±extr, Person Separation Index with and without extremes; α ±extr, Cronbach alpha with and without extremes; ICC, intraclass correlation coefficient; LB, lower bound; UB, upper bound; floor, % to score at 0 (lowest score); ceiling, % to score 100 (highest score).

Table 5 Participants to Disagree/Agree to NAIL-Q Evaluation Questions

	Disagree		Slightly Agree		Mostly Agree		Strongly Agree		Missing	
	n	%	n	%	n	%	n	%	n	%
1. Asked questions in a respectful way.	11	2.0	33	5.9	135	24.3	363	65.4	13	2.3
2. Easy to understand.	12	2.2	45	8.1	158	28.5	327	58.9	13	2.3
3. Was thorough.	34	6.1	63	11.4	189	34.1	260	46.8	9	1.6
4. Asked questions that are important to me.	57	10.3	107	19.3	176	31.7	203	36.6	12	2.2
5. Should be used by doctors treating people with nail conditions.	48	8.6	139	25.0	179	32.3	181	32.6	8	1.4
6. Would help my doctor better understand my nail condition.	69	12.4	119	21.4	195	35.1	160	28.8	12	2.2

Principal Component Analysis supported a single factor for all 7 scales, with factor loadings ≥ 0.78 to 0.94. Pearson correlations between NAIL-Q scales and the EQ-5D-5L scores met the COSMIN criteria for five scales Appearance ($r = 0.18$, $n = 615$), Distress ($r = 0.31$, $n = 614$), Physical: Fingernails ($r = 0.30$, $n = 182$), Physical: Toenails ($r = 0.40$, $n = 433$), and Outcome ($r = 0.10$, $n = 427$). The two remaining scales had correlations that were weaker than hypothesized (Symptoms, $r = 0.26$, $n = 615$; Strength: Fingernails, $r = 0.10$, $n = 182$).

Table 1 summarizes the construct validity findings, and Tables S6–S11 provide the detailed results. All 6 hypotheses were supported by the data. As nail condition severity increased, NAIL-Q scores were incrementally lower (see Table S6). Those with a nail condition that affected both sides reported lower scores than those whose nail condition was on one side (see Table S7). As pain increased, symptom scale scores decreased (see Table S8). Participants who reported feeling more embarrassed about their nail condition reported worse NAIL-Q scores (see Table S9). Additionally, as interference with activities increased, scores on the NAIL-Q scales decreased (see Table S10). Finally, following treatment, scores on the Outcome scale were higher for those who reported their nail condition as better compared with those whose condition was the same or worse following treatment (see Table S11).

Finally, Table 5 shows the proportion of participants to agree/disagree with the six evaluation questions. Most participants agreed that the NAIL-Q asked questions in a respectful way, was easy to understand, was thorough, asked important questions, should be used by doctors treating people with nail conditions, and would help doctors better understand nail conditions.

Discussion

Nail disorders are common and comprise approximately 10% of all dermatological conditions.⁴³ The NAIL-Q is a rigorously developed and tested PROM that can be used in research and clinical care to measure outcomes that matter to people with nail conditions. The NAIL-Q was developed using a patient-centred approach in line with guidelines for PROM development.¹² Our analysis refined a set of independently functioning scales with strong psychometric properties. The evaluation questions provided additional evidence of content validity by showing the NAIL-Q was easy to understand, asked relevant and important questions in a respectful way, and should be used to inform clinical care.

A recent review assessed the content measured by seven nail-specific PROMs.^{19–25,44} Most items in existing PROMs measure physical and psychological concerns.⁴⁴ While the NAIL-Q includes such concerns, it also covers additional concepts, including a scale to measure satisfaction with nail appearance. In clinical trials of treatments for nail conditions, how the nail looks (eg, how thick, clear, even, healthy it looks) represents a proximal outcome important to patients but is largely overlooked in existing PROMs.⁴⁴ Although, five of the seven existing nail-specific PROMs measure nail appearance,^{19–21,23,25} only the onychomycosis-specific PROM from Lubek et al²⁰ and the OnyCOE-tTM²³ provide a separate score for this construct.^{28,44} While such a concept could be measured using objective measures, today it is important in clinical trials to also measure outcomes from the patient perspective.⁴⁴ In addition to outcomes,

measuring patients' experience of care is integral to providing patient-centered care. Previously developed nail-specific PROMs include only a limited number of items that address treatment outcomes,⁴⁴ with only the psoriasis-specific NAPPA-PBI and onychomycosis-specific Lubek et al²⁰ instrument scoring this construct.²⁵ The NAIL-Q, in contrast, includes a scale that can be used post-treatment for any nail condition to evaluate satisfaction with nail treatment (eg, pleased with result, worth the time/effort, would recommend to others, better than expected). We expect this scale will provide useful feedback to physicians and researchers about different nail treatments from the patient perspective.

An important advantage of the NAIL-Q is that its scales were designed to be used across all nail conditions, with most scales applicable to both fingernails and toenails. Other nail-specific PROMs were designed for use in psoriasis and onychomycosis.^{28,44} Using standardized PROM measurements across nail conditions will allow comparability of results between conditions and across studies and could also help facilitate the implementation of PROMs into clinical practice for purposes such as benchmarking, value-based payment, and shared decision-making.^{18,45–47}

Limitations

Our phase 1 study included only four males. However, this limitation was addressed in our phase 2 study, which included more gender diversity; 43.8% identified as male, and 2.3% identified as nonbinary. Furthermore, the DIF analysis by gender showed that items with DIF had negligible impact on scoring. The qualitative sample to develop the NAIL-Q only included people from Canada and the USA. The field-test sample, on the other hand, was open to anyone fluent in English on the Prolific platform. While a strength of the study was the testing of NAIL-Q in people from multiple countries, a limitation is that we were only able to examine DIF by country for the UK and USA. Additional research to explore the psychometric performance of the NAIL-Q in countries where our sample was limited in size is warranted. Another limitation is that since the phase 2 sample was community-based, it was not possible to confirm nail diagnoses clinically, nor the severity ratings, as both variables were self-reported. Future research could examine the psychometric performance of the NAIL-Q in a clinic-based sample with confirmed diagnoses and objective measures of nail severity. Future research to measure responsiveness (ability to measure change) and to determine minimally important differences for NAIL-Q scales is also needed.

There are many advantages to recruiting research participants using online platforms such as Prolific. Such studies can accrue large diverse international samples quickly and at low cost. Online platforms are ideal for TRT studies as it is possible to control the collection of retest data on a specific day and for a specific number of participants. While crowdsourcing platforms may be a useful means to engage research participants, there are limitations. Many factors can affect data quality, including lack of attention, comprehension, honesty, and reliability.⁴⁸ We used Prolific as it has been shown to provide high-quality data compared to other similar platforms.⁴⁸

Conclusion

The NAIL-Q is a rigorously designed nail-specific PROM developed using a modern psychometric approach.¹² The NAIL-Q can be used to measure outcomes for adults with any fingernail or toenail condition. The NAIL-Q is available on the following website: www.qportfolio.org/nail-q/.

Key Points

- The NAIL-Q was carefully designed using a modern psychometric approach (Rasch Measurement Theory) to measure concepts that are relevant to people with nail conditions.
- Most participants found that the NAIL-Q was easy to understand, asked relevant and important questions in a respectful way, and thought it should be used to inform clinical care.
- The NAIL-Q provides seven independently functioning scales that measure nail appearance, physical and psychosocial concerns, and satisfaction with the treatment that can be used to inform clinical trials, research, and patient care.

Data Sharing Statement

The datasets generated during and/or analysed during the current study are not publicly available due to participant privacy, and legal requirements under data transfer agreements.

Ethics Approval

This study was approved by the Hamilton Integrated Research Ethics Board approvals #7370 and #14622 and from Weill Cornell Medicine (New York, NY, USA) and we certify that the study was performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

Consent for Participation

All participants provided electronic informed consent at the start of the online survey.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This study was funded by research funds provided to A Klassen from the Department of Pediatrics, McMaster University.

Disclosure

The NAIL-Q is owned by McMaster University and Weill Cornell Medicine. The developers could potentially receive a share of any license revenues as royalties based on their institutions' inventor sharing policy. Anne Klassen provides research consulting services to the pharmaceutical industry through EVENTUM Research. The other authors have no conflicts of interest to report for this work.

References

1. Wollina U, Nenoff P, Haroske G, Haenssle HA. The diagnosis and treatment of nail disorders. *Dtsch Arztebl Int.* 2016;113(29–30):509–518. doi:10.3238/arztebl.2016.0509
2. Gupta AK, Mays RR, Versteeg SG, et al. Update on current approaches to diagnosis and treatment of onychomycosis. *Expert Rev Anti Infect Ther.* 2018;16(12):929–938. doi:10.1080/14787210.2018.1544891
3. Lipner SR, Scher RK. Onychomycosis – a small step for quality of care. *Curr Med Res Opin.* 2016;32(5):865–867. doi:10.1185/03007995.2016.1147026
4. Armstrong AW, Harskamp CT, Armstrong EJ. Psoriasis and metabolic syndrome: a systematic review and meta-analysis of observational studies. *J Am Acad Dermatol.* 2013;68:654–662. doi:10.1016/j.jaad.2012.08.015
5. Jiaravuthisan MM, Sasseville D, Vender RB, et al. Psoriasis of the nail: anatomy, pathology, clinical presentation, and a review of the literature on therapy. *J Am Acad Dermatol.* 2007;57(1):1–27. doi:10.1016/j.jaad.2005.07.073
6. Baran R, Dawber RPR, de Berker DAR, Haneke E, Tosti A. *Baran and Dawber's Diseases of the Nails and Their Management.* John Wiley & Sons; 2008.
7. Gupta AK, Versteeg SG, Shear NH. Onychomycosis in the 21st Century: an update on diagnosis, epidemiology, and treatment. *J Cutan Med Surg.* 2017;21(6):525–539. doi:10.1177/1203475417716362
8. Stewart CR, Algu L, Kamran R, et al. The impact of nail psoriasis and treatment on quality of life: a systematic review. *Skin Appendage Disord.* 2021;7(2):83–89. doi:10.1159/000512688
9. Stewart CR, Algu L, Kamran R, et al. Effect of onychomycosis and treatment on patient-reported quality-of-life outcomes: a systematic review. *J Am Acad Dermatol.* 2021;85(5):1227–1239. doi:10.1016/j.jaad.2020.05.143
10. Stewart CR, Algu L, Kamran R, et al. Patient satisfaction with treatment for onychocryptosis: a systematic review. *Skin Appendage Disord.* 2020;6(5):272–279. doi:10.1159/000508927
11. Belyayeva E, Gregoriou S, Chalikias J, et al. The impact of nail disorders on quality of life. *Eur J Dermatol.* 2013;23(3):366–371. doi:10.1684/ejd.2013.2048
12. US Food and Drug Administration. Guidance for industry patient reported outcome measures: use in medical product development to support labeling claims; 2009. Available from: <https://www.fda.gov/media/77832/download>. Accessed October 14, 2023.
13. Snyder C, Crossnohere N, King M, et al. The PROTEUS-Trials Consortium: optimizing the use of patient-reported outcomes in clinical trials. *Clin Trials.* 2022;19(3):277–284. doi:10.1177/17407745221077691
14. Calvert M, Kyte D, Price G, et al. Maximising the impact of patient reported outcome assessment for patients and society. *BMJ.* 2019;364. doi:10.1136/bmj.k5267
15. Snyder CF, Aaronson NK, Choucair AK, et al. Implementing patient-reported outcomes assessment in clinical practice: a review of the options and considerations. *Qual Life Res.* 2012;21(8):1305–1314. doi:10.1007/s11136-011-0054-x
16. Gibbons C, Porter I, Gonçalves-Bradley DC, et al. Routine provision of feedback from patient-reported outcome measurements to healthcare providers and patients in clinical practice. *Cochrane Database Syst Rev.* 2021;(10). doi:10.1002/14651858.CD011589.pub2

17. Maruszczuk K, Aiyegbusi OL, Torlinska B, et al. Systematic review of guidance for the collection and use of patient-reported outcomes in real-world evidence generation to support regulation, reimbursement and health policy. *J Patient Rep Outcomes*. 2022;6(1):1–11. doi:10.1186/s41687-022-00466-7
18. Squitieri L, Bozic KJ, Pusic AL. The role of patient-reported outcome measures in value-based payment reform. *Value Health*. 2017;20(6):834–836. doi:10.1016/j.jval.2017.02.003
19. Drake LA, Patrick DL, Fleckman P, et al. The impact of onychomycosis on quality of life: development of an international onychomycosis-specific questionnaire to measure patient quality of life. *J Am Acad Dermatol*. 1999;41(2):189–196. doi:10.1016/s0190-9622(99)70047-2
20. Lubeck DP, Gause D, Schein JR, et al. A health-related quality of life measure for use in patients with onychomycosis: a validation study. *Qual Life Res*. 1999;8(1):121–129. doi:10.1023/a:1026429012353
21. Turner RR, Testa MA. Measuring the impact of onychomycosis on patient quality of life. *Qual Life Res*. 2000;9(1):39–53. doi:10.1023/a:1026429012353
22. Warshaw EM, Foster JK, Cham PM, et al. NailQoL: a quality-of-life instrument for onychomycosis. *Int J Dermatol*. 2007;46(12):1279–1286. doi:10.1111/j.1365-4632.2007.03362.x
23. Potter LP, Mathias SD, Raut M, et al. The OnyCOE-r™ questionnaire: responsiveness and clinical meaningfulness of a patient-reported outcomes questionnaire for toenail onychomycosis. *Health Qual Life Outcomes*. 2006;4(1):1–8. doi:10.1186/1477-7525-4-50
24. Ortonne JP, Baran R, Corvest M, et al. Development and validation of nail psoriasis quality of life scale (NPQ10). *J Eur Acad Dermatol Venereol*. 2010;24(1):22–27. doi:10.1111/j.1468-3083.2009.03344.x
25. Augustin M, Blome C, Costanzo A, et al. Nail Assessment in Psoriasis and Psoriatic Arthritis (NAPPA): development and validation of a tool for assessment of nail psoriasis outcomes. *Br J Dermatol*. 2014;170(3):591–598. doi:10.1111/bjd.12664
26. Prinsen CAD, Mokkink LB, Bouter LM, et al. COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res*. 2018;27(5):1147–1157. doi:10.1007/s11136-018-1798-3
27. Terwee CB, Prinsen CA, Chiarotto A, et al. COSMIN methodology for evaluating the content validity of patient-reported outcome measures: a Delphi study. *Qual Life Res*. 2018;27(5):1159–1170. doi:10.1007/s11136-018-1829-0
28. Kamran R, Algu L, Leveille CF, et al. Patient-reported outcome measures for patients with nail conditions: a systematic review of the psychometric evidence. *Arch Dermatol Res*. 2022;314(3):223–237. doi:10.1007/s00403-021-02222-1
29. Rasch G. *Studies in Mathematical Psychology: 1. Probabilistic Models for Some Intelligence and Attainment Tests*. Copenhagen, Denmark: Danmarks pædagogiske Institut; 1960.
30. Hobart J, Cano S. Improving the evaluation of therapeutic interventions in multiple sclerosis: the role of new psychometric methods. *Health Technol Assess*. 2009;13:177. doi:10.3310/hta13120
31. Regnault A, Willgoss T, Barbic S; International Society for Quality of Life Research Mixed Methods Special Interest Group. Towards the use of mixed methods inquiry as best practice in health outcomes research. *J Patient Rep Outcomes*. 2018;2(1):1–4. doi:10.1186/s41687-018-0043-8
32. Thorne S, Kirkham SR, MacDonald-Emes J. Interpretive description: a noncategorical qualitative alternative for developing nursing knowledge. *Res Nurs Health*. 1997;20(2):169–177. doi:10.1002/(SICI)1098-240X(199704)20:2<169::AID-NUR9>3.0.CO;2-I
33. Pope C, Ziebland S, Mays N. Qualitative research in health care. Analysing qualitative data. *Br Med J*. 2000;320(7227):114–116. doi:10.1136/bmj.320.7227.114
34. Sandelowski M. Theoretical saturation. In: Given LM, editor. *The Sage Encyclopedia of Qualitative Methods*. Vol. 1. Thousand Oaks, CA: Sage; 2008:875–876.
35. Willis GB. *Cognitive Interviewing in Practice: Think-Aloud, Verbal Probing, and Other Techniques: Cognitive Interviewing*. Thousand Oaks: Sage Publications; 2005:42–65.
36. Collins D. Pretesting survey instruments: an overview of cognitive methods. *Qual Life Res*. 2003;12(3):229–238. doi:10.1023/a:1023254226592
37. Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–381. doi:10.1016/j.jbi.2008.08.010
38. Shaw JW, Johnson JA, Coons SJ. US valuation of the EQ-5D health states: development and testing of the D1 valuation model. *Med Care*. 2005;43:203–220. doi:10.1097/00005650-200503000-00003
39. Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika*. 1951;16:297–334. doi:10.1007/BF02310555
40. Nunnally JC. *Psychometric Theory*. 3rd ed. New York, NY: McGraw-Hill; 1994.
41. Caskin CJ, Happell B. On exploratory factor analysis: a review of recent evidence, an assessment of current practice, and recommendations for future use. *Int J Nurs Stud*. 2014;51(3):511–521. doi:10.1016/j.ijnurstu.2013.10.005
42. Kim HY. Statistical notes for clinical researchers: assessing normal distribution (2) using skewness and kurtosis. *Restor Dent Endod*. 2013;38(1):52–54. doi:10.5395/rde.2013.38.1.52
43. Bansal Goyal N, Chavan RB, Belgaumkar VA. Clinico-etiological study of nail disorders at a tertiary care center in Maharashtra, India. *J Skin Stem Cell*. 2021;8(3):e119437. doi:10.5812/jssc.119437
44. Abid K, Algu L, Kamran R, et al. Content analysis of patient-reported outcome measures used in patients with nail conditions: a systematic review. *JAMA Dermatol*. 2021;157(12):1509–1511. doi:10.1001/jamadermatol.2021.4539
45. Mercieca-Bebber R, King MT, Calvert MJ, et al. The importance of patient-reported outcomes in clinical trials and strategies for future optimization. *Patient Relat Outcome Meas*. 2018;1:353–367. doi:10.2147/PROM.S156279
46. Withers KL, Puntoni S, O’Connell S, et al. Standardising the collection of patient-reported experience measures to facilitate benchmarking and drive service improvement. *Patient Exp J*. 2018;5(3):16–24. doi:10.35680/2372-0247.1268
47. Noonan VK, Lyddiatt A, Ware P, et al. Montreal accord on patient-reported outcomes (PROs) use series—Paper 3: patient-reported outcomes can facilitate shared decision-making and guide self-management. *J Clin Epidemiol*. 2017;89:125–135. doi:10.1016/j.jclinepi.2017.04.017
48. Peer E, Rothschild D, Gordon A, et al. Data quality of platforms for online behavioral research. *Behav Res Methods*. 2022;54(4):1643–1662. doi:10.3758/s13428-021-01694-3

Clinical, Cosmetic and Investigational Dermatology

Dovepress

Publish your work in this journal

Clinical, Cosmetic and Investigational Dermatology is an international, peer-reviewed, open access, online journal that focuses on the latest clinical and experimental research in all aspects of skin disease and cosmetic interventions. This journal is indexed on CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dermatology-journal>