




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Chronic Related Group classification system as a new public health tool to predict risk and outcome of COVID-19 in patients with systemic rheumatic disease: a population-based study of more than forty thousand patients



Enrico De Lorenzis Paolo Parente Gerlando Natalello Salvatore Soldati Silvia Laura Bosello Andrea Barbara Chiara Sorge Svetlana Axelrod Lucrezia Verardi Pier Giacomo Cerasuolo Giusy Peluso Antonella Gemma Marina Davoli Donatella Biliotti Vincenzo Bruzzese Mauro Goletti Mirko Di Martino Maria Antonietta D'Agostino

PII: S1297-319X(22)00157-9

DOI: <https://doi.org/doi:10.1016/j.jbspin.2022.105497>

Reference: BONSOI 105497

To appear in: *Joint Bone Spine*

Received Date: 1 July 2022

Please cite this article as: Lorenzis ED, Parente P, Natalello G, Soldati S, Bosello SL, Barbara A, Sorge C, Axelrod S, Verardi L, Cerasuolo PG, Peluso G, Gemma A, Davoli M, Biliotti D, Bruzzese V, Goletti M, Martino MD, D'Agostino MA, Chronic Related Group classification system as a new public health tool to predict risk and outcome of COVID-19 in patients with systemic rheumatic disease: a population-based study of more than forty thousand patients, *Joint Bone Spine* (2022), doi: <https://doi.org/10.1016/j.jbspin.2022.105497>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier.

Chronic Related Group classification system as a new public health tool to predict risk and outcome of COVID-19 in patients with systemic rheumatic disease: a population-based study of more than forty thousand patients

Enrico De Lorenzis^{1,2}, Paolo Parente¹, Gerlando Natalello², Salvatore Soldati³, Silvia Laura Bosello², Andrea Barbara^{1,6}, Chiara Sorge³, Svetlana Axelrod⁴, Lucrezia Verardi², Pier Giacomo Cerasuolo², Giusy Peluso², Antonella Gemma¹, Marina Davoli³, Donatella Biliotti¹, Vincenzo Bruzzese^{1,5}, Mauro Goletti¹, Mirko Di Martino^{3*}, Maria Antonietta D'Agostino^{2*}

*Mirko Di Martino and Maria Antonietta D'Agostino equally contributed to this work.

1. Local public health office Roma 1 - Rome, Italy
2. Division of Rheumatology, Catholic University of the Sacred Heart, Fondazione Policlinico Universitario A. Gemelli IRCCS - Rome, Italy
3. Department of Epidemiology, Lazio Regional Health Service - Rome, Italy
4. Institute of Leadership, University of Sechenov, - Moscow, Russia
5. Department of Internal Medicine, Rheumatology and Gastroenterology, Nuovo Regina Margherita Hospital - Rome, Italy
6. PhD in Infectious Diseases, Microbiology and Public Health, Cycle XXXVI, Department of Public Health and Infectious Disease, Sapienza University of Rome - Rome, Italy

Correspondence to Professor Maria-Antonietta D'Agostino - Division of Rheumatology, Università Cattolica del Sacro Cuore, Policlinico Universitario Agostino Gemelli IRCCS, 00187 Roma, Italy;

mariaantonietta.dagostino@unicatt.it

Keywords: COVID-19, systemic autoimmune rheumatic disease, epidemiology, public health.

Since the COVID-19 outbreak, public health authorities have looked for the best evidence on infection risk and prognosis to guide their choices. The interpretation of the observational studies that variably reported increased infection rates and a poor prognosis in patients with systemic rheumatic diseases (SRD) has been limited by factors such as the selection of patients in the care of tertiary referral centres, the small available sample sizes for the less prevalent diseases, the description of SRDs as a single broad category, and the neglected influence of comorbidities [1].

The use of big healthcare data has become essential in gathering crucial information for a reliable identification of high-risk groups. The Chronic Related Group (CReG) system is an experimental approach of classification of chronic patients created to predict the medical resources needed to ensure their care. This system automatically assigns a diagnosis to a subject according to medical administrative records over a pre-set period. Specifically, CReG system relies on the registration and integration of disease-specific codes used to determine the share of healthcare costs, hospital discharge diagnoses codes and access to the prescription of drugs or therapeutic procedures uniquely associated with a specific condition [2,3].

In this analysis, we compared incidences and 30-day outcomes of 40490 SRD patients (Table 1) to 4716119 subjects dwelling in the Lazio Region, the second most populated region of Italy that includes the Rome metropolitan area. SRDs and comorbidity diagnoses were derived from the CReG classification while data on COVID-19 infection from a regional digital network. The risk was expressed as incidence rate ratio adjusted for demographics and comorbidities. We focused on the period from the 20th of February 2020 to the 31st of December 2020 to selectively assess a cohort of unvaccinated patients.

Table 2 reports peculiar patterns in terms of incidence, hospitalisation, intensive care unit (ICU) admission and death for the different SRDs. COVID-19 risk was increased in patients with Psoriatic Arthritis and Undifferentiated Connective Tissue Disease, possibly as the result of reduced adherence to protecting behaviours. These conditions are indeed less frequently treated with immunosuppressants or associated with visceral involvement, circumstances that have been reported to lead to highly perceived individual risk [4,5,6]. The hospitalization risk was higher in patients with Axial Spondylarthritis, Systemic Erythematosus Lupus (SLE), Systemic Vasculitis, intensive care unit (ICU) admission risk was higher in Systemic Erythematosus Lupus and primary Sjögren's syndrome patients, while increased mortality was reported in patients with Rheumatoid Arthritis, SLE, primary Sjögren Syndrome, and Scleroderma. It can be argued that the patients who are more likely to present pulmo-renal complications are more susceptible to worse outcomes. The high prevalence of lung fibrosis and the specific vasculopathy could explain the especially high mortality in scleroderma patients.

In conclusion, we showed how the CReG system classification allows the identification of high-risk SRD patients on a large scale and highlights the heterogeneity in their clinical behaviours. This methodology could be fruitfully extended to the assessment of other potential SRD-related complications such as cancer and cardiovascular events.

References

1. Grainger R, Kim AHJ, Conway R, Yazdany J, Robinson PC. COVID-19 in people with rheumatic diseases: risks, outcomes, treatment considerations. *Nat Rev Rheumatol*. 2022;18(4):191-204. doi:10.1038/s41584-022-00755-x
2. Sorlini M, Perego L, Silva S, Marinoni G, Zucchi A, Barbaglio G, et al. The chronic related groups (CReG) model for ensuring continuity of care for chronically ill patients: pilot experience of the planning stage in Bergamo (Italy). *Ig Sanita Pubbl*. 2012;68(6):841-861
3. Rea F, Corrao G, Ludergrani M, Cajazzo L, Merlino L. A new population-based risk stratification tool was developed and validated for predicting mortality, hospital admissions, and health care costs. *J Clin Epidemiol*. 2019;116:62-71. 9
4. Kipps S, Paul A, Vasireddy S. Incidence of COVID-19 in patients with rheumatic disease: is prior health education more important than shielding advice during the pandemic?. *Clin Rheumatol*. 2021;40(4):1575-1579.
5. Cook C, Cox H, Fu X, Zhang Y, Stone JH, Choi HK, et al. Perceived Risk and Associated Shielding Behaviors in Patients With Rheumatoid Arthritis During the Coronavirus 2019 Pandemic [published online ahead of print, 2021 Sep 8]. *ACR Open Rheumatol*. 2021;10.1002/acr2.11340.0.
6. Mahil SK, Yates M, Langan SM, Yiu ZZN, Tsakok T, Dand N, et al. Risk-mitigating behaviours in people with inflammatory skin and joint disease during the COVID-19 pandemic differ by treatment type: a cross-sectional patient survey. *Br J Dermatol*. 2021;185(1):80-90.

Table 1: Prevalence of SRDs in Lazio and demographics of the affected patients

SRD Diagnosis	Total Patients n	Prevalence %	Overlapping SRDs n (%)	Female n (%)	Age 18-35 n (%)	Age 36-59 n (%)	Age 60-79 n (%)	Age ≥80 n (%)
RA	14838	0.31	1463 (9.86)	11581 (79.05)	1005 (0.10)	4755 (0.23)	7243 (0.58)	1835 (0.46)
PsA	11273	0.24	867 (7.72%)	6862 (61.07)	611 (0.06)	5192 (0.25)	4915 (0.40)	519 (0.13)
axSpA	2422	0.05	305 (12.59)	1118 (46.16)	255 (0.03)	1276 (0.06)	763 (0.06)	128 (0.03)
SLE	3936	0.08	682 (17.33)	3388 (86.08)	412 (0.04)	2261 (0.11)	1141 (0.09)	122 (0.03)
pSS	3393	0.07	989 (29.15)	3208 (94.55)	114 (0.01)	1324 (0.06)	1628 (0.13)	327 (0.08)
SSc	2256	0.05	522 (23.14)	2056 (91.13)	109 (0.01)	836 (0.04)	1107 (0.09)	204 (0.05)
MSD	952	0.02	296 (31.09)	797 (83.72)	83 (0.01)	418 (0.02)	405 (0.03)	46 (0.01)
UCTD	2977	0.06	695(23.25)	2745 (92.21)	386 (0.04)	1673 (0.08)	853 (0.07)	65 (0.02)
SV	1560	0.03	118 (7.56)	978 (62.69)	206 (0.02)	560 (0.03)	610 (0.05)	184 (0.05)

Abbreviations: SRD Systemic Rheumatic Disease, SD Standard Deviation, RA Rheumatoid Arthritis, PsA Psoriatic Arthritis, axSpA Axial Spondylarthritis, SLE Systemic Lupus Erythematosus, pSS primary Sjögren Syndrome, SSc Systemic Sclerosis, MSD Myositis-Spectrum Disorders, UCTD Undifferentiated Connective Tissue Disease, SV Systemic Vasculitis.

Table 2: Infection rates and thirty-day hospitalisation, ICU admission, and death rates in SRD patients with COVID-19

Diagnosis	n	Tested patients n (%)	Infection		Hospitalization		ICU admission		Death	
			Cases n	Adjusted IRR (95% CI)	Cases n	Adjusted IRR (95% CI)	Cases n	Adjusted IRR (95% CI)	Cases n	Adjusted IRR (95% CI)
RA	13375	3218 (24.06)	453	1.09 (0.99-1.19)	94	1.18 (0.96-1.45)	16	1.47 (0.90-2.41)	30	1.50 * (1.04-2.17)
PsA	10370	2747 (26.49)	394	1.21 *** (1.10-1.33)	60	1.15 (0.89-1.48)	12	1.54 (0.87-2.72)	9	0.89 (0.46-1.73)
axSpA	2117	575 (27.16)	84	1.21 (0.98-1.50)	15	1.89 * (1.14-3.14)	1	0.82 (0.12-5.82)	2	1.02 (0.25-4.12)
SLE	3254	908 (27.90)	128	1.14 (0.96-1.36)	24	2.16 *** (1.45-3.22)	5	3.67 ** (1.52-8.83)	5	2.67 * (1.10-6.44)
pSS	2404	640 (26.62)	84	1.12 (0.90-1.38)	17	1.58 (0.98-2.54)	5	4.13 ** (1.71-9.96)	6	2.51 * (1.12-5.62)
SSc	1734	528 (30.45)	46	0.84 (0.63-1.12)	10	1.23 (0.66-2.31)	0	-	6	4.60 *** (2.06-10.29)
MSD	656	170 (25.91)	22	1.03 (0.68-1.56)	4	1.78 (0.67-4.74)	0	-	0	-
UCTD	2282	726 (31.81)	95	1.26 * (1.03-1.54)	10	1.45 (0.78-2.70)	0	-	0	-
SV	1442	409 (28.35)	53	0.99 (0.75-1.30)	14	1.81 * (1.07-3.06)	2	1.33 (0.33-5.38)	4	2.31 (0.86-6.18)
Overlapping SRDs	2856	836 (29.27)	100	1.06 (0.87-1.28)	20	1.79 ** (1.16-2.78)	6	3.96 *** (1.78-8.84)	5	2.45 * (1.02-5.91)
No SRDs	4675629	1054387 (22.55)	149092	1.00 (0.99-1.00)	17517	1.00 (0.98-1.02)	2570	0.99 (0.93-1.05)	3821	0.99 (0.92-1.08)
Lazio Region	4716119	1065144 (22.59)	150551	-	17785	-	2617	-	3888	

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: SRD Systemic Rheumatic Disease, SD Standard Deviation, RA Rheumatoid Arthritis, PsA Psoriatic Arthritis, axSpA Axial Spondylarthritis, SLE Systemic lupus erythematosus, ICU intensive care unit, Lupus Erythematosus, pSS primary Sjögren Syndrome, SSc Systemic Sclerosis, MSD Myositis-Spectrum Disorders, UCTD Undifferentiated Connective Tissue Disease, SV Systemic Vasculitis.