

## Novel Use for Hemostatic Net in Split-thickness Skin Grafts

Liz Carolina Pava, MD\*; Juana Mendoza, MD†; Diana Martínez, MD‡; Stefania Rodríguez-Bermeo, MD‡; Valeria Barrios, MD‡; Viviana Gómez-Ortega, MD‡†

Skin grafts are a widely known and used technique in the field of plastic surgery.<sup>1</sup> According to the International Burn Injury Database, there are 1000 skin grafts in patients with burns per year in England.<sup>2</sup> The most common cause of skin graft failure is hematoma, followed by infection, seroma, and the presence of shearing forces.<sup>1</sup> Numerous methods for fixation with tie-over bolster, negative pressure, tissue glue, and others are found in the literature because immobilization is essential to prevent graft loss.<sup>3</sup> However, there is a high rate of graft loss that is still unreported.

The hemostatic net was first mentioned in cosmetic procedures to reduce hematoma in the early postoperative period.<sup>4</sup> This is achieved because the suture performed provides an almost total occlusion of the surgical area and avoids fluid accumulation.<sup>5</sup> However, the use of this procedure has not been described in split-thickness skin grafts in burn patients.

We used a modified hemostatic net technique to favor optimal postsurgical results in acutely burned patients. This technique consists of a series of continuous and transfixing sutures with Monocryl 3-0, 0.5 cm apart from each other, until achieving a net shape in the whole grafted area (Fig. 1). (See Video, Supplemental Digital Content 1, which displays a plastic surgeon performing hemostatic net over a split thickness skin graft in a burn patient.) The grafts are bolstered with the conventional technique. We considered that the fixation technique with internal sutures can be performed with other types of absorbable suture material, and achieve results as good as those using Monocryl.

The concept of internal sutures was first described by Barouide and Ferreira in abdominoplasties in 1998, as a technique that avoids the eventual fluid deposits.<sup>4</sup> In 2012 based on the concept of mechanical closure, Auersvald et al described a surgical procedure



Fig. 1. Hemostatic net in a split-thickness skin graft in a burn patient.

of transcutaneous continuous and transfixing sutures of skin flaps in rhytidoplasty, creating a hemostatic net that leads to almost total occlusion of the treated area.<sup>5</sup> They found that this technique was an efficient method for the prevention of early hematomas, without an increase in ischemia or necrosis of flaps.<sup>4</sup> In 2018, the same authors described the use of this method in facelift surgery, with the conclusion that hemostatic net is a safe and effective method to reduce hematoma in the early postoperative period by not altering the perfusion of the flaps.<sup>4</sup>

Further research is needed to better define the advantage of the hemostatic net with the suggested modifications in this field. We propose a novel indication for a modified hemostatic net technique as an effective method for prevention of hematomas and seromas in skin grafts.

From the \*Plastic and Reconstructive Surgery, Universidad del Bosque, Bogotá, Colombia; †Burn Unit Fundación Santa Fé de Bogotá, Universidad de los Andes, Bogotá, Colombia; and ‡Plastic and Reconstructive Surgery Department, Fundación Santa Fé de Bogotá University Hospital, Universidad de los Andes, Bogotá, Colombia.

Received for publication July 30, 2021; accepted September 7, 2021. Copyright © 2021 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal. *Plast Reconstr Surg Glob Open* 2021;9:e3893; doi: 10.1097/GOX.0000000000003893; Published online 26 October 2021.

Related Digital Media are available in the full-text version of the article on [www.PRSGlobalOpen.com](http://www.PRSGlobalOpen.com).

*Viviana Gómez-Ortega, MD*

Plastic and Reconstructive Surgery  
 Department and Burn Unit  
 Fundación Santa Fé de Bogotá  
 Universidad de los Andes  
 Bogotá, Colombia  
 Carrera 7 No. 117-15  
 Bogotá 111071, Colombia  
 E-mail: [Viviana.gomez@fsfb.org.co](mailto:Viviana.gomez@fsfb.org.co)

#### DISCLOSURE

*The authors have no financial interest to declare in relation to the content of this article.*

#### REFERENCES

1. Khan AA, Khan IM, Nguyen PP, et al. Skin graft techniques. *Clin Podiatr Med Surg.* 2020;37:821–835.
2. Hollén L, Greenwood R, Kandiyali R, et al. The SILKIE (Skin grafting Low friction Environment) study: a non-randomised proof-of-concept and feasibility study on the impact of low-friction nursing environment on skin grafting success rates in adult and paediatric burns. *BMJ Open.* 2018;8:e021886.
3. Yen YH, Lin CM, Hsu H, et al. Skin graft fixation using hydrofiber (Aquacel Extra). *Ann Plast Surg.* 2018;80:616–621.
4. O'Daniel T, Auersvald A, Auersvald L. Hemostatic net in facelift surgery. *Der MKG-Chirurg.* 2019;12:78–85.
5. Auersvald A, Auersvald LA, Biondo-Simões MLP. Hemostatic net: an alternative for the prevention of hematoma rhytidoplasty. *Rev Bras Cir Plast.* 2012;27:22–30.