



# Spiroconjugated 1,2,3-triazolo[5,1-*b*]1,3,4-thiadiazine stimulates functional activity of fibroblasts under skin injury regeneration

Irina M Petrova<sup>1</sup>, Sofya Iu Chebanova<sup>1</sup>, Sergey L Khatsko<sup>1,2,\*</sup>, Tatyana A Kalinina<sup>1</sup>,  
Dmitry V Zaitsev<sup>1,3</sup>, and Tatyana V Glukhareva<sup>1</sup>

<sup>1</sup>Ural Federal University named after the first President of Russia B. N. Yeltsin, Yekaterinburg, 620026, Russia

<sup>2</sup>Federal State Budgetary Scientific Institution "Ural Federal Agrarian Scientific Research Centre, Ural Branch of Russian Academy of Sciences", Yekaterinburg, 620142, Russia.

<sup>3</sup>Ural State Mining University, Yekaterinburg, 620144, Russia.

## Abstract

**Background and purpose:** One of the most important mechanisms of tissue regeneration is the high functional activity of cells, including proliferation. Currently, there are practically no effective skin cell activators on the pharmaceutical market. The purpose of this work was to demonstrate the stimulating effect of spiroconjugated 1,2,3-triazolo[5,1-*b*]1,3,4-thiadiazine (STT) on the functional activity of fibroblasts.

**Experimental approach:** STT containing ointment for dermal application was made. To assess *in vivo* effect of the STT a linear wound model in rats was tested. A combination of histological techniques and mechanical testing was employed to estimate the stimulating effect of STT on the functional activity of fibroblasts.

**Findings/Results:** The STT significantly increased the number of fibroblasts as well as the density and order of produced collagen fibers in the dermis during the wound healing process. As a result, a tissue was formed at the site of damage with the structure corresponding to normal skin. In addition, skin functions were restored, in particular mechanically.

**Conclusion and implications:** The results suggested the stimulating effect of the STT on fibroblast activity and demonstrated its potential for skin regeneration.

**Keywords:** Collagen; Fibroblasts, Ki-67; Mast cells; Skin wound; 1,2,3-Triazolo-1,3,4-thiadiazine; Vimentin.

## INTRODUCTION

Skin injury is an acute issue in modern medicine. Damage skin is characterized by changes in morphology and loss of functionality. Successful cutaneous wound repair requires a series of tightly coordinated steps including inflammation, tissue formation, and remodeling (1-3). Increasingly, some studies reveal the role of different tissue elements' interaction in regeneration. It is known that fibroblasts are the most important element of granulation tissue. These cells determine the processes of formation and remodeling of new tissue. Fibroblasts are also significant because they coordinate the activities of other cells in different phases of regeneration. The fibroblasts' involvement in recovery processes is determined by the level of their proliferative and synthetic activity (1,2,4). Restoration of the skin fibrous component is also

crucial because it provides the mechanical characteristics of the skin and maintains dermis architectonics (5).

Because of the regeneration complexity a failure of wound healing takes place. The repair process mainly becomes recurrently deficient due to extended trauma, infections, and inflammation (6). Despite advances in skin wound treatment, a therapeutic effect is not always achieved. In this line, extensive research is underway, focused on searching for regeneration stimulators, so far, the number of effective substances is limited to a few examples. Standard wound healing therapy relies mainly on proliferation activators of natural origin. For example, to accelerate the regeneration Solcoseryl<sup>®</sup> is widely used (7).

### Access this article online



Website: <http://rps.mui.ac.ir>

DOI: 10.4103/RPS.RPS\_74\_23

\*Corresponding author: S.L. Khatsko  
Tel: +7-9097048774, Fax: +7-3433899703  
Email: [sergey.khatsko@urfu.ru](mailto:sergey.khatsko@urfu.ru)

Chitosan-based products showed reduced inflammation, improved angiogenesis, and accelerated epithelialization (8). The popular substance methyluracil also has a stimulating effect on cell proliferation and microcirculation (9).

Previously, compounds that selectively accelerate the proliferation of human skin fibroblasts *in vitro* were discovered (10). It was found that ethyl 5-(4-ethoxybenzoyl)-5,7-dihydrospiro[1,2,3]triazolo[5,1-*b*][1,3,4]thiadiazine-6, 1'-cyclopentane]-3-carboxylate (STT) stimulates skin regeneration (5,11). However, the way to achieve this effect is still unclear. In this paper, we evaluated regenerative process parameters that can indicate a possible mechanism of STT action and help determine further approaches to its study.

## MATERIALS AND METHODS

### Animals

The experiment utilized 63 male outbred laboratory rats ( $434.7 \pm 8.8$  g) and complied with the Principles of Laboratory Animal Care. All the experiments were permitted by the Ethics Committee of the Institute of Natural Sciences and Mathematics, UrFU, Russia (Ethic No. 2/2022).

Animals were randomly divided into 4 groups, including 5 (morphological studies) or 4 (mechanical studies) individuals: (1) intact rats (INT); (2) untreated rats as the control group (CONTR); (3) rats exposed to vaseline (VAS); and (4) rats exposed to STT (STT).

### Chemicals

Vaseline and Analgin were obtained from Samaramedprom JSC (Russia) and Pharmstandard-UfaVITA JSC (Russia), respectively. The synthesis of STT was carried out at the Institute of Chemical Engineering (UrFU, Russia) (12).

### Ointment preparing

Ointment for skin application was prepared by dissolving 208 mg of the 1,2,3-triazolo-1,3,4-thiadiazine in 100  $\mu$ L of dimethyl sulfoxide, adding this solution to 50 g of Vaseline, and stirring to a homogeneous

consistency. To avoid the effects of ointment base, a group of animals (VAS) were exposed to it. Vaseline was chosen due to its beneficial properties for wound healing (13,14).

### Experimental procedures


The study included linear wound modeling. The skin from the back was prepared by hair plucking and rinsed with ethanol 96%. A linear wound of  $50 \pm 2$  mm long was created using a surgical scalpel. Before the wound was applied, rats were injected intramuscularly with 0.2 mL Analgin 50%.

After wound modeling, animals were exposed to an experimental ointment (0.4% of STT) or Vaseline. Ointments were applied epicutaneously at the dose of 0.2 g once a day for 14 days. Animals were taken out of the experiment on the 14<sup>th</sup> and 21<sup>st</sup> days by an overdose of diethyl ether.

### Histological and immunohistochemical analysis

A strip of skin tissue was obtained and fixed in 10% neutral formalin. Tissue specimens were embedded in paraffin wax, sectioned (4  $\mu$ m), and stained with hematoxylin-eosin (H&E), picrofuchsin (Van Gieson's), and Azure B, to visualize the neotissue formation, collagen deposition, and mast cells number, respectively.

To estimate the cell proliferation intensity and fibroblast synthetic activity, immunohistochemical staining with antibodies against Ki-67 (SP6, rabbit monoclonal antibody, Cell Marque, USA) and vimentin (V9, mouse monoclonal antibody, Cell Marque, USA) was performed according to manufacturer's protocols. Briefly, 4- $\mu$ m-thick sections were placed on poly-L-lysine-coated adhesive microscope slides (Thermo Scientific-Menzel, Germany) and then dewaxed in xylene and rehydrated in graded alcohols. Sections were incubated in Trilogy™ (Cell Marque, USA) to antigens unmasking for 20 min at 92 °C and in H<sub>2</sub>O<sub>2</sub> solution for 10 min at room temperature. After that, sections were washed in tris buffered saline (TBS) immunohistochemistry wash buffer + Tween® 20 (Cell Marque, USA) for 1 min and incubated with primary antibodies for 60 min at 30 °C.



**Fig. 1.** Mechanical test.

Sections were washed three times with TBS before incubation with secondary antibodies (N-Histofine® Simple Stain MAX PO (MULTI), Nichirei Biosciences INC, Japan) for 30 min at 30 °C. Sections were washed three extra times in TBS and coated with DAB chromogen (DAB Substrate Kit, Cell Marque, USA), and incubated for approximately 30-60 s. After that sections were washed in distilled water and stained with hematoxylin.

Zeiss Primostar provided histological analysis. The average vessels and cell number (fibroblasts, mast cells, Ki-67- and vimentin-positive cells), collagen fibers thickness, and vessel area were analyzed by TopView software according to 10 randomly selected fields with conversion to 1 mm<sup>2</sup>. Mast cells were defined as dark purple cells with granular cytoplasm (stained with Azur B). Fibroblasts were defined as cells with a branched basophilic cytoplasm surrounding an elliptical nucleus (stained with H&E). Cells with a brown-stained nucleus and yellow-stained cytoplasm were considered Ki-67 positive and vimentin-positive, respectively.

### **Mechanical tests**

Mechanical tests were conducted on a Shimadzu AGX 50kN tensile testing machine (Japan). The analysis of the sample's mechanical strength was carried out using the method for assessing the deformation behavior of the material under uniaxial compression (15,16). According to a special cardboard stencil, skin samples were cut out. The sample

edges were rigidly fixed in the grips of the testing machine (7). After that, a uniaxial tensile load was applied to the sample (Fig. 1). The time of one test was on average  $71.342 \pm 4.211$  s. Upon reaching the point of maximum mechanical stress and the onset of irreversible deformation (ruptures), the tests were stopped.

### **Statistical analysis**


All data were expressed as the mean  $\pm$  SEM. Statistical analysis was performed using Statistica 12.0 software by Kruskal-Wallis test with Dunn's post-hoc analysis. *P*-values  $< 0.05$  were considered statistically significant. Mathematica software (Wolfram language) was used to visualize the received data.

## **RESULTS**

### **H&E and Azure B staining**

The study aimed to evaluate the effect of the chemical compound STT on the fibroblast's functional activity under linear skin wound regeneration. The points of the experiment were chosen to assess the skin condition at the final stages of regeneration.

On the 14<sup>th</sup> day, intense fibroblast proliferation was observed in the STT group. Acanthoses were detected, but on the 21<sup>st</sup> day, the epidermis acquired a uniform thickness. The dermis was divided into papillary and reticular layers. Numerous hair follicles as well as glandular structures were found. The scar was characterized by maturity with a fibrous component predominance (Fig. 2A).



**Fig. 2.** (A) Hematoxylin and eosin staining, magnification: 100 ×; (B) the number of fibroblasts; (C) Azure B staining, magnification: 400 ×; (D) the number of mast cells. Data are presented mean ± SEM, n = 5. \*\**P* < 0.01 Indicates significant differences in comparison with the INT group; and #*P* < 0.05 between the indicated groups. INT, Intact; CONTR, control; VAS, vaseline; STT, spiroconjugated 1,2,3-triazolo[5,1-b]1,3,4-thiadiazine.


The fibroblast number increased in STT groups by the 14<sup>th</sup> day relative to the CONTR and VAS groups (Fig. 2B). The high content of cells persisted by the 21<sup>st</sup> day. Changes were noted in the mast cell population (Fig. 2C). On the 14<sup>th</sup> day, the number of mast cells significantly increased in the STT group comparing other groups (Fig. 2D).

### Immunohistochemical staining


The cell implication in regeneration is determined mainly by their proliferative activity (4,5). To assess the intensity of cell proliferation, the Ki-67 marker was used (17). Ki-67 positive cells were determined in the epidermis germ layer, and most of the cells were observed in the STT group on the 14<sup>th</sup> day (Fig. 3A), which was significantly higher than the values of the CONTR and VAS groups. It was typical that in the earlier period of regeneration, there were more dividing cells (Fig. 3B). However, in the

STT group, the number of cells remained significantly high even by the 21<sup>st</sup> day and it also applied to the dermis (Fig. 3D).

Fibroblasts are an important link that unites various processes during regeneration (5). *Via* vimentin production, fibroblasts coordinate cell proliferation, collagen accumulation, keratinocyte transdifferentiation, and re-epithelialization (4,18-20). The number of vimentin-positive dermal cells on the 14<sup>th</sup> day of the experiment was high (Fig. 4A and B), which correlated with the fibroblasts number (Fig. 2B). By the 21<sup>st</sup> day, the number of vimentin-positive cells sharply decreased in the CONTR and VAS groups, so it was significantly less than the values of the STT group on the 14<sup>th</sup> day. The microvasculature normalization served as reliable marker of successful regeneration. Here, on the 21<sup>th</sup> day of the experiment we observe normal indicators of vessels number and area (Fig. 4C and D).



**Fig. 3.** (A) Immunohistochemical staining of Ki-67 in the epidermis, magnification: 400 ×; (B) number of Ki-67 positive cells in the epidermis; (C) immunohistochemical staining of Ki67 in the dermis, magnification: 400 ×; (D) number of Ki-67 positive cells in the dermis. Data are presented mean ± SEM, n = 5. <sup>\*\*</sup>*P* < 0.01 and <sup>\*\*\*</sup>*P* < 0.001 indicate significant differences in comparison with the control group; <sup>##</sup>*P* < 0.01 versus VAS group. INT, Intact; CONTR, control; VAS, vaseline; STT, spiroconjugated 1,2,3-triazolo[5,1-b]1,3,4-thiadiazine.




**Fig. 4.** (A) Immunohistochemical staining of vimentin in the dermis, magnification: 400 ×; (B) the number of vimentin-positive cells in the dermis; (C) mean vessels area; and (D) vessels number. Data are presented mean ± SEM, n = 5. <sup>\*\*\*</sup>*P* < 0.001 indicates a significant difference in comparison with the respective control group; <sup>###</sup>*P* < 0.001 versus VAS group; <sup>@@@</sup>*P* < 0.001 against INT group. INT, Intact; CONTR, control; VAS, vaseline; STT, spiroconjugated 1,2,3-triazolo[5,1-b]1,3,4-thiadiazine.

### Van Gieson's staining

Assessment of the skin's fibrous component state can serve as a marker of both the successful restoration of the skin's mechanical functions and the normal scar formation. Collagen deposition became more abundant with prolongation of the time after injury. By the 21<sup>st</sup> day in the STT group, most of the dermis was occupied by collagen fibers. The papillary region was composed of thin fibers collected in loose bundles. The reticular dermis represented by thick and crimped fibers formed certain direction bundles. The fiber dye was more intense in the CONTR and VAS groups. The morphology and distribution of collagen fibers approximated those in the normal dermis (Fig. 5A). The collagen fiber thickness showed no significant differences between the INT and STT groups on the 14<sup>th</sup> day. By the 21<sup>st</sup> day in the STT group fiber bundles were significantly thicker in comparison with the CONTR group (Fig. 5B).

### Mechanical tests

Skin characteristics such as resistance to deformation or failure under applied loads can be determined from mechanical tests (16,21). Mechanical tests revealed a general trend in the dependence of the collagen fiber thickness on the skin strength properties. Therefore, on the 14<sup>th</sup> day, all samples had weak mechanical characteristics that were significantly different from the intact value (Fig. 5C). By the 21<sup>st</sup> day, the mechanical properties of the skin samples improved. STT group had the highest strength characteristics: at a maximum load of  $4.903 \pm 1.554$  MPa, skin stretched by  $34.874 \pm 4.829\%$ , which significantly differed from the values of the CONTR group (at  $3.772 \pm 0.640$  MPa, samples stretched by  $49.673 \pm 14.06\%$ ) and approached the INT values (at  $4.844 \pm 0.206$  MPa the percentage of stretch was  $33.36 \pm 4.92$ ) (Fig. 5D).



**Fig. 5.** (A) Van Gieson's staining, magnification: 400 ×; (B) thickness of collagen fiber bundles; (C) mechanical tests on the 14th day of the experiment; (D) mechanical tests on the 21<sup>st</sup> day of the experiment. Data are presented mean ± SEM, n = 4. \*\**P* < 0.01 and \*\*\**P* < 0.001 indicates significant difference in comparison with the INT group; ##*P* < 0.01 versus control group. INT, Intact; CONTR, control; VAS, vaseline; STT, spiroconjugated 1,2,3-triazolo[5,1-b]1,3,4-thiadiazine.

## DISCUSSION

The results of this study were consistent with the previously described effect of STT associated with the activation of fibroblasts and scar formation with a developed fibrous component (11). Here, additional parameters were evaluated that can more fully describe the state of the tissue during the regeneration and indicate the possible mechanisms of STT action. On the 14<sup>th</sup> day, the epidermis in the STT group was characterized by acanthoses, which were associated with epithelium remodeling. Upon injury, keratinocytes at the wound edges form epithelial tongues that move and interact with the dermal cells and matrix to re-establish coverage of the wound bed (3,22). In the STT group, this process was probably more intense, which was confirmed by the high content of Ki-67-positive cells in the epidermis. In addition, a more intensive recovery was observed by the skin derivatives restoration, which could also be associated with increased proliferation of keratinocytes.

In comparison with the previous study, it was found that the fibroblasts number decreased only by the 21<sup>st</sup> day of the experiment, which may indicate an active restructuring of the scar. The presence of a large number of proliferating cells was correlated with the number of Ki-67-positive cells. Furthermore, a high number of vimentin-positive cells in the STT group may indicate a high functional activity of cells, in particular synthetic. Vimentin participates in numerous processes as an integrator, with functions in cell adhesion, migration, and differentiation (16-18,23-27). Thus, in the dermis, by the 21<sup>st</sup> day, an increase in the number of Ki-67 positive cells was found in the STT group. Activated fibroblasts through vimentin may coordinate the migratory and proliferative activity of other cells, for example, progenitor cells involved in scar restructuring. This correlates with studies where activated fibroblasts have been shown to facilitate cell migration and tissue reconstruction (3,22,24).

Except for vimentin, synthetic activity is certainly associated with collagens. By the 21<sup>st</sup> day, the structure of the fibrous component approached the norm. It is worth mentioning

that no sclerotic changes were detected, and microcirculation indicators on the 21<sup>st</sup> day significantly corresponded to intact ones (Fig. 4C and D). Fibrosis may occur due to intense collagenases, but this did not occur in the STT group, which was probably due to the accelerated scar restructuring. Thus, it is known that rapid scar maturation and remodeling prevent the formation of hypertrophic or keloid scars.

In addition, possible interactions between fibroblasts and mast cells played a certain role here. Mast cells are known to be involved in scar remodeling processes. Thus, it is reported that mast cells are responsible for fibroblast proliferation inhibition, which prevents keloid scar formation (28). Considering our results, possibly a large number of activated-STT fibroblasts was a kind of trigger for mast cells to increase their functional activity. As a result, a better-quality scar with better mechanical characteristics was formed. As long as about 60% of the skin is collagen, it is primarily responsible for mechanical properties (29). The structure of collagen fibers in the scar differs from normal skin, which leads to functional disorders. The fibers in the scar are less differentiated, sharply thickened, and randomly located. In keloid scars, an abundance of collagen is noted not only in the reticular but also in the papillary layer (30). In our work, it was demonstrated for CONTR and VAS groups, which presented weak mechanical characteristics. However, in the STT group, the mechanical properties of the skin improved. Probably, due to the STT stimulation of fibroblasts, a mature scar with a predominance of the fibrous component was formed. Considering the described trends, it seems relevant in future studies to study key biomarkers of fibroblast to reveal the mechanisms of their activation under STT treatment. The mechanism of STT action can be clarified by gene expression analyses (such as matrix metalloproteinase 9 and matrix remodeling associated 7 genes) or protein quantification (vimentin, collagens, and fibroblast activation protein  $\alpha$ ), which play an important role in fibroblast activation during regeneration (1,31).

## CONCLUSION

The results of the current study revealed the beneficial effect of STT on skin wound regeneration. Morphological studies indicated that STT was responsible for scar formation with a structure similar to normal skin. An increase in the fibroblasts number, high levels of vimentin, and an intensification of collagenogenesis point to the stimulating effect of STT on the functional activity of fibroblasts. However, detailed mechanisms remain to be explored in future studies.

### Acknowledgments

The work was supported by the Ural Federal University named after the first President of Russia B.N. Yeltsin as part of the Enhancement Program Competitiveness, Decree 211 of the Government of the Russian Federation (Contract No. 02.A03.21.0006).

### Conflict of interest statement

The authors declared no conflicts of interest in this study.

### Authors' contributions

I.M. Petrova performed all histological analyses, data acquisition and analysis, statistical analysis, literature search, and wrote the first draft of the manuscript; S.I. Chebanova assisted in microscopic studies; S.L. Khatsko performed all experimental studies with the animals and provided close supervision and helpful discussions about the project direction; T.A. Kalinina synthesized STT and prepared the skin ointment; D.V. Zaitsev assisted with mechanical testing and interpretation of results; and T.V. Glukhareva participated in the concept, design, and coordination of the study and also provided proofreading of the manuscript. All authors read and approved the finalized article.

## REFERENCES

1. Shen Y, Ning J, Zhao L, Liu W, Wang T, Yu J, *et al.* Matrix remodeling associated 7 proteins promote cutaneous wound healing through vimentin in coordinating fibroblast functions. *Inflamm Regen.* 2023;43(5):1-11. DOI: 10.1186/s41232-023-00256-8.
2. Gurtner GC, Werner S, Barrandon Y, Longaker MT. Wound repair and regeneration. *Nature.* 2008;453(7193):314-321. DOI: 10.1038/nature07039.
3. Reinke JM, Sorg H. Wound repair and regeneration. *Eur Surg Res.* 2012;49(1):35-43. DOI: 10.1159/000339613.
4. Cheng F, Shen Y, Mohanasundaram P, Lindstrom M, Ivaska J, Ny T, *et al.* Vimentin coordinates fibroblast proliferation and keratinocyte differentiation in wound healing via TGF-beta-Slug signaling. *Proc Natl Acad Sci USA.* 2016;113(30):4320-4327. DOI: 10.1073/pnas.1519197113.
5. Petrova IM, Zaitsev DV, Zhdanova AV, Khatsko SL, Vysokova OA, Kalinina TA, *et al.* The assessment 1,2,3-triazolo- [5,1-b]-1,3,4-thiadiazine derivative influence on restoration of the fibrous skin component during regeneration caused by burn injury. *Russ J of Biomech.* 2022;26(2):31-41. DOI: 10.15593/RJBiomech/2022.2.03.
6. Otterco AN, Andrade AL, Brassolatti P, Pinto KNZ, Araujo HSS, Parizotto NA. Photobiomodulation mechanisms in the kinetics of the wound healing process in rats. *J Photochem Photobiol B.* 2018;183:22-29. DOI: 10.1016/j.jphotobiol.2018.04.010.
7. Eissa AE, Zaki MM, Saeid S, Abdelsalam M, Ali HM, Moustafa AA, *et al.* *In vitro* evaluation of the efficacy of hemodialysate (Solcoseryl®) as a wound healing agent in *Nile tilapia (Oreochromis niloticus)*. *Int J Vet Sci.* 2013;1(2):57-64. DOI: 10.1016/j.ijvsm.2013.09.003.
8. Yuan J, Hou Q, Chen D, Zhong L, Dai X, Zhu Z, *et al.* Chitosan/LiCl composite scaffolds promote skin regeneration in full-thickness loss. *Sci China Life Sci.* 2020;63(4):552-562. DOI: 10.1007/s11427-018-9389-6.
9. Nozdrin VI, Belousova TA, Iatskovskii AN. Morphological aspects of dermatotrophic action of methyluracil applied epicutaneously. *Morfologiya.* 2002;122(5):74-78. PMID: 12530313.
10. Kalinina TA, Bystrykh OA, Pozdina VA, Glukhareva TV, Ulitko MV, Morzherin YY. Synthesis of spiro derivatives of 1,2,3-triazolo[5,1-b][1,3,4]thiadiazines and biological activity thereof. *Chem Heterocycl.* 2015;51(6):589-592. DOI: 10.1007/s10593-015-1742-1.
11. Vysokova OA, Zhdanova AV, Petrova IM, Medvedeva SY, Kalinina TA, Khatsko S, *et al.* Wound-healing effect of spiroconjugated [1,2,3]triazolo[5,1-b][1,3,4] thiadiazine on a linear skin wound model. *Pharm Chem J.* 2019;53(6):642-645. DOI: 10.1007/s11094-019-02054-4.
12. Kalinina TA, Bystrykh OA, Glukhareva TV, Morzherin YY. Transformation of 1,2,3-thiadiazolyl hydrazones as method for preparation of 1,2,3-triazolo[5,1-b][1,3,4]thiadiazines. *J Heterocycl Chem.* 2015;54(1):137-146. DOI: 10.1002/jhet.2554.



13. Morales-Burgos A, Loosemore MP, Goldberg LH. Postoperative wound care after dermatologic procedures: a comparison of 2 commonly used petrolatum-based ointments. *J Drugs Dermatol*. 2013;12(2):163-164. PMID: 23377388.
14. Fang QQ, Wang XF, Zhao WY, Shi BH, Lou D, Chen CY, *et al*. Development of a chitosan-vaseline gauze dressing with wound-healing properties in murine models. *Am J Trop Med Hyg*. 2020;102(2):468-475. DOI: 10.4269/ajtmh.19-0387.
15. Bai H, Kyu-Cheol N, Wang Z, Cui Y, Liu H, Liu H, *et al*. Regulation of inflammatory microenvironment using a self-healing hydrogel loaded with BM-MSCs for advanced wound healing in rat diabetic foot ulcers. *J Tissue Eng*. 2020;11:1-13. DOI: 10.1177/2041731420947242.
16. Rogel MR, Soni PN, Troken JR, Sitikov A, Trejo HE, Ridge KM. Vimentin is sufficient and required for wound repair and remodeling in alveolar epithelial cells. *FASEB J*. 2011; 25(11): 3873-3883. DOI: 10.1096/fj.10-170795.
17. Bleaken BM, Menko AS, Walker JL. Cells activated for wound repair have the potential to direct collective invasion of an epithelium. *Mol. Biol. Cell*. 2016;27(3):451-465. DOI: 10.1091/mbc.E15-09-0615.
18. Walker JL, Bleaken BM, Romisher AR, Alnwibit AA, Menko AS. In wound repair vimentin mediates the transition of mesenchymal leader cells to a myofibroblast phenotype. *Mol Biol Cell*. 2018;29(13):1555-1570. DOI: 10.1091/mbc.E17-06-0364.
19. Karimi A, Rahmati SM, Navidbakhsh M. Mechanical characterization of the rat and mice skin tissues using histostructural and uniaxial data. *Bioengineered*, 2015;6(3):153-160. DOI: 10.1080/21655979.2015.1036202.
20. Chao CYL, Ng GYF, Cheung KK, Zheng YP, Wang LK, Cheing GLY. *In vivo* and *ex vivo* approaches to studying the biomechanical properties of healing wounds in rat skin. *J Biomech Eng*. 2013;135(10):101009,1-8. DOI: 10.1115/1.4025109.
21. Hudson LG, Newkirk KM, Chandler HL, Choi C, Fossey SL, Parent AE, *et al*. Cutaneous wound reepithelialization is compromised in mice lacking functional Slug (Snai2). *J Dermatol Sci*. 2009; 56(1):19-26. DOI: 10.1016/j.jdermsci.2009.06.009.
22. Boehnke K, Mirancea N, Pavesio A, Fusenig NE, Boukamp P, Stark HJ. Effects of fibroblasts and microenvironment on epidermal regeneration and tissue function in long-term skin equivalents. *Eur J Cell Biol*. 2007;86(11-12):731-346. DOI: 10.1016/j.ejcb.2006.12.005.
23. Ivaska J, Pallari HM, Nevo J, Eriksson JE. Novel functions of vimentin in cell adhesion, migration, and signaling. *Exp Cell Res*. 2007;313(10):2050-2062. DOI: 10.1016/j.yexcr.2007.03.040.
24. Mendez MG, Kojima SI, Goldman RD. Vimentin induces changes in cell shape, motility, and adhesion during the epithelial to mesenchymal transition. *FASEB J*. 2010;24(6):1838-1851. DOI: 10.1096/fj.09-151639.
25. Dave JM, Bayless KJ. Vimentin as an integral regulator of cell adhesion and endothelial sprouting. *Microcirculation*. 2014;21(4):333-344. DOI: 10.1111/micc.12111.
26. Battaglia RA, Delic S, Herrmann H, Snider NT. Vimentin on the move: new developments in cell migration. *F1000Res*. 2018;7:F1000,1-10. DOI: 10.12688/f1000research.15967.1.
27. Meenakshi J, Jayaraman V, Ramakrishnan KM, Babu M. Keloids and hypertrophic scars: a review. *Indian J Plast Surg*. 2005;38(2):175-179. DOI: 10.1097/00006534-198911000-00021.
28. Yang W, Sherman VR, Gludovatz B, Schaible E, Stewart P, Ritchie RO, *et al*. On the tear resistance of skin. *Nat Commun*. 2015;6(1): 1-10. DOI: 0.1038/ncomms7649.
29. Haydont V, Bernard BA, Fortunel NO. Age-related evolutions of the dermis: clinical signs, fibroblast and extracellular matrix dynamics. *Mech Ageing Dev*. 2019;177:150-156. DOI: 10.1016/j.mad.2018.03.006.
30. Stunova A, Vistejnova L. Dermal fibroblasts- a heterogeneous population with regulatory function in wound healing. *Cytokine Growth Factor Rev*. 2018;39:137-150. DOI: 10.1016/j.cytogfr.2018.01.003.
31. Boink MA, Roffel S, Breetveld M, Thon M, Haasjes MSP, Waaijman T, *et al*. Comparison of advanced therapy medicinal product gingiva and skin substitutes and their *in vitro* wound healing potentials. *J Tissue Eng Regen Med*. 2018;12(2):e1088–e1097. DOI: 10.1002/term.2438.