



Article

Acute Stress Response of Sheep to Shearing Procedures: Dynamic Change of Cortisol Concentration and Protein Electrophoretic Pattern

Francesca Arfuso ¹, Francesco Fazio ¹, Lucas Chikhi ², Guillaume Aymond ², Giuseppe Piccione ^{1,*} and Claudia Giannetto ¹

¹ Department of Veterinary Sciences, University of Messina, 98168 Messina, Italy; farfuso@unime.it (F.A.); ffazio@unime.it (F.F.); claudia.giannetto1@unime.it (C.G.)

² École Nationale Vétérinaire de Toulouse, Chemin des Capelles, 31300 Toulouse, France; chikhilucas@gmail.com (L.C.); guillaume.aymond_18@envt.fr (G.A.)

* Correspondence: gpiccione@unime.it; Tel.: +39-090-676-6830

Simple Summary: Farm animals are daily subjected to a wide range of abiotic stressors. Shearing, one practice of routine management procedures, is applied periodically to harvest sheep's wool. During shearing, short-term acute stress is likely to occur. This study aimed to evaluate the change in hematological parameters, serum cortisol concentration and serum protein electrophoretic pattern in Comisana sheep following shearing procedures. The findings obtained in the current study suggest a linkage between the endocrine and immune systems and an acute phase response during the shearing procedure.



Citation: Arfuso, F.; Fazio, F.; Chikhi, L.; Aymond, G.; Piccione, G.; Giannetto, C. Acute Stress Response of Sheep to Shearing Procedures: Dynamic Change of Cortisol Concentration and Protein Electrophoretic Pattern. *Animals* **2022**, *12*, 862. <https://doi.org/10.3390/ani12070862>

Academic Editors: Daniela Alberghina and Silvana Mattiello

Received: 10 February 2022

Accepted: 28 March 2022

Published: 29 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The current study aimed to investigate the influence of acute stress by shearing procedures on hematological parameters, serum cortisol concentration and serum protein electrophoretic pattern in Comisana sheep. A total of 20 not pregnant and not lactating adult ewes, aged 3–4 years old and with a mean bodyweight of 55.50 ± 3.50 kg, were enrolled in the study. From each animal, blood samples were collected before shearing (T_{PRE}) and 5 (T_{POST5}) and 60 (T_{POST60}) minutes after the end of the shearing procedure in order to assess the values of hematological parameters, serum cortisol, total proteins and protein fractions, including albumin, α -, β 1-, β 2- and γ -globulins. According to statistical analysis results, albumin values were lower at T_{POST60} than T_{POST5} ($p < 0.01$), whereas α - and β 2-globulins and the A/G ratio were higher at T_{POST60} with respect to T_{PRE} ($p < 0.01$) and T_{POST5} ($p < 0.01$). A higher serum concentration of cortisol was found at T_{POST5} and T_{POST60} than T_{PRE} ($p < 0.01$), and at T_{POST60} than T_{POST5} ($p < 0.01$). The serum cortisol values were negatively correlated with the serum values of albumin, β 1-globulins and A/G ratio at T_{POST60} , and positively correlated with α - and β 2-globulins at T_{POST5} and T_{POST60} . The decrease in the albumin concentration and the increase in the α - and β 2-globulins observed in ewes after shearing with respect to the baseline values suggests an acute phase response in shorn ewes. Additionally, the correlation found between the serum cortisol concentration and the serum protein fractions confirmed the immunomodulatory effect of this hormone, emphasizing the linkage between the endocrine and immune systems during an acute stress condition.

Keywords: acute stress; sheep; shearing; stress response; serum protein fractions; electrophoresis

1. Introduction

Farm animals are daily subjected to a wide range of abiotic stressors and management practices which can often cause stress to the animal with negative consequences on its health status and welfare [1,2]. Shearing, one practice of routine management procedures in European countries [3], is applied to harvest the sheep's wool to ameliorate the resistance of sheep to high temperatures in summer and for hygienic reasons [4]. The shearing

procedure represents short-term acute stress for animals [2,5] who develop behavioral, autonomic, endocrine and immune responses to maintain homeostasis [2–11]. Stress responses are related not only to the nature and the intensity of the stressful stimulus but also to individual response predispositions or temperament [6,7]. A deep knowledge of stress responses in farm animals allows practitioners to understand the welfare status of the animals and to create appropriate housing and husbandry conditions to improve production efficiency and product quality [8]. Blood is the tissue mostly investigated to assess the animal health status. Specifically, the evaluation of the hematological profile of an animal is considered a useful and sensitive index to screen physiological and pathological modifications [9]. Under acute stress conditions, the endocrine system is stimulated, and, consequently, an increase in corticosteroids occurs [10,11]. The rise of glucocorticoids in the bloodstream also indicates an inflammation state accompanied by a clear increase and/or decrease in a number of plasma proteins known as acute-phase globulins [12]. It is well-known that glucocorticoids play a crucial permissive role in the synthesis of acute-phase globulins. The acute-phase response is part of the first line of defense against injury and/or stressors as it falls within the non-specific immune system [13,14]. The acute-phase serum protein response could be pictured by the serum protein electrophoretic pattern [15]. An acute-phase response encompasses the increase in positive acute-phase proteins and the decrease in negative acute-phase proteins. Thus, different globulin fractions are not the only parameters to be considered in order to evaluate the severity of an acute-phase response [16,17]. The physiological adaptation of sheep to shearing has been extensively investigated in the field of hematochemical response as well as a thermal homeostasis alteration of the sheared animal's thermoregulation [4,18–28]. However, the early stress response of sheep to shearing has been poorly investigated. To the best of the authors' knowledge, no studies on the influence of this common management practice on serum protein fractions are available in the scientific literature so far. In view of the above consideration, the present study aimed to determine the influence of acute stress by shearing procedures on hematological parameters, serum cortisol concentrations and serum protein electrophoretic patterns in Comisana sheep.

2. Materials and Methods

2.1. Animals and Experimental Design

For the study, a flock of about 100 Comisana breed sheep was chosen during the mild-dry season of May–June in Sicily, Italy (40°41' N; 14°26' E, 9 m above sea level). The sheep are shorn once a year, in late spring, according to the traditional techniques of Sicilian sheep breeding. The animals of the chosen flock grazed on natural pasture were penned at night, when they received a concentrated commercial food supplement of 200 g per head (crude protein (CP) 20.4% and 12.5 MJ metabolizable energy (ME)/kg dry matter (DM)). The sheep had free access to water and hay (CP 11.1% and 7.2 MJ ME/kg DM). A total of 20 adult ewes (55.50 ± 3.50 kg, aged 3–4 years old) shorn in the previous year and familiar with the handling required for venipuncture were chosen from the flock enrolled in the study. The enrolled ewes were defined as clinically healthy with no evidence of disease and free from internal and external parasites.

The shearing was performed on 10 June 2021 (ambient temperature 28 ± 2 °C and relative humidity $64 \pm 6\%$); the procedure was performed on all sheep on the same day starting at 7:00 a.m., and it took about 1 h and 50 min to complete the procedure on all ewes. Shearing was executed by hand using traditional shearing scissors in a 15 m × 10 m pen, and it lasted approximately 5 min for each sheep. At the end of the shearing procedure, the animals were released into an adjacent enclosure after being marked by a progressive number through an appropriate colored spray to identify them.

2.2. Sampling Procedures and Laboratory Analysis

From each sheep, blood was drawn by jugular venipuncture (needle gauge of 20G) and collected into one vacutainer tube containing EDTA and two tubes with clot activator (Terumo Corporation, Tokyo, Japan) before shearing (T_{PRE}) and 5 (T_{POST5}) and 60 (T_{POST60}) min after the end of shearing. The time taken to collect blood from each animal was 40 s, whereas the time between blood sampling of different animals was approximately 10 min. On EDTA whole blood samples, the values of white blood cells (WBCs), red blood cells (RBCs), hematocrit (Hct), hemoglobin (Hb), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration, mean corpuscular volume (MCV) and platelets (PLTs) were assessed by means of an automated hematology analyzer (HeCo Vet C; SEAC, Florence, Italy). Samples from the first tube with a clot activator were allowed to clot for 20 min at room temperature and, thereafter, were centrifuged at $2325 \times g$ for 15 min. By visual inspection, all the obtained sera were non-hemolysed, and they were analyzed to assess the total protein concentration through a commercially available kit (Biosystems S.A., Barcelona, Spain; biuret method using bovine album as a protein standard at a concentration of 6.02 g/dL) by means of an automated ultraviolet (UV) spectrophotometer (Slim; SEAC, Florence, Italy). In order to evaluate serum protein fractions, an electrophoresis analysis was performed using an automated system (Selvet24, Seleco Engineering, Naples, Italy) according to the procedures described by the manufacturer. The protein fractions were divided into albumin, α -, β 1-, β 2-, and γ -globulins according to previous findings on ovine species [29,30]. Relative protein concentrations within each fraction were determined as the optical absorbance percentage; then, the absolute concentration (g/dL) and albumin/globulin ratio (A/G) were calculated using the total protein concentration. Samples from the second tube with a clot activator were allowed to clot overnight at 4 °C before centrifuging at $1000 \times g$ for 20 min at 2–8 °C. The obtained sera concentration of cortisol was evaluated using an ELISA kit specific for ovine species (Bovine/Sheep Cortisol ELISA kit, Elabscience Biotechnology Inc. Kampenhout, Belgium) through a microwell plate reader (Sirio, SEAC, Florence, Italy). Calibrators and samples were run in duplicate, and samples exhibited parallel displacement to the standard curve for both ELISA analyses. Both the intra- and the inter-assay coefficients of variation were <10%.

2.3. Statistical Analysis

All data were normally distributed (Kolmogorov–Smirnov test, $p > 0.05$), and one-way analysis of variance (ANOVA) for repeated measures was used to assess significant changes in hematological parameters, serum cortisol concentration and serum protein profile in ewes at the three time points herein considered (T_{PRE} , T_{POST5} and T_{POST60}). Bonferroni's post hoc comparison was applied to the obtained significances. Pearson's correlation analysis was applied to assess whether the serum cortisol concentration correlated with the values of serum total proteins and their fractions measured in ewes before shearing and 5 and 60 min after the end of shearing procedures. The degree of correlation between these parameters was investigated by means of a linear regression model ($y = a + bx$). p -values < 0.05 were considered statistically significant. The software Prism v. 9.00 (Graphpad Software Ltd., San Diego, CA, USA, 2020) was used to perform statistical analysis.

3. Results

The results are shown as mean values \pm standard deviation (SD).

Statistical analysis of data showed no significant change ($p > 0.05$) in hematological parameters investigated in ewes following the shearing procedure (Table 1). The representative protein electrophoretograms obtained from serum samples of Comisana ewes before and after 5 and 60 min from the end of the shearing procedure are displayed in Figure 1. As reported in Table 2, albumin values were statistically significantly lower at T_{POST60} than T_{POST5} ($p < 0.01$); additionally, α - and β 2-globulins and the A/G ratio were higher at T_{POST60} with respect to T_{PRE} ($p < 0.01$) and T_{POST5} ($p < 0.01$). No statistically significant changes were found in the serum concentration of total proteins ($p > 0.05$), β 1- ($p > 0.05$) and

γ -globulins ($p > 0.05$) throughout the experimental period. Higher serum concentrations of cortisol were found at T_{POST5} ($p < 0.01$) and T_{POST60} ($p < 0.01$) than T_{PRE} , and at T_{POST60} ($p < 0.01$) than T_{POST5} (Table 2). No significant correlation was found between the serum cortisol concentration and the hematological parameters measured in investigated ewes before (T_{PRE}) ($p > 0.05$) and after the shearing procedure (T_{POST5} and T_{POST60}) ($p > 0.05$) (Table 3). The serum cortisol values were not correlated with the serum concentration of total proteins and γ -globulins measured in investigated ewes at T_{PRE} ($p > 0.05$), T_{POST5} ($p > 0.05$) and T_{POST60} ($p > 0.05$), whereas the concentration of this hormone was negatively correlated with the serum values of albumin ($p < 0.05$), β 1-globulins ($p < 0.05$) and the A/G ratio ($p < 0.05$) at T_{POST60} , and positively correlated with α - ($p < 0.05$) and β 2-globulins ($p < 0.05$) at T_{POST5} and T_{POST60} . The linear regression model confirmed the statistically significant correlations found among selected parameters (Figures 2–4).

Table 1. White blood cells (WBCs), red blood cells (RBCs), hematocrit (Hct), hemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) and platelets (PLTs) measured in ewes before shearing (T_{PRE}) and after 5 and 60 min (T_{POST5} and T_{POST60}) from the end of shearing. Data are expressed as mean values \pm standard deviation (SD).

Hematological Parameters	Experimental Period		
	T_{PRE}	T_{POST5}	T_{POST60}
WBCs ($\times 10^3/\mu\text{L}$)	7.82 \pm 1.58	7.48 \pm 1.45	7.83 \pm 1.61
RBCs ($\times 10^6/\mu\text{L}$)	10.71 \pm 0.78	10.50 \pm 0.62	10.47 \pm 0.81
Hct (%)	36.55 \pm 4.72	35.81 \pm 4.92	34.41 \pm 3.98
Hb (g/dL)	9.78 \pm 0.91	9.68 \pm 1.05	9.59 \pm 0.97
MCV (fL)	35.60 \pm 5.14	35.62 \pm 5.16	34.50 \pm 5.20
MCH (pg)	9.51 \pm 0.87	9.60 \pm 0.90	9.57 \pm 0.86
MCHC (%)	26.94 \pm 0.40	27.19 \pm 0.83	28.03 \pm 1.78
PLTs ($\times 10^3/\mu\text{L}$)	604.85 \pm 168.48	636.08 \pm 119.30	575.45 \pm 101.08

Table 2. Serum cortisol, total proteins, albumin, α -, β 1-, β 2-, γ -globulins and the A/G ratio were measured in ewes before shearing (T_{PRE}) and after 5 and 60 min (T_{POST5} and T_{POST60}) from the end of the shearing procedure, together with statistically significant changes. Data are expressed as mean values \pm standard deviation (SD).

Serum Parameters	Experimental Period		
	T_{PRE}	T_{POST5}	T_{POST60}
Cortisol (ng/mL)	94.99 \pm 4.14	100.3 \pm 1.93 ^a	106.5 \pm 2.98 ^{a,b}
Total proteins (d/dL)	7.05 \pm 0.80	7.65 \pm 0.78	7.10 \pm 0.68
Albumin (g/dL)	3.30 \pm 0.28	3.71 \pm 0.68	2.93 \pm 0.38 ^b
α -globulins (g/dL)	0.26 \pm 0.06	0.28 \pm 0.05	0.40 \pm 0.08 ^{a,b}
β 1-globulins (g/dL)	0.90 \pm 0.11	0.93 \pm 0.10	1.03 \pm 0.14
β 2-globulins (g/dL)	0.58 \pm 0.06	0.68 \pm 0.08	0.88 \pm 0.10 ^{a,b}
γ -globulins (g/dL)	2.01 \pm 0.33	2.05 \pm 0.31	1.85 \pm 0.13
A/G ratio	0.89 \pm 0.10	0.96 \pm 0.23	0.71 \pm 0.11 ^b

Note: statistical significances: ^a versus T_{PRE} ($p < 0.01$); ^b versus T_{POST5} ($p < 0.01$).




Figure 1. Representative cellulose acetate electrophoretograms of serum proteins were obtained from ewes before shearing (T_{PRE}) and after 5 (T_{POST5}) and 60 (T_{POST60}) min from the end of the shearing procedure.

Table 3. Coefficients of correlation (Pearson's r and p -value) among the serum concentration of cortisol, the levels of white blood cells (WBCs), red blood cells (RBCs), hematocrit (Hct), hemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) and platelets (PLTs) measured in ewes before shearing (T_{PRE}) and after 5 and 60 min (T_{POST5} and T_{POST60}) from the end of shearing. p -values < 0.05 were considered statistically significant.

		WBCs ($\times 10^3/\mu\text{L}$)	RBCs ($\times 10^6/\mu\text{L}$)	Hct (%)	Hb (g/dL)	MCV (fL)	MCH (pg)	MCHC (%)	PLTs ($\times 10^3/\mu\text{L}$)
Cortisol (ng/mL)	T_{PRE}	$r = 0.24$ $p = 0.31$	$r = -0.99$ $p = 0.70$	$r = -0.02$ $p = 0.95$	$r = -0.20$ $p = 0.40$	$r = -0.10$ $p = 0.67$	$r = -0.30$ $p = 0.20$	$r = -0.13$ $p = 0.59$	$r = -0.20$ $p = 0.40$
	T_{POST5}	$r = -0.44$ $p = 0.06$	$r = -0.37$ $p = 0.11$	$r = 0.18$ $p = 0.46$	$r = 0.10$ $p = 0.68$	$r = 0.31$ $p = 0.18$	$r = 0.43$ $p = 0.06$	$r = 0.12$ $p = 0.61$	$r = -0.29$ $p = 0.21$
	T_{POST60}	$r = -0.24$ $p = 0.30$	$r = 0.27$ $p = 0.26$	$r = -0.44$ $p = 0.06$	$r = -0.15$ $p = 0.54$	$r = -0.16$ $p = 0.50$	$r = 0.13$ $p = 0.61$	$r = 0.06$ $p = 0.81$	$r = -0.26$ $p = 0.27$




Figure 2. Linear regression model results found between the serum values of cortisol and the serum concentration of serum total proteins, albumin, α -, β 1-, β 2-, γ -globulins and the A/G ratio measured in ewes before the shearing procedure (T_{PRE}).




Figure 3. Linear regression model results found between the serum values of cortisol and the serum levels of total proteins, albumin, α -, β 1-, β 2-, γ -globulins and the A/G ratio measured in ewes after 5 min from the end of the shearing procedure (T_{POST5}).




Figure 4. Linear regression model results found between the serum values of cortisol and the serum levels of serum total proteins, albumin, α -, β 1-, β 2-, γ -globulins and the A/G ratio measured in ewes after 60 min from the end of the shearing procedure (T_{POST60}).

4. Discussion

Shearing includes several procedures, such as animal capture, separation, immobilization and tying, which are a potential source of acute stress.

It has been well established that acute stressors could lead to a transient increase in cells in the bloodstream as a consequence of splenic contraction following catecholamine stimulation as the so-called defence alarm reaction [31]. Under stress conditions, there is also a cortisol-mediated release of cells from the bone marrow [31]. According to the results obtained in the current study, hematological parameters did not show differences related to the shearing procedure in ewes. The missing change in hematological indices in stressed ewes could be due to the timing of blood sampling. In the current study, blood was collected before shearing and 5 and 60 min after the end of the stressor experience; hence, the transient increase of cells into the bloodstream could have been missed.

Contrariwise, it was found that shearing influenced some serum protein fractions measured in investigated ewes. The changes in total serum protein concentrations, as well as their fractional distribution and albumin/globulin ratio, are usually the first signs of homeostasis perturbation and/or disease conditions [32]. According to the serum protein electrophoretic pattern previously suggested for ovine species [29], five protein fractions were observed in sera obtained from ewes enrolled in the current study.

The analysis of protein electrophoretograms obtained from serum samples of Comisana ewes showed that a decrease in the albumin concentration and an increase in the α - and β 2-globulins were found in ewes after shearing. These findings suggest an acute phase response onset as an integral part of an acute stress response in ewes. The pro-inflammatory mediators, including cytokines, are the main factors involved in the regulation network of an acute-phase response [12] which stimulates the liver to synthesize some acute-phase proteins (APPs) whose blood levels increase. Thus, they are known as positive APPs. Other proteins, known as negative APPs, display decreasing trends in response to challenges [12]. The α - and β 2-globulins include the main positive APPs (i.e., alpha 1-acid glycoprotein, caeruloplasmin, complement proteins, C-reactive protein, fibrinogen, haptoglobin, serum amyloid A). The β 1-globulins fraction includes transferrin, which decreases in response to challenges like albumin. It is well known that the cortisol concentration rises in sheep subjected to handling and shearing as a stress response [22,27]. In agreement, the serum cortisol concentrations increased in ewes after shearing compared to pre-shearing, highlighting that this procedure would be stressful due to different factors linked to shearing, including handling of animals and isolation [22,33]. It is noteworthy that, in the current study, α - and β 2-globulins were positively correlated with serum cortisol levels at T_{POST5} and T_{POST60} , whereas albumin and β 1-globulins showed a negative correlation with the serum concentration of this hormone at T_{POST60} . Cortisol is known to have immunomodulatory effects and, together with catecholamines, affects the proliferation of B and T cells, cytokine production, and antibody production upon acute stress conditions [6,7].

Studies carried out on rats suggested that glucocorticoids and catecholamine could trigger the synthesis of proteins involved in the acute-phase response [34,35]. In agreement with this, the results obtained in the current study lead us to hypothesize that cortisol could be one of the mediators of mechanisms orchestrating the fine and complex acute phase response of the animal under stress circumstances. Additionally, the findings seem to reinforce the well-established idea that, upon a stressful condition, the immune and endocrine systems cross-talk to re-establish homeostasis of the organism [6].

5. Conclusions

According to the results herein gained, the acute stress caused by the shearing procedure induced a dynamic change in the electrophoretic serum protein pattern and the cortisol concentration. The electropherograms obtained following shearing suggest an acute-phase response as highlighted by the increase of α - and β -globulins, which are known to include many acute-phase proteins. Additionally, the significant relationship found between the serum concentration of cortisol, the albumin values, and α -, β 1-, and

β 2-globulins confirmed the immunomodulatory effect of this hormone and emphasized the linkage between the endocrine and immune systems during acute stress.

Author Contributions: Conceptualization, F.A.; methodology, F.A., C.G., L.C., G.A. and F.F.; software, F.A.; validation, F.A., C.G. and G.P.; formal analysis, F.A.; investigation, F.A.; resources, C.G.; data curation, F.A., C.G. and F.F.; writing—original draft preparation, F.A.; writing—review and editing, F.A. and C.G.; visualization, G.P.; supervision, C.G. and G.P.; project administration, G.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: All treatments, housing, and animal care were carried out in accordance with the standards recommended by the European Directive 2010/63/EU for animal experiments.

Informed Consent Statement: A written Informed Consent is obtained from the owner of the animals.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Le Neindre, P.; Boivin, X.; Boissy, A. Handling of extensively kept animals. *Appl. Anim. Behav. Sci.* **1996**, *49*, 73–81. [[CrossRef](#)]
2. Yardimci, M.; Sahin, E.H.; Cetingul, I.S.; Bayram, I.; Aslan, R.; Sengor, E. Stress responses to comparative handling procedures in sheep. *Animal* **2013**, *7*, 143–150. [[CrossRef](#)] [[PubMed](#)]
3. Dýrmundsson, O.R. Shearing time of sheep with special reference to conditions in northern Europe: A review. *Bùvisindi Icel. Agric. Sci.* **1991**, *5*, 39–46.
4. Piccione, G.; Casella, S.; Alberghina, D.; Zumbo, A.; Pennisi, P. Impact of shearing on body weight and serum total proteins in ewes. *Span. J. Agric. Res.* **2010**, *8*, 342–346. [[CrossRef](#)]
5. Sanger, M.E.; Doyle, R.E.; Hinch, G.N.; Lee, C. Sheep exhibit a positive judgement bias and stress-induced hyperthermia following shearing. *Appl. Anim. Behav. Sci.* **2011**, *131*, 94–103. [[CrossRef](#)]
6. Etim, N.N.; Evans, E.I.; Offiong, E.E.A.; Williams, M.E. Stress and the neuroendocrine system: Implications for animal well-being. *Am. J. Biol. Life Sci.* **2013**, *1*, 20–26.
7. Kumar, B.; Manuja, A.; Aich, P. Stress and its impact on farm animals. *Front. Biosci. Elite* **2012**, *4*, 1759–1767. [[CrossRef](#)]
8. Blokhuis, H.J.; Hopster, H.; Geverink, N.A.; Korte, S.M.; van Reenen, C.G. Studies of stress in farm animals. *Comp. Haematol. Int.* **1998**, *8*, 94–101. [[CrossRef](#)]
9. Etim, N.N.; Williams, M.E.; Akpabio, U.; Offiong, E.E.A. Haematological Parameters and Factors Affecting Their Values. *Agric. Sci.* **2014**, *2*, 37–47. [[CrossRef](#)]
10. Wingfield, J.C.; Kitaysky, A.S. Endocrine responses to unpredictable environmental events: Stress or anti-stress hormones? *Integr. Comp. Biol.* **2002**, *42*, 600–609. [[CrossRef](#)]
11. Lay, D.C.; Friend, T.H.; Bowers, C.L.; Grissom, K.K.; Jenkins, O.C. A comparative physiological and behavioral study of freeze and hot-iron branding using dairy cows. *J. Anim. Sci.* **1992**, *70*, 1121–1130. [[CrossRef](#)] [[PubMed](#)]
12. O'Connell, T.X.; Horita, T.J.; Kasravi, B. Understanding and interpreting serum protein electrophoresis. *Am. Fam. Physician* **2005**, *71*, 105–112. [[PubMed](#)]
13. Reczyńska, D.; Zalewska, M.; Czopowicz, M.; Kaba, J.; Zwierzchowski, L.; Bagnicka, E. Acute phase protein levels as an auxiliary tool in diagnosing viral diseases in ruminants—A Review. *Viruses* **2018**, *10*, 502. [[CrossRef](#)]
14. Greguła-Kania, M.; Kosior-Korzecka, U.; Patkowski, K.; Juszcuk-Kubiak, E.; Plewik, M.; Gruszecki, T.M. Acute-phase proteins, cortisol and haematological parameters in ewes during the periparturient period. *Reprod. Domest. Anim.* **2020**, *55*, 393–400. [[CrossRef](#)] [[PubMed](#)]
15. Azim, W.; Azim, S.; Ahmed, K.; Shafi, H.; Rafi, T.; Luqman, M. Diagnostic significance of serum protein electrophoresis. *Biomedica* **2004**, *20*, 40–44.
16. Ceciliani, F.; Ceron, J.J.; Eckersall, P.D.; Sauerwein, H. Acute phase proteins in ruminants. *J. Proteom.* **2012**, *75*, 4207–4231. [[CrossRef](#)] [[PubMed](#)]
17. Bertoni, G.; Trevisi, E.; Han, X.; Bionaz, M. Effects of inflammatory conditions on liver activity in puerperium period and consequences for performance in dairy cows. *J. Dairy Sci.* **2008**, *91*, 3300–3310. [[CrossRef](#)]
18. Piccione, G.; Caola, G.; Refinetti, R. Effect of shearing on the core temperature of three breeds of Mediterranean sheep. *Small Rum. Res.* **2002**, *46*, 211–215. [[CrossRef](#)]
19. Piccione, G.; Caola, G. Influence of shearing on the circadian rhythm of body temperature in the sheep. *J. Vet. Med. A* **2003**, *50*, 235–240. [[CrossRef](#)]

20. Piccione, G.; Casella, S.; Fazio, F.; Pennisi, P. Effect of shearing on some haematochemical parameters in ewes. *Czech J. Anim. Sci.* **2008**, *53*, 106–111. [[CrossRef](#)]
21. Piccione, G.; Fazio, F.; Casella, S.; Pennisi, P.; Caola, G. Influence of shearing on oxidative stress and some physiological parameters in ewes. *Anim. Sci. J.* **2011**, *82*, 481–485. [[CrossRef](#)] [[PubMed](#)]
22. Hargreaves, A.L.; Huston, G.D. The stress response in sheep during routine handling procedures. *Appl. Anim. Behav. Sci.* **1990**, *26*, 83–90. [[CrossRef](#)]
23. Pennisi, P.; Costa, A.; Biondi, L.; Avondo, M.; Piccione, G. Influence of the fleece on thermal homeostasis and on body condition in Comisana ewe lambs. *Anim. Res.* **2004**, *53*, 13–19. [[CrossRef](#)]
24. Aleksiev, Y. The effect of shearing on the behaviour of some physiological responses in lactating Plevan blackhead ewes. *Bulg. J. Agric. Sci.* **2009**, *15*, 446–452.
25. Beatty, D.T.; Barnes, A.; Fleming, P.A.; Taylor, E.; Maloney, S.K. The effect of fleece on core and rumen temperature in sheep. *J. Therm. Biol.* **2008**, *33*, 437–443. [[CrossRef](#)]
26. Casella, S.; Giudice, E.; Passantino, A.; Zumbo, A.; Di Pietro, S.; Piccione, G. Shearing induces secondary biomarkers responses of thermal stress in sheep. *Anim. Sci. Pap. Rep.* **2016**, *34*, 73–80.
27. Hefnawy, A.; Helal, M.; Sabek, A.; Shousha, S. Clinical, behavioral and biochemical alterations due to shearing stress in Ossimi sheep. *J. Vet. Med. Sci.* **2018**, *80*, 1281–1286. [[CrossRef](#)]
28. Piccione, G.; Lutri, L.; Casella, S.; Ferrantelli, V.; Pennisi, P. Effect of shearing and environmental conditions on physiological mechanisms in ewes. *J. Environ. Biol.* **2008**, *29*, 877–880.
29. Kaneko, J.J.; Harvey, J.W.; Bruss, M.L. *Clinical Biochemistry of Domestic Animals*, 5th ed.; Academic Press Inc.: San Diego, CA, USA, 1997.
30. Nagyová, V.; Arfuso, F.; Rizzo, M.; Nagy, O.; Piccione, G. Stability of ovine serum protein fractions under different storage conditions. *J. Vet. Diagn. Investig.* **2017**, *29*, 312–315. [[CrossRef](#)]
31. Jern, C.; Wadenvik, H.; Mark, H.; Hallgren, J.; Jern, S. Haematological changes during acute mental stress. *Br. J. Haematol.* **1989**, *71*, 153–156. [[CrossRef](#)]
32. Dede, S.; Altug, N.; Deger, Y.; Ozdal, N.; Ceylan, E. Serum biochemical profile and protein fractions in cattle with Theileriosis. *Rev. Méd. Vét.* **2014**, *165*, 137–143.
33. Carcangiu, V.; Vacca, G.M.; Parmeggiani, A.; Mura, M.C.; Pazzola, M.; Dettori, M.L.; Bini, P.P. The effect of shearing procedures on blood levels of growth hormone, cortisol and other stress haematochemical parameters in Sarda sheep. *Animal* **2008**, *2*, 606–612. [[CrossRef](#)]
34. Thompson, W.L.; Abeles, F.B.; Beall, F.A.; Dinterman, R.E.; Wannemacher, R.W., Jr. Influence of the adrenal glucocorticoids on the stimulation of synthesis of hepatic ribonucleic acid and plasma acute-phase globulins by leucocytic endogenous mediator. *Biochem. J.* **1976**, *156*, 25–32. [[CrossRef](#)] [[PubMed](#)]
35. Van Gool, J.; Boers, W.; Sala, M.; Ladiges, N.C. Glucocorticoids and catecholamines as mediators of acute-phase proteins, especially rat alpha-macrofoetoprotein. *Biochem. J.* **1984**, *220*, 125–132. [[CrossRef](#)] [[PubMed](#)]