

EDGE ARTICLE

Cite this: *Chem. Sci.*, 2022, 13, 8871

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 6th May 2022

Accepted 4th July 2022

DOI: 10.1039/d2sc02541h

rsc.li/chemical-science

Asymmetric synthesis of chromanone lactones via vinylogous conjugate addition of butenolide to 2-ester chromones†

Yi Li, Shuang Xin, Rui Weng, Xiaohua Liu * and Xiaoming Feng *

Chiral chromanone lactones are a class of natural products with important biological activity. We report a direct diastereo- and enantioselective vinylogous conjugate addition of butenolide to 2-ester substituted chromones. The transformation proceeded well in the presence of as low as 1 mol% of a chiral *N,N'*-dioxide/Sc^{III} complex, 3 Å MS and a catalytic amount of hexafluoroisopropanol (HFIP). The scope of Michael acceptors includes a variety of substituted chromones at different positions, and the desired chromanone lactones upon reduction are afforded in good yield and diastereoselectivity, and excellent enantioselectivity (up to 99% ee). The strategy could be used in the concise synthesis of blennolide C and gonytolide A, C and G.

Introduction

Chromanone lactones represent a family of natural products with important biological activity, and the structure characterized as 4-chromanone bearing a five-membered lactone at the 2-position could also give access to tetrahydroxanthone natural products (Scheme 1a).¹ In view of their intriguing bioactivities and structural diversity, the total synthesis of the chromanone lactone series has drawn active interest.² The vinylogous conjugate addition of butanolide or its synthetic equivalents to chromones provided a straightforward route for this purpose. Porco and coworkers initiated a benzopyrylium species involving Mukaiyama addition with silyoxyfurans from 5-hydroxyl-2-carboxylate chromones activated by diisopropylsilyl ditriflate (Scheme 1b),³ and the transformations following the resolution enabled the asymmetric total synthesis of several mono- or dimeric natural products.⁴ The strategy was expanded to the vinylogous addition of silyoxyfurans to chromones promoted with TMSI by Zhang's and Li's groups, which was used for the synthesis of racemic natural products.⁵ A stereo-divergent variety using a chiral copper(i)-complex catalyst and silyoxyfurans was realized by Shibasaki very recently.⁶

On the other hand, the asymmetric catalytic preparation of butanolide-containing chromanones from butanolides seems to be more promising and challenging. Previously, Trost's^{7a} and Shibasaki's groups^{7b} reported asymmetric vinylogous addition


of 2-chromones with unactivated 5-alkyl γ -butenolides as the pronucleophiles using Zn-ProPhenon and Cu-Biphep complex catalysts, respectively (Scheme 1c), providing an atom-economical method without preactivation of the reactants and desilylation.⁸ However, as evidenced by the structures presented in Scheme 1a, 2-ester and 5-hydroxyl substitutions are popular in these natural products, but the employment of 2-ester chromones as Michael acceptors overcomes the challenges of steric hindrance for the construction of a quaternary carbon center,⁹ the retro-oxo-Michael reaction and epimerization of the initial adducts for the achievement of high diastereo- and enantioselectivity.^{3,4} Our research group has realized vinylogous additions of both β,γ -unsaturated butenolides and their equivalents catalyzed by the chiral *N,N'*-dioxide/Sc^{III} complex catalysts.¹⁰ In light of the performance of chiral Lewis acid catalysis using Feng *N,N'*-dioxides as ligands, we reasoned that it could be applied for the preparation of optically active chromanone lactones and tetrahydroxanthones (Scheme 1d). In fact, the direct vinylogous addition of butanolide to 2-ester chromones worked well promoted by as low as 1 mol% of the chiral scandium catalyst of *N,N'*-dioxide. The diastereoselectivity in the Mukaiyama addition of 2-ester chromones from Porco's study showed that the *syn*-diastereomer of the silylene adduct is kinetically favorable. Differently, the highly *anti*-selective and enantioselective chromanone lactone is more readily available in the presence of molecular sieves and alcohols in this study. Herein, we disclosed a highly diastereo- and enantioselective vinylogous conjugate addition for the synthesis of chromanone lactones, which could be used in the concise synthesis of (+)-blennolide C, (+)-gonytolide A and other natural product members.

Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China. E-mail: liuxh@scu.edu.cn; xmxfeng@scu.edu.cn


† Electronic supplementary information (ESI) available. CCDC [1957788]. For ESI and crystallographic data in CIF or other electronic format see <https://doi.org/10.1039/d2sc02541h>




a) Selected chromanone lactones and tetrahydroxanthone natural products




b) Vinylogous addition of siloxyfurans to 2-ester-chromones (by Porco)



c) Asymmetric vinylogous addition of butenolides to chromones (by Trost, Shibasaki)



d) Asymmetric vinylogous addition of butenolides to 2-ester-chromones



Scheme 1 Background information about chromanone lactones and related compounds.

Results and discussion

In the initial studies, the vinylogous addition reaction of 2-ester chromone **1a** and 5-*H*-butenolide **2a** was selected as the model reaction to optimize the reaction conditions (Table 1). The direct addition product **3aa** was unstable during the purification with silica gel, and thus the target lactone-substituted chromanone **3a** was isolated as the final product *via* the treatment with $\text{NaBH}_4/\text{NiCl}_2\cdot 6\text{H}_2\text{O}$ in one-pot after the catalytic reaction.^{3,4} The reaction in the presence of a chiral N,N' -dioxide $\text{L}_3\text{-PrPr}_2/\text{Sc}(\text{OTf})_3$ complex could deliver the desired product **3a** with good enantioselectivity after 4 hours in THF at 30 °C (94% ee; entry 1), and although the yield increased gradually with a prolonged reaction time, both the diastereoselectivity and the ee value decreased during the course (entries 2 and 3), indicating that the epimerization of the adduct **3aa** occurred during the process, similar to the observation from Porco's work.³ To minimize the loss of stereoselectivity we further investigated different additives (see the ESI† for details). It was a delight to find that the medium yield (56% yield) with up to 99% ee and 19 : 1 *anti* : *syn* was obtained after 16 hours when 3 Å MS was added to the reaction system (entry 4). Remarkably, the addition

Table 1 Optimization of the reaction conditions^a

(1) $\text{L}^*/\text{Sc}(\text{OTf})_3$ (1:1, 10 mol%)
additive, THF, 30 °C

(2)

$\text{L}_3\text{-PrPr}_2$: X = *i*Pr
 $\text{L}_3\text{-PrEt}_2$: X = Et
 $\text{L}_3\text{-PrMe}_2$: X = Me


$\text{L}_3\text{-RaPr}_2$
 $\text{L}_3\text{-PiPr}_2$

Entry	Conditions	Yield (%)	dr	ee (%)
1	$\text{L}_3\text{-PrPr}_2$, 4 h	19	4 : 1	94
2	$\text{L}_3\text{-PrPr}_2$, 8 h	35	3.6 : 1	90
3	$\text{L}_3\text{-PrPr}_2$, 24 h	42	2.3 : 1	74
4	$\text{L}_3\text{-PrPr}_2$, 3 Å MS, 16 h	56	>19 : 1	>99
5	$\text{L}_3\text{-PrPr}_2$, 3 Å MS, HFIP, 16 h	82	>19 : 1	>99
6	$\text{L}_3\text{-RaPr}_2$, 3 Å MS, HFIP, 16 h	23	1 : 1	>99
7	$\text{L}_3\text{-PiPr}_2$, 3 Å MS, HFIP, 16 h	42	6 : 1	>99
8	$\text{L}_3\text{-PrMe}_2$, 3 Å MS, HFIP, 16 h	70	9 : 1	78
9	$\text{L}_3\text{-PrEt}_2$, 3 Å MS, HFIP, 16 h	79	12 : 1	>99
10 ^b	$\text{L}_3\text{-PrEt}_2$, 3 Å MS, HFIP, 16 h	83	19 : 1	>99
11 ^{b,c}	$\text{L}_3\text{-PrEt}_2$, 3 Å MS, HFIP, 16 h	82	19 : 1	>99

^a Unless otherwise noted, all reactions were carried out with (1) $\text{Sc}(\text{OTf})_3/\text{L}$ (1 : 1, 10 mol%), **1a** (0.1 mmol), and **2a** (3.0 equiv.) in THF (0.1 M) at 30 °C; 3 Å MS (0 or 20 mg); hexafluoroisopropanol (HFIP, 0 or 20 mol%). (2) $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$ (1.0 equiv.), NaBH_4 (2.0 equiv.), THF/MeOH (3/1, 0.25 M), 0 °C, 10 min. The isolated yield of the product **3a** is provided. The ee values were determined by UPC² analysis and dr values were determined by ¹H NMR. ^b At 20 °C. ^c 1 mol% catalyst loading.

of 20 mol% of hexafluoroisopropanol (HFIP) increased the yield of **3a** to 82% (entry 5). With the established grounded system, we evaluated the performance of different N,N' -dioxide ligands, and the selected results are listed in entries 6–9. It was found that the *L*-proline-derived $\text{L}_3\text{-PrPr}_2$ gave better yield and diastereoselectivity than *L*-ramipril-derived $\text{L}_3\text{-RaPr}_2$ (entry 6), and higher outcome in terms of yield, and diastereo- and enantioselectivity than *L*-pipecolic acid-derived $\text{L}_3\text{-PiPr}_2$ (entry 7). Changing the ligand to 2,6-dimethylaniline-based $\text{L}_3\text{-PrMe}_2$ led to a dramatic drop in stereoselectivity (entry 8). Although the use of 2,6-diethyl aniline-based $\text{L}_3\text{-PrEt}_2$ resulted in decreased diastereoselectivity at 30 °C (entry 9), excellent results could be achieved by lowering the temperature to 20 °C (entry 10). Noteworthy, the loading of the catalyst $\text{L}_3\text{-PrEt}_2/\text{Sc}(\text{OTf})_3$ could be reduced to 1 mol%, and the lactone **3a** was isolated in 82% yield, 99% ee and a 19 : 1 *anti* : *syn* ratio (entry 11).

With the optimized reaction conditions in hand (Table 1, entry 11), the scope of chromones in the vinylogous addition reaction was evaluated (Table 2). Particularly, various substitutions at different positions of chromones have nearly no influence on the enantioselectivity, and all the tested reactions (22

Table 2 The substrate scope^{abcd}


^a Unless otherwise noted, the reactions were carried out with **1** (0.1 mmol), **2a** (0.3 mmol), and $L_3\text{-PrEt}_2/\text{Sc}(\text{OTf})_3$ (1 : 1, 1 mol%) in THF (0.1 M) at 20 °C for 16 hours and then with $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (1.0 equiv.) and NaBH_4 (2.0 equiv.) in THF/MeOH (3/1, 0.25 M) at 0 °C for 10 min. The isolated yield of **3** is provided. The ee values were determined by UPC³ and the dr values were determined by ¹H NMR. ^b $L_3\text{-PrEt}_2/\text{Sc}(\text{OTf})_3$ (5 mol%). ^c At 50 °C. ^d DCE/MeCN (3 : 7, 0.1 M) as the solvent.

examples) yielded the desired products with 96% to >99% ee. Generally, electron-withdrawing or donating substitution at 7- or 8-positions did not significantly affect the yield and diastereoselectivity (**3g–3k**; >19 : 1 dr), while 5- or 6-substituted ones led to varied reactivity and diastereoselectivity. A 5-MeO substituent did not impact the selectivity but reduced the yield (**3b**), and the ethyl ester substitution gave similar results (**3b'**), while a 5-Cl substituent resulted in a reduced dr value (**3c**, 9 : 1 dr). Moderate to good yield and excellent enantioselectivity were observed with 6-methoxychromone (**3d**), 6-fluorochromone (**3e**), or 6-bromochromone (**3f**), but the diastereoselectivity was also obviously affected. Chromones with disubstitution at the 6- and 8-positions were also good electrophiles, and the adducts **3l** and **3m** were afforded in a moderate yield and dr value, with 99% and 96% ee, respectively in the presence of 5 mol% catalyst. With a fused-ring substrate, chromanone lactone **3n** was formed in 72% yield, 12 : 1 dr, and 99% ee.

Owing to the fact that chromanone lactones and the related tetrahydroxanthone natural products contain a hydroxyl group at the 5-position, we then focused on the reactions of 5-MeO-chromones. The disubstituted and tri-substituted chromones were tolerable with excellent enantioselectivity (98–99% ee) and moderate yield with 5 mol% of the scandium catalyst. The 7-, 8-substituted chromanone lactones **3q–3u** could be isolated as nearly single enantiomers (>19 : 1 dr and >99% ee), but 6-substitution resulted in decreased diastereoselectivity (**3o** and

3p). Halogen-substitution at the 6- or 8-position allowed further coupling reactions to prepare the corresponding dimeric natural products. Due to poor solubility of these polysubstituted chromones in tetrahydrofuran, the reaction solvent needed to be changed to a mixture of 1,1,2,2-tetrachloroethane and acetonitrile with increased reaction temperature (50 °C).

To evaluate the synthetic potential of the protocol, chromanone lactone **3a** was prepared on a gram-scale with 1 mol% of



Scheme 2 The gram scale synthesis of **3a** and transformations into natural products.

the $L_3\text{-PrEt}_2/\text{Sc}(\text{OTf})_3$ complex with no deterioration in yield or stereoselectivity (Scheme 2a). Then, the synthetic utility of the chromanone lactones in the related natural products was probed. Treatment of 5-methoxy-7-methyl substituted chromanone lactone **3s** with BBr_3 solution afforded 5-OH based gonytolide C, the prevalent backbone of chromones in natural products. The absolute configuration of **3s** and its precursor **3sa** was assigned as (2*R*,5'*S*) according to the structure of gonytolide C (Scheme 2b).^{2d} Following Porco's work,^{3,4} gonytolide C can be smoothly transformed into the tetrahydroxanthone-type natural product blennolide C *via* NaH promoted Dieckmann cyclization. As an important intermediate, gonytolide C can also be converted to gonytolide A^{4d} and G^{2d} (Scheme 1a).

The major diastereomers obtained in our study differed from those obtained in the vinylogous addition of siloxyfuran to benzopyrylium in Porco's study, which is proposed to be due to the swing favored transition state.³ As shown in Table 1 and S6b in the ESI,[†] the addition of a desiccating agent dramatically increased the *anti*-selectivity. The main function of molecular sieves would be protection of the active catalyst from adventitious water. X-ray crystal structures of the Sc^{3+} complexes of N,N' -dioxide ligands¹¹ manifest octahedral geometry in which one *cis*-position is occupied by two ancillary ligands (usually H_2O and the TfO^- ion). If water takes over one coordination to the scandium center prior to the enolized butanolide intermediate, the reaction could take place *via* transition state (i) as


shown in Scheme 3a where the chromone directly binds to the metal center rather than enolate to undergo addition. The *epi*-transition state (ii) is relatively disfavored due to the steric hindrance and long pair repulsion of the oxygens. As a result, the generation of a *syn*-selective product would be a competitive pathway if a trace amount of water existed. However, if the amount of water is minimized in the reaction system, the catalyst would enable the bonding of both chromone and the enolized butanolide simultaneously. As displayed in Scheme 3a, transition state (iii) to generate an *anti*-selective isomer (Re → Re approach or its antipodal) would be preferred over the generation of an opposite *syn*-isomer *via* transition state (iv) (Si → Re approach or its antipodal) where there exists long pair repulsion of oxygens. In addition, the basic condition in the presence of 3 Å MS might accelerate the enolization of butanolide **2a** to improve the reactivity.

The positive role of HFIP in the reactivity is rationalized in Scheme 3b, and as shown in transition state (v), this weakly coordinated alcohol could act as a proton transfer agent to accelerate the isomerization of the enol intermediate after vinylogous addition, giving the adduct **3aa** and regenerating the catalyst for another catalytic cycle. Epimerization of the adduct is common as observed in the previous study,³ which is proposed *via* an isomerization process due to the enolization of the lactone unit (vi). But under the optimized reaction conditions, the epimerization of **3aa** was not observed (see the ESI[†] for details). In addition, the retro-oxo-Michael reaction should be avoided,^{4d} and the byproduct was detected during the column chromatography with silica gel, which could not undergo addition to regenerate the adduct **3aa** under the catalyst system (Scheme 3b).

To explain the observed high enantioselectivity, we propose a model based on the crystal structure of the N,N' -dioxide/ $\text{Sc}(\text{OTf})_3$ complexes¹¹ and previous DFT calculations.¹² The scandium binds the chromone **1a** at the horizontal position, and the enolate of the butanolide substitutes the solvent or counterion at the top. As depicted in transition state (vii) of Scheme 3c, the reaction occurs *via* the Re → Re approach in the open front left quadrant of the catalyst, which leads to the major (2*R*,5'*S*) adduct. In contrast, the Si → Si approach (viii) to yield the (2*S*,5'*R*) adduct is disfavored due to a steric clash of the ester group of chromone and enolized butanolide with the left amide subunit of the catalyst.

Conclusions

In summary, we developed an efficient asymmetric vinylogous addition of butanolide to 2-ester chromones. The reaction proceeds well under mild reaction conditions with good to excellent diastereo- and enantioselectivity (up to 99% ee) and the yields are generally satisfactory. As low as 1 mol% of the chiral N,N' -dioxide/ Sc^{III} complex catalyst enables the generation of a variety of tolerable substituted chromones at different positions. The 5-MeO substituted ones provide access to chromanone lactones, which could be rapidly transformed into optically enriched natural products, including blennolide C, gonytolide C and other related natural products.



Scheme 3 Proposed explanation for the role of additives and selectivity.

Data availability

Further details of experimental procedure, ^1H , $^{13}\text{C}\{^1\text{H}\}$ and $^{19}\text{F}\{^1\text{H}\}$ NMR, UPCC spectra are available in the ESI.†

Author contributions

Y. L. performed the experiments. R. W. repeated data. X. M. F. and X. H. L. supervised the project. X. M. F., X. H. L., S. X. and Y. L. co-wrote the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We appreciate the National Natural Science Foundation of China (21890723 and 21921002) for financial support.

Notes and references

- (a) F. McPhee, P. S. Caldera, G. W. Bemis, A. F. McDonagh, I. D. Kuntz and C. S. Craik, *Biochem. J.*, 1996, **320**, 681; (b) S. Bräse, A. Encinas, J. Keck and C. F. Nising, *Chem. Rev.*, 2009, **109**, 3903; (c) I. N. Siddiqui, A. Zahoor, H. Hussain, I. Ahmed, V. U. Ahmad, D. Padula, S. Draeger, B. Schulz, K. Meier, M. Steinert, T. Kurtán, U. Flörke, G. Pescitelli and K. Krohn, *J. Nat. Prod.*, 2011, **74**, 365; (d) H. Kikuchi, M. Isobe, M. Sekiya, Y. Abe, T. Hoshikawa, K. Ueda, S. Kurata, Y. Katou and Y. Oshima, *Org. Lett.*, 2011, **13**, 4624; (e) T. El-Elimat, M. Figueroa, A. H. Raja, T. N. Graf, S. M. Swanson, J. O. Falkinham III, M. C. Wani, C. J. Pearce and N. H. Oberlies, *Eur. J. Org. Chem.*, 2015, 109; (f) N. Yodsing, R. Lekphrom, W. Sangsopha, T. Aimi and S. Boonlue, *Curr. Microbiol.*, 2018, **75**, 513; (g) L. Xie, M. Li, D. Liu, X. Wang, P. Wang, H. Dai, W. Yang, W. Liu, X. Hu and M. Zhao, *Molecules*, 2019, **24**, 393.
- (a) M. C. Bröhmer, E. Bourcet, M. Nieger and S. Bräse, *Chem.–Eur. J.*, 2011, **17**, 13706; (b) K. S. Masters and S. Bräse, *Chem. Rev.*, 2012, **112**, 3717; (c) L. F. Tietze, L. Ma, J. R. Reiner, S. Jackenkroll and S. Heidemann, *Chem.–Eur. J.*, 2013, **19**, 8610; (d) G. Sudhakar, S. Bayya, V. D. Kadam and J. B. Nanubolu, *Org. Biomol. Chem.*, 2014, **12**, 5601; (e) K. Adachi, S. Hasegawa, K. Katakawa and T. Kumamoto, *Tetrahedron Lett.*, 2017, **58**, 4479.
- T. Qin, R. P. Johnson and J. A. Porco Jr, *J. Am. Chem. Soc.*, 2011, **133**, 1714.
- (a) T. Qin and J. A. Porco Jr, *Angew. Chem.*, 2014, **126**, 3171; *Angew. Chem., Int. Ed.*, 2014, **53**, 3107; (b) T. Qin, S. L. Skraba-Joiner, Z. G. Khalil, R. P. Johnson, R. J. Capon and J. A. Porco Jr, *Nat. Chem.*, 2015, **7**, 234; (c) T. Qin, T. Iwata, T. T. Ransom, J. A. Beutler and J. A. Porco Jr, *J. Am. Chem. Soc.*, 2015, **137**, 15225; (d) X. W. Wu, T. Iwata, A. Scharf, T. Qin, K. D. Reichl and J. A. Porco Jr, *J. Am. Chem. Soc.*, 2018, **140**, 5969.
- (a) L. A. Stubbing, F. F. Li, D. P. Furkert, V. E. Caprio and M. A. Brimble, *Tetrahedron*, 2012, **68**, 6948; (b) J. Liu, Z. Li, P. Tong, Z. Xie, Y. Zhang and Y. Li, *J. Org. Chem.*, 2015, **80**, 1632.
- During the preparation of this manuscript, an enantioselective version was reported, see J. Cui, R. Oriez, H. Noda, T. Watanabe and M. Shibasaki, *Angew. Chem., Int. Ed.*, 2022, **61**, e202203128.
- (a) B. M. Trost, E. Gnanamani, C. A. Kalnmals, C. I. Hung and J. S. Tracy, *J. Am. Chem. Soc.*, 2019, **141**, 1489; (b) J. Cui, N. Kumagai, T. Watanabe and M. Shibasaki, *Chem. Sci.*, 2020, **11**, 7170.
- The review of butenolides: (a) B. Mao, M. Fañanás-Mastral and B. L. Feringa, *Chem. Rev.*, 2017, **117**, 10502, select recent examples of chiral butenolides; (b) D. Zhang, X. Wang, M. C. Zhang, Z. H. Kang, G. L. Xiao, X. F. Xu and W. H. Hu, *CCS Chem.*, 2020, **2**, 432; (c) C. G. Yu, P. Ji, Y. T. Zhang and X. Meng, *Org. Lett.*, 2021, **23**, 7656; (d) Y. C. Chan, X. Y. Wang, Y. P. Lam, J. Wong, Y. L. Steve Tse and Y. Y. Yeung, *J. Am. Chem. Soc.*, 2021, **143**, 12745.
- (a) D. Baek, H. Ryu, J. Y. Ryu, J. Lee, B. M. Stoltz and S. Hong, *Chem. Sci.*, 2020, **11**, 4602; (b) Y. Guan, T. Buiydas, R. F. Lalisie, J. W. Attard, R. Ali, C. Stern, C. M. Hadad and A. E. Mattson, *ACS Catal.*, 2021, **11**, 6325.
- (a) L. Zhou, L. L. Lin, J. Ji, M. S. Xie, X. H. Liu and X. M. Feng, *Org. Lett.*, 2011, **13**, 3056; (b) Q. Zhang, X. Xiao, L. L. Lin, X. H. Liu and X. M. Feng, *Org. Biomol. Chem.*, 2011, **9**, 5748; (c) J. Ji, L. L. Lin, L. Zhou, Y. H. Zhang, Y. B. Liu, X. H. Liu and X. M. Feng, *Adv. Synth. Catal.*, 2013, **355**, 2764; (d) J. Ji, L. L. Lin, Q. Tang, T. F. Kang, X. H. Liu and X. M. Feng, *ACS Catal.*, 2017, **7**, 3763; (e) Q. Tang, L. L. Lin, J. Ji, H. P. Hu, X. H. Liu and X. M. Feng, *Chem.–Eur. J.*, 2017, **23**, 16447.
- (a) Y. L. Liu, D. J. Shang, X. Zhou, X. H. Liu and X. M. Feng, *Chem.–Eur. J.*, 2009, **15**, 2055; (b) CCDC: 1957788 ($\text{L}_3\text{-PiPr}_2/\text{Sc}(\text{OTf})_3$).
- (a) F. Tan, M. P. Pu, J. He, J. Z. Li, J. Yang, S. X. Dong, X. H. Liu, Y. D. Wu and X. M. Feng, *J. Am. Chem. Soc.*, 2021, **143**, 2394. The review of chiral N,N' -dioxides see ; (b) X. H. Liu, L. L. Lin and X. M. Feng, *Acc. Chem. Res.*, 2011, **44**, 574; (c) X. H. Liu, L. L. Lin and X. M. Feng, *Org. Chem. Front.*, 2014, **1**, 298; (d) X. H. Liu, H. F. Zheng, Y. Xia, L. L. Lin and X. M. Feng, *Acc. Chem. Res.*, 2017, **50**, 2621; (e) X. H. Liu, S. X. Dong, L. L. Lin and X. M. Feng, *Chin. J. Chem.*, 2018, **36**, 791; (f) S. X. Dong, X. H. Liu and X. M. Feng, *Acc. Chem. Res.*, 2022, **55**, 4125.