

Commentary

Childhood infections and asthma: at the crossroads of the hygiene and Barker hypotheses

Kelan G Tantisira and Scott T Weiss

Channing Laboratory, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA

Correspondence: Kelan G Tantisira, MD, Channing Laboratory, 181 Longwood Avenue, Boston, MA 02115, USA. Tel: +1 617 525 0863; fax: +1 617 525 0958; e-mail: rekgt@channing.harvard.edu

Received: 18 July 2001

Revisions requested: 26 July 2001

Revisions received: 1 August 2001

Accepted: 1 August 2001

Published: 13 September 2001

Respir Res 2001, **2**:324-327

© 2001 BioMed Central Ltd
(Print ISSN 1465-9921; Online ISSN 1465-993X)

Abstract

The hygiene hypothesis states that childhood asthma develops as a result of decreased exposure to infectious agents during infancy and early childhood. This results in the persistence of the neonatal T helper lymphocyte 2 immunophenotype, thereby predisposing the child to atopic disease. While multiple studies support the hygiene hypothesis in asthma ontogeny, the evidence remains inconclusive; multiple other environmental exposures in early childhood also alter predisposition to asthma. Moreover, the current paradigm for asthma development extends far beyond simple childhood environmental exposures to include fetal development, genetic predisposition, and interactions of the developmental state and genetics with the environment.

Keywords: asthma, child, fetal programming, gene by environment, infection

Introduction

In 1989, David Strachan described decreases in the prevalence of childhood hay fever and atopic dermatitis in association with the presence of older siblings [1]. He concluded that "declining family size, improved household amenities, and higher standards of personal cleanliness have reduced the opportunities for cross-infection in young families. This may have resulted in more widespread clinical expression of atopic disease" [1]. This, and subsequent observations, led to the formulation of the 'hygiene hypothesis'. The biological basis for the hygiene hypothesis lies in the induction of a T helper lymphocyte 1 (Th1) population by bacterial and viral infections, and the resultant deviation from the T helper lymphocyte 2 (Th2) immune responses involved in IgE-mediated allergy [2].

Asthma can be defined as a combination of airway inflammation, often as a result of allergic sensitization, and airway hyperresponsiveness. While family size and child-

hood infections have generally not been associated with airway responsiveness, because of its close relationship with atopy the risk for childhood asthma has also been hypothesized to relate to these factors. Family size and/or attendance at daycare have consistently been associated with decrements in the relative risk of asthma [3-5], although a few studies have demonstrated increased asthma risk with daycare, probably due to an increased prevalence of lower respiratory tract infections [6]. Lower respiratory tract infections in early childhood have uniformly been associated with an increased risk of subsequent asthma [3,5,7]. While the association of non-respiratory childhood infections and the risk of asthma have been generally supportive of a protective effect, the evidence remains inconclusive. In a case-control study of 1659 Italian military cadets, the relative risk of atopy decreased with exposure to orofecal microbes, including *Helicobacter pylori*, *Toxoplasma gondii*, and hepatitis A virus, as diagnosed by serology [8]. Allergic asthma was

present in only one of the 245 cadets positive for at least two of these serologies. A decreased risk of atopy was not noted in relation to the airborne respiratory viral serologies evaluated. While exposure to measles [4] and *Mycobacterium tuberculosis* [9] have also been reported to protect against asthma development, no specific infection to date has consistently been demonstrated to support the tenets of the hygiene hypothesis.

The focus article

The article by Illi *et al* [10] provides convincing data in support of the hygiene hypothesis. In a longitudinal birth cohort of 1314 children followed to the age of 7 years, Illi *et al* compared the prevalence of doctor-diagnosed asthma, current wheeze, and airway hyperresponsiveness with the occurrence of various categories of infection during the first 3 years of life. As expected, lower respiratory tract infections were positively associated with asthma (odds ratio [OR] = 4.46, 95% confidence interval [CI] = 2.07–9.64 for four or more infections versus one or no infection), wheeze (OR = 3.97, 95% CI = 2.06–7.64), and airway responsiveness (OR = 2.14, 95% CI = 1.03–4.43). As a group, however, non-lower respiratory viral infections demonstrated a strong protective effect against the same outcomes (i.e. asthma [OR = 0.16, 95% CI = 0.05–0.54 for eight or more infections versus one or no infection], wheeze [OR = 0.46, 95% CI = 0.14–1.49], and airway responsiveness [OR = 0.24, 95% CI = 0.09–0.68]). The effect was strongest for rhinorrhea and herpetic infections, and was not noted with bacterial, fungal, or gastrointestinal infections. The limitations of this study included follow-up of only 71% of infants (allowing for potential bias) and no direct measures of infectious burden. Additionally, the cohort's primary study design was to evaluate infants at high risk for atopy, potentially limiting the generalizability of the results. Nevertheless, this was a well-designed study, the particular strengths of which included consistency of associations across several asthma phenotypes, including airway hyperresponsiveness. Moreover, the associations noted were strong and there appeared to be a dose–response effect between the number of infections and the outcome. For instance, the relative odds for airway responsiveness were 0.50 for two to four viral infections versus one or no infections, 0.34 for five to seven infections, and 0.24 for eight or more infections.

A broader paradigm for asthma development


While early exposure to infectious burden may affect the Th1/Th2 balance in the developing neonate, other environmental risk factors for the development of childhood asthma may also affect immune system ontogeny. These risk factors include the protective effect of early exposures to farm animals (via endotoxin) [11] and to household pets (via immune tolerance) [12], and the increased asthma risk associated with antibiotic usage (via suppression of the

gut flora) [13]. However, other early childhood exposures increasing risk for the development of asthma, such as household polyvinylchloride exposure and environmental tobacco smoke, do not have readily apparent effects on the immune system. Overall, it is clear that variations in early life environment are significant risk factors for the development of childhood asthma, but that these variations are not sufficient to cause asthma by themselves. The evolving paradigm supports a combination of genetics, *in utero* development, and early life environment in the origin of asthma (Fig. 1).

The idea that fetal programming can affect the subsequent development of chronic disease was popularized by Barker and colleagues [14], and it is often referred to as the Barker hypothesis. This hypothesis of fetal origins proposes that these diseases originate through adaptations that the fetus makes when it is undernourished. Such diseases may be consequences of 'programming', whereby a stimulus or insult at a critical, sensitive period of early life results in long-term changes in physiology or metabolism [15]. A prominent example of this is the increased risk of asthma in low birth weight infants [7,16]. While the Barker hypothesis focuses on alterations in the developing fetus due to nutrition, there is evolving evidence that many other factors involving the *in utero* environment and fetal gene by environment interactions can affect fetal programming and the subsequent development of disease.

The maternal–fetal interface appears to play a particularly prominent role in the subsequent development of asthma; risk of childhood asthma is greater for infants with a maternal history of asthma than those with a paternal history of asthma [17]. Whether this maternal influence is primarily genetic, environmental, or both has yet to be fully elucidated. Maternal imprinting, a phenomenon whereby the maternal genes are preferentially expressed in the fetus, has been noted in linkage studies of atopy [18] and asthma [19], and probably explains the familial aggregation pattern of pulmonary function noted within asthmatics [20]. Maternal environmental characteristics, as they relate to the fetus, including smoking [21] and infections [22] during pregnancy, have also been strongly associated with subsequent asthma development.

Within the developing fetus, a weak Th2 response normally develops as a result of fetal priming to help maintain pregnancy. *In utero* exposure to allergens may significantly enhance the Th2 response [23,24]. The critical period for this programmed response of the fetus to allergens is thought to be from 5 to 7 months of gestation [25]. The fetal response to allergens may also vary according to genetic predisposition. In a recent study of fetal IgE production, endogenous production of IgE was noted at as early as 8 weeks' gestation, but only in those fetuses with at least one *IL4RA**A1902G allele [26]. This genotype

Figure 1

Overview of the development of childhood asthma. Fetal environmental and genetic influences lead to a specific immunophenotype in the newborn. Subsequent interactions with external environmental exposures (including infections), in conjunction with genetic predisposition, lead to the development of asthma. It is probable that all three components (developmental, genetic, and environmental) are necessary for asthma to occur.

would predispose to early elevations of IgE levels, which have been correlated with asthma and atopy risk [27].

The result of interactions between genetics and the *in utero* environment is a Th2 skewed immunophenotype in the neonate. Those infants with IL-13 producing Th2 lymphocytes may be particularly predisposed to develop asthma [28]. The risk of atopy and asthma is related to failure of the neonate to generate interferon- γ and the resultant failure to transition from the Th2 to the Th1 immunophenotype [29,30]. This failure to transition probably occurs only within genetically susceptible individuals [29], under environmental influences from both the *in utero* and postnatal state (Table 1). One common hypothesis to support this gene by environment interaction in the neonate has been the role of endotoxin and polymorphisms of the CD14 gene. Endotoxin is a component of Gram-negative bacterial cell walls, is fairly ubiquitous in nature, and is an accurate indicator of the cleanliness of indoor environments in urban areas. CD14 recognizes and binds to the endotoxin. A C \rightarrow T polymorphism at position 159 of the 5' flanking region of the CD14 gene has been identified. The homozygous TT genotype has been associated with increases in the serum CD14 level, decreases in serum IgE concentrations, and decreases in the number of positive skin tests in atopic individuals [31]. Whether this association is also true in the development of asthma is under active investigation.

Conclusion

Overall, there appears to be supportive evidence for the role of early exposure to non-respiratory infections as a protective factor against the development of childhood asthma. However, this is likely to be only one of several independent environmental risk factors for asthma in the neonate. Moreover, these postnatal environmental risk factors are themselves only part of a greater scheme that includes fetal development and genetic predisposition. Together, these three broad influences (developmental,

Table 1

Environmental risk factors in the development of asthma

| <i>In utero</i> environmental factors | Postnatal environmental factors |
|---------------------------------------|------------------------------------|
| Maternal diet | Infant diet (breast versus bottle) |
| Maternal/fetal infections | Maternal smoking |
| Bacterial | Allergens |
| Viral | Endotoxin |
| Parasitic | Daycare |
| Maternal smoking | Farm animals |
| Allergens | Antibiotics |
| Endotoxin | |

genetic, and environmental), along with their complex interactions, are currently the most important factors in the ontogeny of childhood asthma.

Acknowledgement

Dr Tantisira is supported by the National Institutes of Health: 2T32 HL07427, Clinical Epidemiology of Lung Diseases.

References

1. Strachan DP: **Hay fever, hygiene, and household size.** *BMJ* 1989, **299**:1259-1260.
2. Romagnani S: **Human TH1 and TH2 subsets: regulation of differentiation and role in protection and immunopathology.** *Int Arch Allergy Immunol* 1992, **98**:279-285.
3. Nafstad P, Magnus P, Jaakkola JJ: **Early respiratory infections and childhood asthma.** *Pediatrics* 2000, **106**:E38.
4. Bodner C, Anderson WJ, Reid TS, Godden DJ: **Childhood exposure to infection and risk of adult onset wheeze and atopy.** *Thorax* 2000, **55**:383-387.
5. Ponsonby AL, Couper D, Dwyer T, Carmichael A, Kemp A: **Relationship between early life respiratory illness, family size over time, and the development of asthma and hay fever: a seven year follow up study.** *Thorax* 1999, **54**:664-669.
6. Nystad W, Skrondal A, Magnus P: **Day care attendance, recurrent respiratory tract infections and asthma.** *Int J Epidemiol* 1999, **28**:882-887.
7. Gold DR, Burge HA, Carey V, Milton DK, Platts-Mills T, Weiss ST: **Predictors of repeated wheeze in the first year of life: the relative roles of cockroach, birth weight, acute lower respiratory illness, and maternal smoking.** *Am J Respir Crit Care Med* 1999, **160**:227-236.
8. Matricardi PM, Rosmini F, Riondino S, Fortini M, Ferrigno L, Rapicetta M, Bonini S: **Exposure to foodborne and orofecal microbes versus airborne viruses in relation to atopy and allergic asthma: epidemiological study.** *BMJ* 2000, **320**:412-417.
9. von Hertzen LC: **Puzzling associations between childhood infections and the later occurrence of asthma and atopy.** *Ann Med* 2000, **32**:397-400.
10. Illi S, von Mutius E, Lau S, Bergmann R, Niggemann B, Sommerfeld C, Wahn U: **Early childhood infectious diseases and the development of asthma up to school age: a birth cohort study.** *BMJ* 2001, **322**:390-395.
11. Von Ehrenstein OS, Von Mutius E, Illi S, Baumann L, Bohm O, von Kries R: **Reduced risk of hay fever and asthma among children of farmers.** *Clin Exp Allergy* 2000, **30**:187-193.
12. Nafstad P, Magnus P, Gaarder PI, Jaakkola JJ: **Exposure to pets and atopy-related diseases in the first 4 years of life.** *Allergy* 2001, **56**:307-312.
13. von Mutius E, Illi S, Hirsch T, Leupold W, Keil U, Weiland SK: **Frequency of infections and risk of asthma, atopy and airway hyperresponsiveness in children.** *Eur Respir J* 1999, **14**:4-11.

14. Barker DJ, Martyn CN: **The maternal and fetal origins of cardiovascular disease.** *J Epidemiol Community Health* 1992, **46**:8-11.
15. Barker DJ: **In utero programming of chronic disease.** *Clin Sci (Colch)* 1998, **95**:115-128.
16. Brooks AM, Byrd RS, Weitzman M, Auinger P, McBride JT: **Impact of low birth weight on early childhood asthma in the United States.** *Arch Pediatr Adolesc Med* 2001, **155**:401-406.
17. Litonjua AA, Carey VJ, Burge HA, Weiss ST, Gold DR: **Parental history and the risk for childhood asthma. Does mother confer more risk than father?** *Am J Respir Crit Care Med* 1998, **158**:176-181.
18. Kurz T, Strauch K, Heinzmann A, Braun S, Jung M, Ruschendorf F, Moffatt MF, Cookson WO, Inacio F, Ruffilli A, Nordskov-Hansen G, Peltre G, Forster J, Kuehr J, Reis A, Wienker TF, Deichmann KA: **A European study on the genetics of mite sensitization.** *J Allergy Clin Immunol* 2000, **106**:925-932.
19. Daniels SE, Bhattacharya S, James A, Leaves NI, Young A, Hill MR, Faux JA, Ryan GF, le Souef PN, Lathrop GM, Musk AW, Cookson WO: **A genome-wide search for quantitative trait loci underlying asthma.** *Nature* 1996, **383**:247-250.
20. Holberg CJ, Morgan WJ, Wright AL, Martinez FD: **Differences in familial segregation of FEV1 between asthmatic and nonasthmatic families. Role of a maternal component.** *Am J Respir Crit Care Med* 1998, **158**:162-169.
21. Weitzman M, Gortmaker S, Walker DK, Sobol A: **Maternal smoking and childhood asthma.** *Pediatrics* 1990, **85**:505-511.
22. Xu B, Pekkanen J, Jarvelin MR, Olsen P, Hartikainen AL: **Maternal infections in pregnancy and the development of asthma among offspring.** *Int J Epidemiol* 1999, **28**:723-727.
23. Jones AC, Miles EA, Warner JO, Colwell BM, Bryant TN, Warner JA: **Fetal peripheral blood mononuclear cell proliferative responses to mitogenic and allergenic stimuli during gestation.** *Pediatr Allergy Immunol* 1996, **7**:109-116.
24. Piccinni MP, Mecacci F, Sampognaro S, Manetti R, Parronchi P, Maggi E, Romagnani S: **Aeroallergen sensitization can occur during fetal life.** *Int Arch Allergy Immunol* 1993, **102**:301-303.
25. Donovan CE, Finn PW: **Immune mechanisms of childhood asthma.** *Thorax* 1999, **54**:938-946.
26. Lima JO, Zhang L, Atkinson TP, Philips J, Dasanayake AP, Schroeder HW Jr: **Early expression of epsilon, CD23 (FcepsilonRII), IL-4Ralpha, and IgE in the human fetus.** *J Allergy Clin Immunol* 2000, **106**:911-917.
27. Hansen LG, Halken S, Host A, Moller K, Osterballe O: **Prediction of allergy from family history and cord blood IgE levels. A follow-up at the age of 5 years. Cord blood IgE. IV.** *Pediatr Allergy Immunol* 1993, **4**:34-40.
28. Spinozzi F, Agea E, Russano A, Bistoni O, Minelli L, Bologni D, Bertotto A, de Benedictis FM: **CD4+IL13+ T lymphocytes at birth and the development of wheezing and/or asthma during the 1st year of life.** *Int Arch Allergy Immunol* 2001, **124**:497-501.
29. Holt PG, Macaubas C, Prescott SL, Sly PD: **Primary sensitization to inhalant allergens.** *Am J Respir Crit Care Med* 2000, **162**:S91-S94.
30. Prescott SL, Macaubas C, Smallacombe T, Holt BJ, Sly PD, Holt PG: **Development of allergen-specific T-cell memory in atopic and normal children.** *Lancet* 1999, **353**:196-200.
31. Baldini M, Lohman IC, Halonen M, Erickson RP, Holt PG, Martinez FD: **A Polymorphism* in the 5' flanking region of the CD14 gene is associated with circulating soluble CD14 levels and with total serum immunoglobulin E.** *Am J Respir Cell Mol Biol* 1999, **20**:976-983.