

Poster presentation

Open Access

Exploring sparse connectivity in the motor system using multivariate autoregression analysis

Rafael Rodriguez-Rojas^{*1}, Mayrim Vega-Hernandez², Agustín Lage², Jose Sanchez², Maylen Carballo¹, Jorge Bosh² and Pedro Valdes-Sosa²

Address: ¹Brain Images Processing Group, International Center for Neurological Restoration, Havana, Cuba and ²Neurophysics Department, Cuban Neuroscience Center, Havana, Cuba

Email: Rafael Rodriguez-Rojas^{*} - rafael.rguez@infomed.sld.cu

^{*} Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P138 doi:10.1186/1471-2202-8-S2-P138

© 2007 Rodriguez-Rojas et al; licensee BioMed Central Ltd.

Background

Multivariate autoregressive (MAR) models can be used in the identification of causal relations from functional MRI time series. Connectivity information is extracted from large neural networks combining graphical modeling methods and Granger causality. The aim of this paper is to demonstrate the feasibility of working with the MAR models to identify functional circuits in the human motor

system, and demonstrates their application to data of motor performance in patients with Parkinson's disease (PD).

Methods

In this work we incorporate a family of linear methods called penalized linear regression that were designed to deal with problems having a large set of variables (i.e.




Figure 1

Realistic rendering of effective connections in the motor system. Note involvement of areas related to motor performance. **MI**: Primary motor cortex, **S1**: Primary somatosensory cortex, **PM**: Premotor cortex, **SMA**: Supplementary motor area, **ASC**: Associative parietal cortex, **STR**: Striatum, **TAL**: Thalamus, **CER**: Cerebellum.

Table 1: Effects of L-Dopa treatment on strength of inter-regional path coefficients

Connection	OFF Medication	ON Medication
ASC – PRE	0.035	0.789
TAL – PM	0.062	0.154
TAL – SMA	0.093	0.256
STR – TAL	0.018	-0.386
TAL – SMA	0.093	0.256

brain structures) and a relatively small set of observations (i.e. fMRI time points). One parkinsonian patient with early stage akinetic PD was studied by fMRI during the "drug-off" state and after reaches the "drug-on" state (table 1).

Results

The statistically most relevant connections from the connectivity matrix, in parkinsonian state, are summarized in the realistic rendering shown in Figure 1. These results indicate that the components of the basal ganglia-thalamus-cortical circuit were functionally connected to each other, but also functional connections in the cortico-cerebello-thalamo-cortical pathway are evidenced.

Conclusion

In opposition to widely spread methods for connectivity analysis, the proposed algorithm does not rely on pre-conditioned connections between regions from anatomical models. The penalized regression techniques expand the basic idea of ordinary least squares by means of the addition of new terms to the minimization equation. Our results support that MAR models form a valuable and feasible approach to study functional circuits in the human motor system, in normal and disease condition.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

