

Meeting abstract

Specific effects of Lef-1 splice variants on the regulation of gene expression in pancreatic cancer cells

S Jesse*¹, K Giehl¹, A König² and A Menke¹

Address: ¹Universitätsklinikum Ulm, Innere Medizin I, AG Molekulare Zellbiologie, Ulm, Germany and ²Abteilung für Gastroenterologie und Endocrinologie, Universitätsklinikum Giessen und Marburg, Germany

* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

Cell Communication and Signaling 2009, **7**(Suppl 1):A32 doi:10.1186/1478-811X-7-S1-A32

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A32>

© 2009 Jesse et al; licensee BioMed Central Ltd.

The lymphoid enhancer factor (Lef-1) belongs to the nuclear transducers of canonical Wnt-signalling in embryogenesis and cancer. Lef-1 acts, in cooperation with beta-catenin, as a context-dependent transcriptional activator or repressor thereby influencing multiple cellular functions such as proliferation, differentiation and migration.

Here we report an increased Lef-1 expression in human pancreatic cancer, which correlates with advanced tumour stages. As demonstrated by RT-PCR analysis, pancreatic carcinoma exhibit two different transcripts present in pancreatic carcinomas. One transcript was identified as the full length Lef-1 (Lef-1 FL), whereas the second, shorter transcript, lacked exon VI (Lef-1 exon VI) compared to the published sequence. Comparative analysis of these two Lef-1 variants revealed different cellular effects after transient expression in pancreatic carcinoma cells. Forced expression of Lef-1 exon VI in pancreatic carcinoma cells inhibited E-cadherin expression and resulted in reduced cellular aggregation and increased cell migration compared to cells expressing full length Lef-1. Expression of Lef-1 FL, but not the newly identified Lef-1 exon VI, induced expression of the cell cycle regulating proteins c-myc and cyclin D1 and resulted in enhanced cell proliferation.

Thus, our findings implicate that expression of alternatively spliced isoforms of Lef-1 are involved in the determination of proliferative or migratory characteristics of pancreatic carcinoma cells.