



Review Article

A bird's-eye overview of molecular mechanisms regulating feed intake in chickens—with mammalian comparisons



Anastasiia R. Volyanskaya ^{a, b, *}, Ilya R. Akberdin ^{a, b, c}, Mikhail A. Kulyashov ^{a, b, c}, Ivan S. Yevshin ^{b, c}, Michael N. Romanov ^{d, e}, Elena I. Shagimardanova ^f, Oleg A. Gusev ^{f, g, h}, Fedor A. Kolpakov ^{b, c}

^a Department of Natural Sciences, Novosibirsk State University, Novosibirsk, Russia

^b Biosoft.Ru, Ltd., Novosibirsk, Russia

^c Sirius University of Science and Technology, Sirius, Russia

^d School of Biosciences, University of Kent, Canterbury, UK

^e L.K. Ernst Federal Research Centre for Animal Husbandry, Dubrovitsy, Podolsk, Russia

^f Regulatory Genomics Research Center, Institute of Fundamental Medicine and Biology, Kazan Federal University, Kazan, Russia

^g Life Improvement By Future Technologies (LIFT) Center, Moscow, Russia

^h Intractable Disease Research Center, Juntendo University, Tokyo, Japan

ARTICLE INFO

Article history:

Received 6 September 2023

Received in revised form

29 December 2023

Accepted 10 January 2024

Keywords:

Chicken

Feed intake

Hypothalamus

Neuropeptide

Signaling pathway

Hormone

ABSTRACT

In recent decades, a lot of research has been conducted to explore poultry feeding behavior. However, up to now, the processes behind poultry feeding behavior remain poorly understood. The review generalizes modern expertise about the hormonal regulation of feeding behavior in chickens, focusing on signaling pathways mediated by insulin, leptin, and ghrelin and regulatory pathways with a cross-reference to mammals. This overview also summarizes state-of-the-art research devoted to hypothalamic neuropeptides that control feed intake and are prime candidates for predictors of feeding efficiency. Comparative analysis of the signaling pathways that mediate the feed intake regulation allowed us to conclude that there are major differences in the processes by which hormones influence specific neuropeptides and their contrasting roles in feed intake control between two vertebrate clades.

© 2024 The Authors. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Molecular mechanisms regulating feed intake during vertebrate ontogenesis are essential for maintaining growth and meat production in livestock, including poultry (Swennen et al., 2007; Everaert et al., 2019; Richards and Proszkowiec-Węglarz, 2007). Therefore, disentangling these mechanisms orchestrating feeding

behavior and energy expenditure is important for commercial breeding and meat industry.

Eating behavior is controlled by central and peripheral regulation, which is coordinated by the nervous and digestive systems. Appetite regulation is provided through the perception of peripheral signals from the external environment and internal physiological signals that convey information about energy and nutritional status (Honda, 2021). The integration of hormonal and nutritional metabolic inputs that control feeding behavior and energy homeostasis is carried out by neural networks in the hypothalamus, “the center of satiety and hunger control”.

Several neuropeptides are expressed in hypothalamic neurons when stimulated by hormones such as ghrelin, insulin, and leptin, providing long-term regulation of eating behavior. Hormone signaling through the hypothalamic neuronal networks is closely related to the adenosine-monophosphate-activated protein kinase/mammalian target of rapamycin (AMPK/mTOR) signaling pathway, which serves as the main sensor of cellular energy. To date, several

* Corresponding author.

E-mail address: a.voliaskaia@alumni.nsu.ru (A.R. Volyanskaya).

Peer review under responsibility of Chinese Association of Animal Science and Veterinary Medicine.



studies have tackled individual components of signaling pathways that mediate the formation of feed intake in poultry and mammals. However, the understanding of molecular mechanisms regulating feed intake in chickens remains very limited.

It is worth noting that the molecular mechanisms and factors that regulate feed intake in birds are not comprehensively investigated compared to mammals. In this regard, we overview state-of-the-art knowledge and data on this topic by combining essential differences between two classes of vertebrates and highlighting blank spots in the regulatory mechanisms for chickens.

2. The hypothalamus as a central regulator of feed intake

In birds as well as mammals, the hypothalamus is crucial for controlling feeding behavior by integrating all peripheral and central signals and generating satiety or hunger states. The central nervous system (CNS) receives information about the nutritional and metabolic state via a variety of peripheral signals, including peptide hormones. These signals influence a number of hypothalamic neuropeptides and complex neural circuits in the hypothalamus, which set off the appropriate responses related to feed intake (Kuenzel et al., 1999). The regulation of chicken feed intake and energy homeostasis appears to be comparable to that of mammals, which is implemented by means of neuropeptides produced in the hypothalamic nuclei (Boswell, 2005; Richards and Proszkowiec-Weglarczyk, 2007). Both mammals and birds are assumed to have satiety areas in the ventromedial and lateral hypothalamus (Kuenzel et al., 1999). In addition to these hypothalamic structures, the paraventricular nucleus (PVN) and the infundibular nucleus (IN), the avian equivalent of the mammalian arcuate nucleus (ARC), are also involved in the control of feed intake (Tachibana and Tsutsui, 2016). Many common neuropeptides between mammals and chickens have been shown to be involved in the control of feed intake (Denbow and Cline, 2015). Two types of hypothalamic neuropeptides that regulate feeding behavior can be identified: some suppress eating behavior and are called anorexigenic neuropeptides, whereas others stimulate eating behavior and are called orexigenic neuropeptides.

Two types of neuron populations in the IN of the hypothalamus are important conduits through which peripheral signals that affect appetite are integrated (Boswell, 2005; Wynne et al., 2005). One of them expresses the pro-opiomelanocortin (POMC) and cocaine- and amphetamine-regulated transcript (CART) mRNAs and the other one the neuropeptide Y (NPY) and agouti-related protein (AgRP) mRNAs (Boswell, 2005). Hypothalamic neuropeptides such as adrenocorticotrophic hormone (ACTH; also, adrenocorticotropin, or corticotropin) and α -melanocyte-stimulating hormone (α -MSH) derived from the POMC precursor and CART are expected to be

anorexigenic in the chickens. NPY and AgRP exert an anabolic effect and are representative candidates for orexigenic neuropeptides (Tachibana, 2016). In addition to orexigenic neuropeptides, these neurons also produce the inhibitory neurotransmitter γ -aminobutyric acid (GABA), which can act in a local circuit to reduce the activity of POMC neurons (Rau and Hentges, 2019). The expression of the neuropeptide genes *AgRP*, *NPY*, *POMC*, and *CART* was determined in the IN of several species of birds, including chickens, with the expression of *AgRP* and *NPY* colocalised in individual IN neurons (Boswell et al., 2002; Boswell and Dunn, 2017; Gerets et al., 2000; Wang et al., 2001; Yuan et al., 2009).

3. Functions of hypothalamic neuropeptides in regulating feed intake

In mammals and several bird species, including chickens, it is well known that the neuropeptide gene expression in neurons of the arcuate nucleus is affected by nutritional status and changes in energy levels (Boswell and Dunn, 2017). In recent decades, numerous studies have been conducted to explore the function of orexigenic AgRP, NPY, and anorexigenic α -MSH neuropeptides that regulate feeding behavior in chickens (Tables 1 and 2).

The *POMC* and *AgRP* genes encoding neuropeptides, along with melanocortin receptors (MC-R), constitute the central melanocortin system (Boswell and Dunn, 2015). MC-R neurons project from the mammalian arcuate nucleus to the PVN, containing a high density of MC-R. Activation of these receptors causes a decrease in feed consumption and an increase in energy use (Gali Ramamoorthy et al., 2015).

Chicken melanocortin signaling in the hypothalamus is mediated by specific subtypes of MC-R, i.e., melanocortin receptors 3 (MC3R) and 4 (MC4R), by binding to such agonists as AgRP and α -MSH/ACTH (Boswell and Dunn, 2015). Neuropeptides α -MSH and ACTH act as activators of chicken melanocortin receptors (Zhang et al., 2017). As expected, the central injection of α -MSH in broiler neonatal chicks followed by administration of an MC4R antagonist led to the reduction of the α -MSH anorexigenic effect, suggesting that its effect is MC4R-mediated (Saneyasu et al., 2011a). Indeed, α -MSH mediated anorexigenic effect in chickens, suppressing feed intake in both broilers and layers after intracerebroventricular (ICV) injection (Cline and Smith, 2007; Honda et al., 2007, 2012; Kawakami et al., 2000; Saneyasu et al., 2011a). In contrast, β -MSH causes an anorexigenic effect in layers (Honda et al., 2012), but not in broiler chickens (Honda et al., 2012; Saneyasu et al., 2011a).

By contrast to α -MSH and ACTH, AgRP can act as an inverse agonist for MC-Rs. Furthermore, AgRP can also antagonise α -MSH/ACTH action on these receptors (Zhang et al., 2017). Indeed, central injection of AgRP with α -MSH attenuated the anorexigenic effect of

Table 1
Neuropeptide central injections demonstrating effects on feeding behavior in chickens.

| Neuropeptide category | Neuropeptide | Type of chicken breed | Breed and age of chicken | Change in feed intake and references |
|---------------------------|-----------------------|-----------------------|------------------------------------|--------------------------------------|
| Orexigenic neuropeptide | NPY | Broiler | 2- and 3-d-old Cobb | Increased (Ando et al., 2001) |
| | | Layer | 2-, 4- and 8-d-old chunky | Increased (Saneyasu et al., 2011b) |
| | AgRP | Broiler | 6-d-old | Increased (Tachibana et al., 2006) |
| | | Layer | 2-, 4- and 8-d-old White Leghorn | Increased (Saneyasu et al., 2011b) |
| Anorexigenic neuropeptide | POMC (α -MSH) | Broiler | 3-d-old Cobb | No change (Tachibana et al., 2001) |
| | | Layer | 4-d-old Boris Brown | Increased (Tachibana et al., 2001) |
| | | Broiler | 5-d-old Cobb 500 | Decreased (Cline and Smith, 2007) |
| | | | 8-d-old chunky | Decreased (Honda et al., 2012) |
| | | | 2-d-old Cobb | Decreased (Kawakami et al., 2000) |
| | | | 1-d-old chunky | Decreased (Saneyasu et al., 2011a) |
| | | Layer | 8-d-old White Leghorn | Decreased (Honda et al., 2007, 2012) |
| | | 6-d-old | Decreased (Tachibana et al., 2007) | |

NPY = neuropeptide Y; AgRP = agouti-related peptide; POMC = proopiomelanocortin; α -MSH = α -melanocyte-stimulating hormone.

Table 2
Changes in expression of neuropeptides in the hypothalamus of feed-restricted chickens.

| Neuropeptide category | Neuropeptide | Type of chicken breed | Breed and age of chicken | Fasting period | Change in neuropeptide expression and references | | |
|-------------------------|---------------------|------------------------|---------------------------|-----------------------------------|--|-------------|-------------------------------|
| Orexigenic neuropeptide | NPY | Broiler | 14-d-old yellow-feathered | 48 and 24 h | Increased (Fang et al., 2014) | | |
| | | | 7-d-old Arbor Acres | 48 h | Increased (Song et al., 2012) | | |
| | | | 1-d-old Ross × Cobb | 48 h | Increased (Higgins et al., 2010) | | |
| | | | 7-d-old Arbor Acres | 24 h | No change (Liu and Zhu, 2012) | | |
| | | | 21-d-old Ross 308 | 12 h | Increased (Kewan et al., 2021) | | |
| | | Layer | 6-wk-old Ross 308 | Chronic feed restriction for 6 wk | No change (Dunn et al., 2013) | | |
| | | | 21-d-old White Leghorn | 12 h | Increased (Kewan et al., 2021) | | |
| | | | 10-d-old White Leghorn | 4 d | Increased (Kameda et al., 2001) | | |
| | | | AgRP | Broiler | 14-d-old yellow-feathered | 48 and 24 h | Increased (Fang et al., 2014) |
| | | | | | 7-d-old Arbor Acres | 48 h | Increased (Liu and Zhu, 2012) |
| | 7-d-old Arbor Acres | 48 h | | | Increased (Song et al., 2012) | | |
| | 1-d-old Ross × Cobb | 48 h | | | Increased (Higgins et al., 2010) | | |
| | 21-d-old Ross 308 | 12 h | | | Increased (Kewan et al., 2021) | | |
| | Layer | 21-d-old White Leghorn | 6-wk-old Ross 308 | Chronic feed restriction for 6 wk | Increased (Dunn et al., 2013) | | |
| | | | 21-d-old White Leghorn | 12 h | No change (Kewan et al., 2021) | | |

NPY = neuropeptide Y; AgRP = agouti-related peptide.

α -MSH in both neonatal broilers and layer chicks (Kawakami et al., 2000; Tachibana et al., 2001). Under ad libitum feeding settings, AgRP injection increased feed consumption in neonatal layer chicks but not in broilers, indicating that the orexigenic impact of endogenous AgRP varies between the two breeds, at least at the neonatal stage (Tachibana et al., 2001).

In both chickens and mammals, NPY is regarded as a powerful stimulant of feeding behavior (Greene et al., 2022). Indeed, central injections of NPY resulted in stimulation of feed intake in broilers (Ando et al., 2001; Saneyasu et al., 2011b) and slow-growing chicks (Saneyasu et al., 2011b; Tachibana et al., 2001). After 4 d of feed deprivation, NPY mRNA and peptide levels increased markedly in the hypothalamic IN nuclei of layers, suggesting its involvement in the regulation of feed intake (Kameda et al., 2001).

Fasting for 24 and 48 h upregulated the hypothalamic NPY and AgRP gene expression and downregulated POMC in yellow-feathered broiler chicks (Fang et al., 2014). In young Arbor Acres broilers fasted for 48 h, the mRNA expression levels of orexigenic neuropeptides were increased too, but the gene expression of POMC was not affected by the starvation (Song et al., 2012). However, newly hatched broiler chicks after the same period of fasting showed a significant increase in POMC mRNA (Higgins et al., 2010). The discrepancies between the effects of fasting on POMC gene expression is supposed to be due to different breeds and ages of broiler chickens.

Differences in the expression of hypothalamic neuropeptides should also be caused by the period of fasting. In Arbor Acres broiler chicks of the same age under 24-h feed restriction conditions, NPY mRNA levels in the hypothalamus were similar to those in ad libitum-fed chicks. At the same time, starvation led to activation of hypothalamic AgRP and inhibition of POMC gene expression. Refeeding following 24 h of fasting increased mRNA levels of POMC, but decreased mRNA levels of AgRP (Liu and Zhu, 2012). One can assume the increase in appetite during fasting was due to the suppression of the anorexigenic POMC gene expression, and the activation of the orexigenic AgRP, but not the NPY. This is confirmed by the fact that after refeeding the POMC mRNA levels were increased and the AgRP mRNA levels were decreased, and serve as an indicator of satiety.

When fasting for a shorter period of time, 12 h, gene expression levels of NPY in both Ross 308 broiler and layer chicks were significantly elevated and returned to control levels after 12 h of refeeding. In contrast, upregulation of AgRP after starvation was observed only in broilers, and these changes were not reversed by refeeding. Simultaneously, starvation did not influence the mRNA

levels of hypothalamic POMC in either layer or broiler chicks (Kewan et al., 2021). It is likely that 12 h of refeeding is not enough to suppress feed intake in broilers, and appetite control is probably achieved through upregulation of AgRP, but not NPY.

Prolonged feed restriction of Ross 308 broilers for 6 wk showed increased levels of AgRP mRNA, which returned to control levels after unlimited access to feed for 2 d. At the same time, observations were found for NPY, although changes in expression level were not as significant (Dunn et al., 2013). Feed restriction did not change the expression of anorexigenic POMC gene, which was also observed during 12-h fasting in another experiment with Ross 308 (Dunn et al., 2013; Kewan et al., 2021).

When chickens were restricted to feed for 7 d, a significant reduction was identified in POMC hypothalamic expression in both Cobb broilers and layer chicks. However, the suppression of POMC gene expression was more pronounced in layers than in broilers (Hen et al., 2006).

In mammals, α -MSH induces a release of corticotropin-releasing factor (CRF) in a hypothalamic PVN, an area that controls both the hypothalamic–pituitary–adrenal axis (HPA) and feeding behavior (Lu et al., 2003). An anorectic action of CRF is observed in mammals, as well as in chickens, for which it was found to suppress feed intake after central administration in both broilers and layer hens (Denbow et al., 1999). In order to ascertain how α -MSH and CRF neurons interact in chickens, there was a study on the effect of ICV α -MSH injection on corticosterone (CORT) secretion, which is the main stress hormone in birds and is produced when HPA is activated (Tachibana et al., 2007). In particular, it was revealed that in layer chickens, CORT release is induced by central administration of α -MSH (Tachibana et al., 2007). An increase in CORT levels was also observed in broiler chickens after ICV injection of β -MSH (Smith et al., 2007). Moreover, the significantly increased level of hypothalamic CRF mRNA was detected in neonatal broilers after central administration of β -MSH, proposing that CRF participates in the β -MSH anorexigenic pathway (Saneyasu et al., 2013).

4. Feed intake is controlled by the AMPK/mTOR signaling pathway in the hypothalamus

In chickens, there is strong evidence for an association between the control of central melanocortin signaling by hypothalamic energy perception and neuropeptide gene expression in the hypothalamus. At the cellular and organismal levels, AMPK is a central energy sensor essential for maintaining energy homeostasis. AMPK controls energy balance by integrating a diverse set of physiological

signals, such as nutrition status and the metabolic effects of hormones. In the hypothalamus, AMPK completes crucial functions in the control of feed intake and maintaining energy balance and body weight (Hardie, 2014). AMPK signaling is activated under low-energy conditions, elevating energy production and reducing energy consumption. AMPK activation restricts energy expenditure by inhibiting anabolic processes and stimulating catabolic processes, in an attempt to restore cellular energy charge.

The mammalian AMPK complex is composed of one α -catalytic subunit and two regulatory β - and γ -subunits. Chicken AMPK (chAMPK) subunits were shown to have considerable homology to the amino acid sequence of human AMPK (Proszkowiec-Weglarz et al., 2006). This may indicate the similarity of AMPK functioning in chickens and mammals (Fig. 1).

Initial research suggests that AMPK activation occurs due to an increase in the cellular ratio of adenosine monophosphate/adenosine triphosphate (AMP/ATP) and under physiological conditions of negative energy balance, including hunger. Furthermore, AMPK can be activated by direct allosteric binding of AMP. However, there are studies demonstrating that some hormones (e.g., ghrelin) can induce AMPK activity regardless of a change in the AMP/ATP ratio. On the contrary, inhibition of AMPK activity is observed under energy-sufficient conditions (feeding, reduction in AMP/ATP, insulin, and leptin) (Ronnett et al., 2009). In addition, there is evidence that AMPK activation is promoted by phosphorylation of Thr172 in the catalytic domain by the upstream kinases: liver kinase B1 (LKB1) and Ca^{2+} /calmodulin-dependent protein kinase β (CaMKK β).

Immunocytochemical analysis revealed the localization of phosphorylated AMPK in IN, PVN, and other hypothalamic nuclei in chickens, which are closely related to the regulation of feed intake and energy homeostasis. While AMPK phosphorylation was shown to be influenced by nutritional status. Restriction of broiler chickens feeding affected the decrease in the ratio of phosphorylated AMPK to the total amount in the hypothalamus. On the contrary, repeated feeding contributed to a decrease in the level of phosphorylated AMPK (Proszkowiec-Weglarz et al., 2006).

Using the immunofluorescence method, it was shown that the LKB1 protein, a major AMPK upstream kinase, was expressed in chicken hypothalamic cells (Zhang et al., 2021). The starvation of broiler chickens contributed to an increase in hypothalamic levels of phospho-LKB1 compared to total LKB1 (Proszkowiec-Weglarz et al., 2006).

Analysis of AMPK gene expression in the brain, including the hypothalamus, of broiler chickens revealed priority expression of the $\alpha 1$, $\beta 2$ and $\gamma 1$ subunit isoforms. However, alterations in energy status (starvation and feed intake) did not contribute to a significant change in the transcription of the AMPK subunit genes (Proszkowiec-Weglarz et al., 2006). A study of the effect of dietary energy level on the AMPK signaling pathway in the hypothalamus of broiler chickens showed that a high-energy diet led to suppression of appetite and expression of the LKB1 and AMPK $\alpha 1$ genes. On the contrary, a low-energy diet increased AMPK $\alpha 2$ mRNA levels and increased appetite (Hu et al., 2019). Based on the above, it can be concluded that energy availability affects hypothalamic chAMPK, as in mammals. Therefore, the data suggest that the LKB1/




Fig. 1. AMPK/mTOR signaling pathway involved in regulation of feed intake in chickens (created with BioRender.com). AMP = adenosine monophosphate; ATP = adenosine triphosphate; LKB1 = liver kinase B1; AMPK = adenosine-monophosphate activated protein kinase; TSC1/TSC2 = tuberous sclerosis complex 1/2; Rheb = Ras homologue enriched in brain; mTOR = mammalian target of rapamycin; Raptor = regulatory-associated protein of mTOR; 4E-BP 1 = 4E-binding protein 1; S6K = S6 kinase; Akt = protein kinase B; mTORC1 = mTOR complex 1; mTORC2 = mTOR complex 2; Rag = recombination-activating gene protein.

AMPK hypothalamic signaling pathway exists, at least in broilers. However, the functionality of the CaMKK β /AMPK pathway in the chicken hypothalamus requires further study.

A crucial role in the regulation of feeding behavior and maintaining energy balance is assigned to mTOR signaling in the hypothalamus, which responds to changes in nutrient status (Cota et al., 2006). mTOR is a serine–threonine kinase and is a component of two multiprotein complexes, mTORC1 and mTORC2, which have different structures and functions. In the hypothalamus, mTORC1 acts as a sensor of changes in nutrient and energy status in rats: its activity increases with feed intake and decreases with fasting (Cota et al., 2006). The activity of mTORC1 is regulated in response to growth factors, hormones (including leptin, insulin, and ghrelin), and nutrient signaling (glucose and amino acids) (Hu et al., 2016). Amino acids can control the activity of mTORC1 through the Rag proteins (recombination-activating gene), that is, a set of small GTPases (Sancak et al., 2008). Branched-chain amino acids, such as L-leucine and L-arginine, are considered potential activators of mTORC1 (Jewell et al., 2013). Intracerebroventricular injection of L-leucine into broilers and layer chicks significantly stimulated feed intake, while L-arginine did not significantly affect broiler chicken feed intake (Kehinde et al., 2022; Wang et al., 2012).

Growth factors and insulin activate the phosphoinositide 3-kinase/protein kinase B (PI3K/Akt) signaling pathway (a detailed description is presented below), causing the phosphorylation of serine/threonine kinase, protein kinase B (Akt). Phosphoinositide-dependent protein kinase-1 (PDK-1) is an upstream kinase of Akt, which activates Akt by phosphorylation at Thr308. However, mTORC2 is required for maximal Akt activation, which is achieved through phosphorylation at Ser 473 (Dibble and Cantley, 2015). This is accomplished by increasing the activity of mTORC2 due to its phosphorylation by Akt, forming a positive feedback loop with each other (Yang et al., 2015). It was identified that feeding after a fast led to a significant elevation of phosphorylated Akt (Thr308), but not Akt (Ser473) levels in the hypothalamus of layer and broiler chickens (Saneyasu et al., 2018, 2019).

Activated Akt inhibits the tuberous sclerosis complex 1/2 (TSC1/2), which acts as a GTPase activating protein for the small GTPase Rheb (Ras homologue enriched in brain), through multiple phosphorylation of the TSC1 subunit. This contributes to mTORC1 activity stimulation by suppressing Rheb (Inoki et al., 2002).

Similar to AMPK, mTORC1 is involved in energy perception (Fig. 1). However, mTORC1 has the opposite effect of AMPK under conditions of high cellular energy levels. Besides that, AMPK stimulation results in mTORC1 inactivation. Low available cellular energy due to glucose restriction inhibits mTORC1 via activation of AMPK. Two AMPK-catalyzed phosphorylation events counteract the activating effects of Akt on mTORC1: (1) phosphorylation of the TSC-2 subunit in the TSC1/TSC2 complex, which suppresses Rheb-GTP-dependent mTORC1 activation; and (2) AMPK-mediated

phosphorylation of the regulatory-associated protein of mTOR (Raptor). The latter is an essential regulatory mTORC1 subunit whose phosphorylation is required for mTORC1 kinase activity inhibition (Xu et al., 2012).

The central administration of compound C, an inhibitor of AMPK, reduced feed intake in broiler chickens and caused a great decrease in hypothalamic AMPK α phosphorylation and an increase in mTOR phosphorylation. This may indicate that AMPK signaling in the hypothalamus participates in the feed intake control in broiler chickens (Hu et al., 2021). There was also a study in which layer chicks were ICV injected with rapamycin, the mTOR inhibitor, causing inhibition of feed intake (Saneyasu et al., 2018). This suggests that rapamycin blocks hypothalamic mTOR signaling in chickens.

mTORC1 senses alterations in nutrient and hormone levels and regulates translational control of protein synthesis by binding the two downstream targets to Raptor and phosphorylating them. Stimulation of mTORC1 signaling resulted in inactivation of the mRNA translation repressor, the 4E binding protein of eukaryotic initiation factor 4E-binding protein 1 (4E-BP1), and activation of ribosomal protein S6 kinases (S6K) (Hay and Sonenberg, 2004). Upon refeeding after fasting, a significant increase in phosphorylated S6K1 and S6 (downstream target of S6K1) was noted in rats (Cota et al., 2006). In layer chicks fed after a 24-h fast, the level of hypothalamic phosphorylated ribosomal protein S6 increased significantly (Saneyasu et al., 2018). However, refeeding after fasting did not alter hypothalamic S6 phosphorylation in broiler chickens (Saneyasu et al., 2019).

5. Hormonal regulation of feed intake

The hypothalamus integrates information from hormones such as insulin, leptin, and other peptide hormones secreted by the gastrointestinal tract, liver, and adipose tissue.

As a rule, the influence of signaling peptides synthesized in the intestines has a short-term effect on appetite, which in turn does not have a significant role in mediating long-term changes in energy balance and body weight. However, some hormones can contribute to long-term changes in energy balance by activating or inhibiting metabolic pathways (Tables 3 and 4). Ghrelin and leptin have been recognized as key hormones that significantly influence the long-term regulation of energy balance in chickens and mammals. Ghrelin is known as a “hunger hormone” in mammals, because it drives short-term food consumption and manages long-term body weight control (Higgins et al., 2007). Unlike mammals, ghrelin has the opposite effect on feeding behavior in chickens and leads to decreased feed intake (Murugesan et al., 2022). Approximately two decades have passed since the discovery of leptin as a satiety hormone in mammals (Friedman and Halaas, 1998). Later on, the leptin gene in chickens was finally identified and cloned (Seroussi et al., 2016). Leptin is believed to serve as a

Table 3
Effects of hormone injection on feeding behavior.

| Hormone | Type of chicken breed | Breed and age of chicken | Change in feed intake and references |
|---------|-----------------------|--|---|
| Insulin | Broiler | 20-d-old Ross 308 | Decreased (Yousefvand et al., 2018) |
| | | 4-d-old Chunky | No change (Shiraishi et al., 2011b) |
| | | 5-d-old | Decreased (Yousefvand et al., 2020) |
| Layer | Broiler | 8-d-old White Leghorn | Decreased (Honda et al., 2007) |
| | | 3- or 4-d-old Single Comb White Leghorn | Decreased (Shiraishi et al., 2008, 2009; 2011b) |
| Leptin | Broiler | 4-d-old Hubbard \times Cobb-500 | No change (Sims et al., 2017) |
| | | 7-wk-old | Decreased (Denbow et al., 2000) |
| | Layer | 4-wk-old Single Comb White Leghorn | Decreased (Denbow et al., 2000) |
| | | 12-wk-old White Rock high body weight line | No change (Kuo et al., 2005) |
| | | 12-wk-old White Rock low body weight line | Decreased (Kuo et al., 2005) |

Table 4
Effects of hormone injection on the expression of hypothalamic neuropeptides.

| Hormone | Type of chicken breed | Neuropeptide | Breed and age of chicken | Change in neuropeptide expression and references |
|---------|-----------------------|--------------|---|--|
| Insulin | Broiler | NPY | – | – |
| | | AgRP | – | – |
| | | POMC | 7-d-old Ross 308 | No change (Saneyasu et al., 2019) |
| | Layer | NPY | 3- or 4-d-old Single Comb White Leghorn | Decreased (Shiraishi et al., 2008) |
| | | | 8-d-old White Leghorn | No change (Honda et al., 2007) |
| | | AgRP | 3- or 4-d-old Single Comb White Leghorn | No change (Shiraishi et al., 2008) |
| | | | 8-d-old White Leghorn | No change (Honda et al., 2007) |
| | | POMC | 3- or 4-d-old Single Comb White Leghorn | Increased (Shiraishi et al., 2008) |
| | | | 8-d-old White Leghorn | Increased (Honda et al., 2007) |
| | | | 7-d-old White Leghorn | Increased (Saneyasu et al., 2019) |
| Leptin | Broiler | NPY | 3-wk-old Ross | Decreased (Dridi et al., 2005) |
| | | AgRP | 3-wk-old Ross | No change (Dridi et al., 2005) |
| | | POMC | 3-wk-old Ross | No change (Dridi et al., 2005) |
| | Layer | NPY | – | – |
| | | AgRP | – | – |
| | | POMC | – | – |

NPY = neuropeptide Y; AgRP = agouti-related peptide; POMC = proopiomelanocortin.

communication link between peripheral fat reserves and the CNS (Friedman, 2014). Nevertheless, recent findings suggest that this relationship does not hold true in chickens (Friedman-Einat and Seroussi, 2019). In chickens, insulin's function is somewhat conserved compared to mammals and, like leptin, insulin is thought to act as an appetite suppressant peptide. Although significant differences exist in insulin sensitivity and glucose homeostasis, chickens are naturally more glucose intolerant and insulin resistant (Seki et al., 2003). However, it is likely that, depending on the age and breed of chickens with high and low growth rates, different effects are found in the influence of these hormones on feed intake in chickens.

5.1. Insulin

In mammals, the pancreatic hormone insulin is defined as an adiposity signal that regulates blood glucose levels (Woods and Seeley, 2001). However, there is evidence that most likely insulin does not function as an adiposity signal in birds. Since components of the insulin signaling pathway in chicken adipose tissue were found to be insulin insensitive (Dupont et al., 2012), plasma insulin levels and the abdominal fat mass were unrelated (Honda et al., 2015).

According to numerous studies ICV insulin injection suppressed feed consumption in slow-growing chickens (Honda et al., 2007; Shiraishi et al., 2008, 2009, 2011b). The ICV insulin administration to Chunky broiler chickens did not affect their feed intake (Shiraishi et al., 2011b). However, in Ross 308 broilers central insulin injection decreased feed consumption in a dose-dependent manner (Yousefvand et al., 2018, 2020). This difference in insulin-mediated feed intake may be due to different breeds of broiler chickens. Peripheral insulin treatment also did not affect changes in the feed intake of broiler chicks (Liu et al., 2016).

Insulin-dependent signaling pathways control eating behavior. In chickens, insulin receptors were located in several structures of the hypothalamus, while in IN, the presence of insulin receptors was found both in anorexigenic POMC/CART neurons and in orexigenic AgRP/NPY neurons (Shiraishi et al., 2011a). The levels of *InsR* expression in the hypothalamus varied between broilers and layer chickens. Under conditions of free access to feed, the expression of *InsR* in broilers is considerably lower compared to layers, which was accompanied by increased insulin concentrations in broilers. Moreover, feed restriction substantially downregulated the *InsR* expression only in layer chicks, which together may indicate insulin resistance in broiler chicks (Shiraishi et al., 2011b).

Insulin receptor structure is conserved between chickens and mammals. The α subunit of *InsR* is the insulin-binding subunit, while the β subunit exhibits insulin-stimulated tyrosine-specific autophosphorylation (Simon and Leroith, 1986). In its inactive form, the insulin receptor exists as a dimer (Ottenmeyer et al., 2000).

Insulin binding to the receptor results in autophosphorylation of tyrosine residues among β subunits, which allows binding to the insulin receptor substrate (IRS) protein family. It was determined that the insulin receptor substrate-1 (*IRS-1*) gene is expressed in the brain of chickens and phosphorylated at tyrosine residues in response to insulin, at least in the LMH cell line derived from a Leghorn chicken hepatocellular carcinoma that was previously shown to be insulin sensitive (Taouis et al., 2009).

The phosphorylation at tyrosine residues activates IRS proteins and enables PI3K recruitment to the cell membrane and subsequent activation. Phosphorylated IRS proteins interact with the PI3K regulatory subunit (p85), which contributes to activation of the PI3K catalytic subunit (p110), allowing it to phosphorylate membrane-bound phosphatidylinositol 4,5-bisphosphate (PIP2) to promote phosphatidylinositol (3,4,5)-triphosphate (PIP3) synthesis (Engelman et al., 2006). This mediates membrane translocation of serine/threonine kinases PDK-1 and Akt that are binding to membrane-bound lipid PIP3, which leads to activation of Akt by phosphorylation at Thr308 (Boucher et al., 2014). Akt activation in response to insulin promotes *IRS-1* phosphorylation on serine residues, which creates a positive feedback loop for insulin action (Gual et al., 2005).

Central injection of LY294002, a PI3K inhibitor, significantly facilitated feed intake in starving layer chicks. In a separate study, injection of LY294002 significantly prevented insulin-induced elevation in hypothalamic phosphorylated Akt activity (Thr308), indicating that LY294002 inhibits PI3K in the hypothalamus of layer chickens (Saneyasu et al., 2018). The research mediated by the signaling pathway in White Leghorn layer hens and Ross 308 broilers showed that ICV insulin injection reduced feed intake and significantly increased Akt and S6 phosphorylation in the chicken hypothalamus (Saneyasu et al., 2018, 2019).

In mammals, the hypothalamic mTOR/S6K signaling acts as a negative regulator of PI3K-related signaling. Because S6Ks phosphorylate *IRS-1* at several serine residues to promote inhibition of insulin signaling at the *IRS-1* level (Blouet et al., 2008). In broiler chickens, it was demonstrated that repeated feeding and central insulin administration led to increased *IRS-1* serine residue phosphorylation but did not affect the phosphorylation of tyrosine residues in skeletal muscles. This suggests the possibility of a

negative feedback mechanism, which may reduce the activity of IRS-1 by increasing the phosphorylation of serine residues (Duchêne et al., 2008).

In mammals, leptin binding to the receptor leads to the activation of the Janus tyrosine kinase 2 (JAK2) protein, which exists in complex with the receptor, which leads to the activation of IRS-1 and IRS-2, which in turn are phosphorylated by JAK2. Therefore, IRS-1 is a cross-component of the insulin and leptin signaling pathways, indicating that leptin and insulin regulatory effects on appetite may be achieved through the IRS-1/PI3K interaction (Barrios-Correa et al., 2018).

The forkhead box protein O1 (FOXO1) transcription factor is the downstream target of Akt. In mammals, hypothalamic FOXO1 activation promotes increased feed consumption and body mass, while FOXO1 inhibition has the opposite effect. FOXO1 acts as an activator of the orexigenic neuropeptides NPY and AgRP transcription, and as an inhibitor of anorexigenic POMC transcription (Kim et al., 2006). This action is opposite to the effects of the leptin-stimulated activated transcription factor signal transducer and transcription 3 (STAT3), which inhibits AgRP and NPY expression and activates POMC (a detailed description is presented below). Additionally, an increase in the level of FOXO1 expression leads to the formation of a complex with activated phosphorylated STAT3 in the nucleus, blocking binding of STAT3 with the POMC promoter, which contributes to the inhibition of POMC expression activation mediated by leptin signaling (Yang et al., 2009).

Insulin suppresses feed intake by activating the PI3K/Akt signal, leading to inhibition of FOXO1 activity. Once fully activated, Akt becomes capable of phosphorylating its targets, including FOXOs (Manning and Cantle, 2007). Activated Akt induces FOXO1 phosphorylation, followed by exclusion from the nucleus and subsequent proteasomal degradation. Therefore, activation of the PI3K/Akt pathway leads to inhibition of FOXO1 activity, which contributes to a decrease in the expression of orexigenic neuropeptides while simultaneously activating STAT3-mediated transcription of POMC by decreasing the antagonistic effect of FOXO1 on STAT3 (Kodani and Nakae, 2020).

The PI3K/Akt-mediated pathway study in broiler chickens indicated that central insulin administration in contrast to mice (Kim et al., 2006) had no impact on phosphorylated FOXO1. At the same time, insulin did not significantly affect the hypothalamic neuropeptide POMC gene expression (Saneyasu et al., 2019). These may indicate that expression of POMC induced by leptin signaling is not associated with the function of FOXO1 in the hypothalamus, which may be one of the reasons for excessive feed intake in broiler chickens. However, ICV insulin administration in layer chicks increased the level of FOXO1 phosphorylation and hypothalamic POMC expression (Saneyasu et al., 2018). Indeed, the level of hypothalamic POMC and CART mRNA significantly increased after central insulin injection in layer chicks (Honda et al., 2007; Shiraishi et al., 2008). At the same time, insulin ICV injection could inhibit or not affect NPY gene expression (Shiraishi et al., 2008; Honda et al., 2007). Furthermore, the central injection of insulin did not change AgRP gene expression (Honda et al., 2007; Shiraishi et al., 2008). A summary of the insulin signaling pathway and observed differences between chickens and mammals has been summarized in Fig. 2.

5.2. Leptin

The adiposity hormone leptin performs a crucial function in the regulation of feeding behavior and energy balance in mammals. Mammalian leptin is predominantly secreted by adipose tissue and acts as a transmitter of body fat information (Friedman, 2014).

However, leptin expression is not limited to adipose tissue, the stomach and duodenum also produce a significant amount of leptin (Cammisotto and Bendan, 2012).

In birds, including commercial breeds of chickens (broiler and layer chickens), unlike mammals, leptin is not expressed in adipose tissue, and the expression levels of this hormone do not correlate with obesity (Bornelöv et al., 2018; Farkašová et al., 2016; Huang et al., 2014; Resnyk et al., 2015). This suggests that leptin is not a key signal for fat stores (Friedman-Einat and Seroussi, 2019).

In most studies, leptin expression levels in birds, including chickens, were shown to be especially high in the brain, including the cerebellum, hypothalamus, and pituitary gland (Farkašová et al., 2016; Friedman-Einat et al., 2014; Huang et al., 2014; Seroussi et al., 2016). At the same time, the leptin expression patterns in these tissues closely correlated with the expression level of LEPR, indicating a paracrine/autocrine mode of action of this hormone in birds (Friedman-Einat and Seroussi, 2019). The expression of chicken leptin and leptin receptor (LEPR) has also been found in the digestive system (duodenum, caecum, ileum, and pancreas). Leptin is observed in the duodenal mucosa, suggesting that it is involved in short-term appetite regulation (Seroussi et al., 2019).

Leptin regulates eating behavior by binding to leptin receptors in hypothalamic neurons, with the ARC nucleus being the main center sensitive to leptin. In mammals, leptin contributes to the activation of POMC/CART neurons and the suppression of the activity of AgRP/NPY neurons through the corresponding signaling transduction pathway (Van Swieten et al., 2014).

There are many studies that examine the effect of leptin injection on feed intake in chickens. However, the results are quite inconsistent, demonstrating both inhibition of feeding and no effect on appetite in birds. Central administration of recombinant human leptin promotes lower feed intake in broilers and layer chickens (Denbow et al., 2000). However, in slow-growing White Rock chickens selected for body weight, human recombinant leptin caused a reduction in feed intake only in chickens with low body weight (Kuo et al., 2005). ICT injection of the incomplete synthetic chicken leptin peptide did not affect feed intake in Hubbard × Cobb-500 broiler chicks (Sims et al., 2017). As found in another study, leptin contributed to a significant inhibition of feed intake in Ross 308 broiler chicks (Adeli et al., 2020). There was also a study that examined the effect of intraperitoneal injection (IP) of recombinant chicken leptin in broilers and layers of two age groups. In young and adult layers, IP leptin administration resulted in appetite inhibition, while young broilers had no significant effect on feed intake (Cassy et al., 2004). Although the differences between the results of different studies remain incompletely identified, it is possible that the breed, age, or source of leptin (human or chicken recombinant leptin) are responsible for the observed distinction.

Leptin-dependent signaling pathways control eating behavior. To date, the leptin signaling pathway involved in the control of feeding behavior in chickens is poorly understood. However, the experimental data demonstrate the conservative basics of similar signaling pathways in mammals (Fig. 3).

The leptin signaling pathway initiates through the binding of leptin to specific receptors, leptin receptors. This, in turn, promotes activation of several signaling pathways, including JAK2/STAT3 and PI3K/IRS/Akt, which mediate the regulation of feed intake and energy homeostasis. The leptin signaling pathway contributes to inhibition of hypothalamic AMPK through the PI3K/Akt pathway, inducing p70S6K-dependent direct phosphorylation of the AMPK α -subunit at Ser 491 t (Dagon et al., 2012). However, ICV leptin administration in broiler chickens activated AMPK, significantly facilitating AMPK phosphorylation at Thr172 of the α -subunit in the




Fig. 2. Overview of leptin and insulin signaling pathways in the hypothalamus of chickens regulating neuropeptide genes (created with BioRender.com). The mechanism of insulin action in chickens may differ from what is known in mammals. FOXO1 signaling is significantly different between broilers and laying hens. In broilers, there is a slight phosphorylation of FOXO1 and no significant change in the expression of the anorexigenic neuropeptide POMC. While the opposite pattern is shown for laying hens, which is similar to the characteristic of mammals. Note: Pattern of a chicken indicates differences in signaling pathway compared to mammals. IR = insulin receptor; IRS-1 = insulin receptor substrate-1; JAK2 = Janus tyrosine kinase 2; PI3K = phosphoinositide 3-kinase; PIP2 = phosphatidylinositol 4,5-bisphosphate; PIP3 = phosphatidylinositol (3,4,5)-triphosphate; PDK-1 = phosphoinositide-dependent protein kinase-1; Akt = protein kinase B; mTORC1 = mammalian target of rapamycin complex 1; mTORC2 = target of rapamycin complex 2; S6K = S6 kinase; FOXO1 = forkhead box protein O1; STAT3 = signal transducer and transcription 3; POMC = proopiomelanocortin; SOCS3 = suppressor of cytokine signaling 3; NPY = neuropeptide Y.

hypothalamus (Piekarski et al., 2018). A summary of the leptin signaling pathway and observed differences between chickens and mammals has been summarized in Fig. 3.

Several isoforms of the leptin receptor (chLEPR) exist in chickens. The long isoform of the leptin receptor (chLEPRb) contains JAK2 and signal transducer and activator of transcription 3 (STAT3) binding motifs and three conserved mammalian tyrosine

residues (Tyr-986, Tyr-1079, and Tyr-1141) associated with intracellular domain phosphorylation. The short isoform of the chLEPR lacks the STAT3 binding motif and contains only the JAK2 binding motif. Only chLEPRb is able to activate the JAK2/STAT3 pathway. It should be noted that in chickens, there is a lack of expression of the short forms of the chLEPR in the brain, while the long isoform demonstrates a high level of expression, including in the




Fig. 3. Detailed leptin signaling pathway in the hypothalamus of chickens (adapted from “Cytokine Signaling through the JAK-STAT Pathway”, by BioRender.com (2023). Retrieved from <https://app.biorender.com/biorender-templates>). The mechanism of leptin action in chickens may differ from what is known in mammals. In SOCS3-mediated inhibition of leptin signaling, chicken SOCS3 probably does not interact with p-Tyr986 in the intracellular domain of chicken LEPR, but directly binds to JAK2. In contrast to mammals, leptin is more likely to affect feeding behavior of chickens through the expression of the orexigenic neuropeptide NPY (but not AgRP), without affecting the expression of anorexigenic POMC. Note: The pattern of a chicken indicates differences in signaling pathway compared to mammals. LEPR = leptin receptor; JAK2 = Janus tyrosine kinase 2; STAT3 = signal transducer and transcription 3; SOCS3 = suppressor of cytokine signaling 3; NPY = neuropeptide Y.

hypothalamus (Liu et al., 2007). Central leptin injection promoted the expression of chLEPRb in the hypothalamus (Piekariski et al., 2018).

In mammals, binding of leptin to the long isoform of the leptin receptor (LEPRb) leads to dimerisation of the receptor subunits. As a result, JAK2 associated with the intracellular domains of receptors is activated through autophosphorylation due to their proximity to each other (Mengie Ayele et al., 2022). Activated JAK2 stimulates the phosphorylation of three tyrosine residues of the LEPRb intracellular domain (Tyr 985, Tyr1077, and Tyr1138) to create binding sites for proteins. It further enables STAT3 to bind to the receptor at phosphorylated Tyr1138. Then STAT3 is phosphorylated by JAK2 (Liu et al., 2021). ChLEPR was revealed to activate the JAK2/STAT3 signaling pathway in vitro. Stimulation with leptin resulted in STAT3 phosphorylation via chLEPR and JAK2 (Adachi et al., 2008). This proves that vertebrates share a similar leptin signaling pathway.

The phosphorylation of STAT3 promotes its dimerisation and translocation to the nucleus, where it acts as a transcriptional regulator of genes, including suppressor of cytokine signaling 3 (SOCS3) and neuropeptides (POMC, AgRP, and NPY) (Banks et al., 2000; Kwon et al., 2016). In chickens, leptin induced STAT3 phosphorylation and its subsequent translocation to the nucleus in COS-7 cells expressing chLEPR (Adachi et al., 2012). SOCS3 acts as a feedback inhibitor of the JAK2/STAT3 signaling by interacting with LEPR or JAK2, thereby blocking STAT3 activation (Bjørnbæk et al., 2000). Like in mammals, SOCS3 in chickens was demonstrated to be a feedback inhibitor of leptin signaling. This mechanism, however, might be a little different from that found in mammals. Chicken SOCS3 inhibits leptin signaling by binding directly to JAK2, then blocking phosphorylation and subsequent activation of STAT3. SOCS3 may not interact with phospho-Tyr986 in the intracellular

domain of chLEPR for leptin signaling inhibition (Adachi et al., 2013).

The transcription factor STAT3 binds to the promoters of genes that encode anorexigenic neuropeptides (POMC) and anorexigenic neuropeptides (AgRP and NPY). STAT3 acts as an activator of POMC expression and promotes the down-regulation of AgRP and NPY expression, thus reducing feed intake and inducing energy expenditure in mammals (Liu et al., 2021).

However, the mechanism of neuropeptide-mediated action of leptin in chickens may differ from that established in mammals. ICT administration of recombinant chicken leptin decreased the hypothalamic expression of the orexigenic neuropeptide NPY in broilers. However, no changes in AgRP and POMC expression were observed (Dridi et al., 2005). This may suggest that, at least in broiler chickens, leptin preferentially acts through orexigenic neuropeptides (NPY, but not AgRP) as opposed to anorexigenic pathways (POMC). In particular, after immunization against chLEPR, the hypothalamic expression of AgRP and NPY was up-regulated, whereas the expression of POMC was significantly down-regulated (Lei et al., 2015).

5.3. Ghrelin

In mammals one of the crucial peptides involved in controlling appetite and energy homeostasis is ghrelin. Ghrelin also has a stimulating effect on growth hormone (GH) secretion (Kojima et al., 1999). In young chicks, the chicken ghrelin injection also transiently increased plasma GH concentrations (Kaiya et al., 2002).

In mammals, ghrelin is an orexigenic hormone released predominantly by the gastric mucosa, although it is widely expressed in many different tissues, including the central nervous system, the gastrointestinal tract, and the pituitary gland (Devesa, 2021). In the

case of layers, chicks had ghrelin mRNA at the highest levels in the proventriculus, which was comparable to the gastric fundus, but not in the gizzard, whose function is to mechanically process feed (Kaiya et al., 2002). This was also shown in another study in which the highest expression was in the proventriculus and then in the pancreas, brain, and intestines in broiler chickens (Richards et al., 2006). These results indicate that the major site of ghrelin synthesis in laying chickens and broiler chickens is the same, regardless of their lineage. Ghrelin immunopositive cells were found in the mucosal layer of the proventriculus, gastrointestinal tract, and chicken hypothalamus (Ahmed and Harvey, 2002; Neglia et al., 2004; Wada et al., 2003).

The role of ghrelin in relation to feeding behavior and energy balance in chickens differs from that in mammals. In mammals, ghrelin acts as an appetite stimulating hormone both after central and peripheral ghrelin injection, but in chickens, central ghrelin administration, in contrast, suppresses feed intake (Furuse et al., 2001; Saito et al., 2002a, 2005; Taati et al., 2010). In mammals, ghrelin modulates feeding behavior through the growth hormone secretagogue receptor (GHS-R) in hypothalamic neurons, including the ARC nucleus, a main center for maintaining energy homeostasis. Ghrelin stimulates orexigenic AgRP/NPY-associated neurons and inhibits anorexic POMC neurons in the hypothalamus, increasing feed intake and body mass (Kageyama et al., 2010).

Several studies revealed that central ghrelin injection suppressed feed consumption in both broilers and layer chickens (Furuse et al., 2001; Saito et al., 2002a, 2005; Taati et al., 2010). Surprisingly, the effect of peripheral ghrelin injections on feed intake shows conflicting results between chicken strains. Intravenous injections of chicken ghrelin did not influence feed intake among layer chickens (Kaiya et al., 2007). In contrast, peripheral injections of ghrelin into newly hatched and young broiler chickens suppressed feed intake (Buyse et al., 2009; Geelissen et al., 2006; Ocioń and Pietras, 2011).

Further evidence points to a fundamental difference between the peripheral action of ghrelin in chickens and mammals. Peripheral injection of ghrelin into broiler chickens resulted in increased expression of the key lipogenic enzyme fatty acid synthase (FAS) and its associated transcription factors, sterol regulatory element binding protein-1 (SREBP-1) and peroxisome proliferator-activated receptor γ (PPAR γ) in the diencephalon. These findings imply that the anorectic action of ghrelin is mediated by central fatty acid metabolism. On the other hand, the decreased expression levels of FAS and both transcription factors were significantly observed in the liver. This result suggests that ghrelin has a peripheral antilipogenic effect in chickens (Buyse et al., 2009). Intravenous ghrelin injection was found to be accompanied by a reduction in respiratory quotient in broiler chicks, while heat production was not changed, suggesting a decrease in de novo lipogenic activity (Geelissen et al., 2006). The impact of ghrelin in animals, which encourages an increase in respiratory quotient and the deposition of fat, is contrary to this antilipogenic function (Kaiya et al., 2013).

Ghrelin-dependent signaling pathways control eating behavior. Ghrelin mediates its actions primarily through growth hormone secretagogue receptor1a (GHS-R1a), stimulating the secretion of growth hormone. A chicken ghrelin receptor was discovered in different peripheral tissues, such as the pancreas, proventriculus, and also the brain, possibly suggesting autocrine/paracrine effects (Richards et al., 2006; Tanaka et al., 2003). Ghrelin receptor mRNA was detected in the hypothalamus (Chen et al., 2007; Sirotkin et al., 2013; Song et al., 2018). However, ghrelin immunoreactivity was present in the chicken hypothalamus, its presence was not found in the IN nucleus (Ahmed and Harvey, 2002).

Feed restriction was found to be able to increase ghrelin and *GHS-R1a* expression in the hypothalamus of layer hens, but the administration of ghrelin only resulted in a significant increase in *GHS-R1a* mRNA levels (Sirotkin et al., 2013). However, there was no significant effect of feeding restriction and refeeding on the hypothalamic expression of ghrelin and GHS-R1a in e broiler chickens (Chen et al., 2007).

Ghrelin transmits signals by binding to GHS-R1a and raising intracellular calcium levels. Chicken ghrelin was identified to elevate the intracellular calcium ion concentration in chicken cells (Tachibana et al., 2011). Ghrelin regulates feeding behavior in mammals through the AMPK signaling pathway. The interaction between ghrelin and AMPK was exerted through an increase in intracellular calcium levels and subsequent activation of CaMKK β which in turn phosphorylated and activated AMPK (Andrews, 2011). In the case of chickens, central injection of ghrelin significantly inhibited AMPK subunits gene expression and phosphorylation of catalytic AMPK subunits in the hypothalamus. An inhibitory effect of ghrelin on the expression of CaMKK β in chickens with low body weight but not high body weight chicks was also observed (Xu et al., 2011). Therefore, it has been proposed that AMPK signaling in the hypothalamus is responsible for the anorexigenic actions of ghrelin.

In mammals, ghrelin was shown to cause higher calcium levels via AMPK-mediated signaling that led to activation of ARC NPY neurons (Kohno et al., 2008). However, ghrelin administration did not affect hypothalamic NPY mRNA in neonatal layers. In addition, co-injection of ghrelin with NPY prevented the rise in feed intake that NPY causes (Saito et al., 2005). Since ghrelin does not activate NPY neurons in the hypothalamus, it can be assumed that there is no orexigenic effect in chickens. Instead, it has been suggested that the inhibitory effect of ghrelin is mediated by the corticotropin-releasing hormone system, rather than through AgRP/NPY neurons. Ghrelin ICV administration activates the hypothalamic–pituitary–adrenal axis, resulting in higher plasma corticosterone levels (Saito et al., 2005). Furthermore, in support of this hypothesis, it was observed that vocalization, which is characteristic of hyperactivity behavior in chickens, significantly increased after ICV ghrelin injection. Here-with, similar behavior was also observed after injection of CRH, which in turn plays an important role in behavioral responses to stressors and in activation of the HPA axis (Saito et al., 2002b).

6. Conclusions and perspectives

Comparative analysis of the molecular mechanisms regulating feed intake has demonstrated that the majority of components and their interactions that orchestrate such complex biological processes in chickens are quite similar to their counterparts in mammals. In general, it can be suggested that the regulation of eating behavior is based on the integration of hormonal signals and nutritional status by the hypothalamus, which forms the state of satiety or hunger. The AMPK/mTOR signaling pathway, which is crucial to maintaining mammalian energy balance, is involved in the regulation of feeding behavior in chickens as well. However, there are conflicting effects of hormones on the regulation of feed intake in fast- and slow-growing chicken breeds. This is also confirmed by the heterogeneous results in the data on the expression of hypothalamic orexigenic and anorexigenic neuropeptides after hormone injection or feeding restriction. It seems that these differences are related to age, breed, period of feed restriction, or source of the hormone used for the injection. However, further systems studies of the signaling pathways involved in feed intake are required, with a focus on the role of hypothalamic neuropeptides in the formation of eating behavior. Moreover, the complex interrelationships between AMPK/mTOR and hormone-

mediated signaling pathways with downstream regulation of neuropeptide expression cause the unintuitive dynamic behavior of the biological system. Therefore, an application of the mathematical modeling approach, including the development of detailed mechanistic and modular, spatially distributed models is pivotal for further investigation of the molecular mechanisms and their impact on feed intake and energy balance in chickens.

Availability of data and materials

All conceptual diagrams in Systems Biology Graphical Notation (SBGN) standard reflecting signaling pathways and molecular mechanisms in Figs. 1–3 are available as a GitLab project at https://gitlab.sirius-web.org/collaboration/Chicken/Feed_intake. These diagrams can be considered as a growth point for further model development of a certain biological system regulating feed intake in chickens.

Author contributions

Anastasiia R. Volyanskaya: Conceptualization, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. **Ilya R. Akberdin:** Conceptualization, Writing - Original Draft, Writing - Review & Editing, Supervision. **Mikhail A. Kulyashov:** Data curation. **Ivan S. Yevshin:** Data curation. **Michael N. Romanov:** Data curation, Writing - Original Draft. **Elena I. Shagimardanova:** Data curation, Project administration. **Oleg A. Gusev:** Conceptualization, Writing - Review & Editing, Supervision, Project administration, Funding acquisition. **Fedor A. Kolpakov:** Conceptualization, Supervision, Project administration, Funding acquisition.

Declaration of competing interest

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, and there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the content of this paper.

Acknowledgement

This work was supported by the Ministry of Science and Higher Education of the Russian Federation (Grant No. 075–15–2021–1344).

References

Adachi H, Murase D, Atomura S, Ohkubo T. Detection of leptin activity in living cells expressing chicken leptin receptor and STAT3. *J Poult Sci* 2012;49:46–50. <https://doi.org/10.2141/jpsa.011085>.

Adachi H, Murase D, Ohkubo T. Inhibitory mechanism of signal transduction through chicken leptin receptor by suppressor of cytokine signaling 3 (SOCS3). *J Poult Sci* 2013;50:262–9. <https://doi.org/10.2141/jpsa.0120166>.

Adachi H, Takemoto Y, Bungo T, Ohkubo T. Chicken leptin receptor is functional in activating JAK–STAT pathway in vitro. *J Endocrinol* 2008;197:335–42. <https://doi.org/10.1677/JOE-08-0098>.

Adeli A, Zendejdel M, Babapour V, Panahi N. Interaction between leptin and glutamatergic system on food intake regulation in neonatal chicken: role of NMDA and AMPA receptors. *Int J Neurosci* 2020;130:713–21. <https://doi.org/10.1080/00207454.2019.1702983>.

Ahmed S, Harvey S. Ghrelin: a hypothalamic GH-releasing factor in domestic fowl (*Gallus domesticus*). *J Endocrinol* 2002;172:117–25. <https://doi.org/10.1677/joe.0.1720117>.

Ando R, Kawakami S, Bungo T, Ohgushi A, Takagi T, Denbow DM, et al. Feeding responses to several neuropeptide Y receptor agonists in the neonatal chick. *Eur J Pharmacol* 2001;427:53–9. [https://doi.org/10.1016/S0014-2999\(01\)01201-8](https://doi.org/10.1016/S0014-2999(01)01201-8).

Andrews ZB. Central mechanisms involved in the orexigenic actions of ghrelin. *Peptides* 2011;32:2248–55. <https://doi.org/10.1016/j.peptides.2011.05.014>.

Banks AS, Davis SM, Bates SH, Myers MG. Activation of downstream signals by the long form of the leptin receptor. *J Biol Chem* 2000;275:14563–72. <https://doi.org/10.1074/jbc.275.19.14563>.

Barrios-Correa AA, Estrada JA, Contreras I. Leptin signaling in the control of metabolism and appetite: lessons from animal models. *J Mol Neurosci* 2018;66:390–402. <https://doi.org/10.1007/s12031-018-1185-0>.

Bjørbaek C, Lavery HJ, Bates SH, Olson RK, Davis SM, Flier JS, et al. SOCS3 mediates feedback inhibition of the leptin receptor via Tyr985. *J Biol Chem* 2000;275:40649–57. <https://doi.org/10.1074/jbc.M007577200>.

Blouet C, Ono H, Schwartz GJ. Mediobasal hypothalamic p70 S6 kinase 1 modulates the control of energy homeostasis. *Cell Metabol* 2008;8:459–67. <https://doi.org/10.1016/j.cmet.2008.10.004>.

Bornelöv S, Seroussi E, Yosefi S, Benjamini S, Miyara S, Ruzal M, et al. Comparative omics and feeding manipulations in chicken indicate a shift of the endocrine role of visceral fat towards reproduction. *BMC Genom* 2018;295:19. <https://doi.org/10.1186/s12864-018-4675-0>.

Boswell T. Regulation of energy balance in birds by the neuroendocrine hypothalamus. *J Poult Sci* 2005;42:161–81. <https://doi.org/10.2141/jpsa.42.161>.

Boswell T, Dunn IC. Regulation of agouti-related protein and pro-opiomelanocortin gene expression in the avian arcuate nucleus. *Front Endocrinol* 2017;8. <https://doi.org/10.3389/fendo.2017.00075>.

Boswell T, Dunn IC. Regulation of the avian central melanocortin system and the role of leptin. *Gen Comp Endocrinol* 2015;221:278–83. <https://doi.org/10.1016/j.ygcen.2014.12.009>.

Boswell T, Li Q, Takeuchi S. Neurons expressing neuropeptide Y mRNA in the infundibular hypothalamus of Japanese quail are activated by fasting and co-express agouti-related protein mRNA. *Mol Brain Res* 2002;100:31–42. [https://doi.org/10.1016/S0169-328X\(02\)00145-6](https://doi.org/10.1016/S0169-328X(02)00145-6).

Boucher J, Kleinfelders A, Kahn CR. Insulin receptor signaling in normal and insulin-resistant states. *Cold Spring Harbor Perspect Biol* 2014;6:a009191. <https://doi.org/10.1101/cshperspect.a009191>.

Buyse J, Janssen S, Geelissen S, Swennen Q, Kaiya H, Darras VM, et al. Ghrelin modulates fatty acid synthase and related transcription factor mRNA levels in a tissue-specific manner in neonatal broiler chicks. *Peptides* 2009;30:1342–7. <https://doi.org/10.1016/j.peptides.2009.04.015>.

Cammisotto P, Bendani M. A review on gastric leptin: the exocrine secretion of a gastric hormone. *Anat Cell Biol* 2012;1:45. <https://doi.org/10.5115/acb.2012.45.1.1>.

Cassy S, Picard M, Crochet S, Derouet M, Keisler DH, Taouis M. Peripheral leptin effect on food intake in young chickens is influenced by age and strain. *Domest Anim Endocrinol* 2004;27:51–61. <https://doi.org/10.1016/j.domaniend.2004.01.004>.

Chen L-L, Jiang Q-Y, Zhu X-T, Shu G, Bin Y-F, Wang X-Q, et al. Ghrelin ligand-receptor mRNA expression in hypothalamus, proventriculus and liver of chicken (*Gallus gallus domesticus*): studies on ontogeny and feeding condition. *Comp Biochem Physiol Mol Integr Physiol* 2007;147:893–902. <https://doi.org/10.1016/j.cbpa.2007.02.017>.

Cline MA, Smith ML. Central α -melanocyte stimulating hormone attenuates behavioral effects of neuropeptide Y in chicks. *Physiol Behav* 2007;91:588–92. <https://doi.org/10.1016/j.physbeh.2007.03.021>.

Cota D, Proulx K, Smith KAB, Kozma SC, Thomas G, Woods SC, et al. Hypothalamic mTOR signaling regulates food intake. *Science* 2006;312:927–30. <https://doi.org/10.1126/science.1124147>.

Dagon Y, Hur E, Zheng B, Wellenstein K, Cantley LC, Kahn BB. p70S6 kinase phosphorylates AMPK on serine 491 to mediate leptin's effect on food intake. *Cell Metabol* 2012;16:104–12. <https://doi.org/10.1016/j.cmet.2012.05.010>.

Denbow DM, Meade S, Robertson A, McMurtry JP, Richards M, Ashwell C. Leptin-induced decrease in food intake in chickens. *Physiol Behav* 2000;69:359–62. [https://doi.org/10.1016/S0031-9384\(99\)00258-9](https://doi.org/10.1016/S0031-9384(99)00258-9).

Denbow DM, Snapir N, Furuse M. Inhibition of food intake by CRF in chickens. *Physiol Behav* 1999;66:645–9. [https://doi.org/10.1016/S0031-9384\(98\)00340-0](https://doi.org/10.1016/S0031-9384(98)00340-0).

Denbow DM, Cline MA. Food intake regulation. In: *Sturkie's avian physiology*. Elsevier; 2015. p. 469–85. <https://doi.org/10.1016/B978-0-12-407160-5.00021-X>.

Devesa J. The complex world of regulation of pituitary growth hormone secretion: the role of ghrelin, klotho, and nesfatins in it. *Front Endocrinol* 2021;12:636403. <https://doi.org/10.3389/fendo.2021.636403>.

Dibble CC, Cantley LC. Regulation of mTORC1 by PI3K signaling. *Trends Cell Biol* 2015;25:545–55. <https://doi.org/10.1016/j.tcb.2015.06.002>.

Dridi S, Swennen Q, Decuypere E, Buyse J. Mode of leptin action in chicken hypothalamus. *Brain Res* 2005;1047:214–23. <https://doi.org/10.1016/j.brainres.2005.04.034>.

Duchêne S, Métayer S, Audouin E, Bigot K, Dupont J, Tesseraud S. Refeeding and insulin activate the AKT/p70S6 kinase pathway without affecting IRS1 tyrosine phosphorylation in chicken muscle. *Domest Anim Endocrinol* 2008;34:1–13. <https://doi.org/10.1016/j.domaniend.2006.09.002>.

Dunn IC, Wilson PW, Smulders TV, Sandilands V, D'Eath RB, Boswell T. Hypothalamic agouti-related protein expression is affected by both acute and chronic experience of food restriction and Re-feeding in chickens. *J Neuroendocrinol* 2013;25:920–8. <https://doi.org/10.1111/jne.12088>.

Dupont J, Métayer-Coustard S, Ji B, Ramé C, Gespach C, Voy B, et al. Characterization of major elements of insulin signaling cascade in chicken adipose tissue: apparent insulin refractoriness. *Gen Comp Endocrinol* 2012;176:86–93. <https://doi.org/10.1016/j.ygcen.2011.12.030>.

- Everaert N, Decuyper E, Buyse J. 3: feed intake and regulation. In: Hendriks WH, Verstegen MWA, Babinszky L, editors. Poultry and pig nutrition. Brill | Wageningen Academic; 2019. p. 59–75. https://doi.org/10.3920/978-90-8686-884-1_3.
- Engelman JA, Luo J, Cantley LC. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet* 2006;7:606–19. <https://doi.org/10.1038/nrg1879>.
- Fang X-L, Zhu X-T, Chen S-F, Zhang Z-Q, Zeng Q-J, Deng L, et al. Differential gene expression pattern in hypothalamus of chickens during fasting-induced metabolic reprogramming: functions of glucose and lipid metabolism in the feed intake of chickens. *Poultry Sci* 2014;93:2841–54. <https://doi.org/10.3382/ps.2014-04047>.
- Farkasová H, Hron T, Pačes J, Pajer P, Elleder D. Identification of a GC-rich leptin gene in chicken. *Agri Gene* 2016;1:88–92. <https://doi.org/10.1016/j.aggene.2016.04.001>.
- Friedman JM, Halaas JL. Leptin and the regulation of body weight in mammals. *Nature* 1998;395:763–70. <https://doi.org/10.1038/27376>.
- Friedman J. 20 years of leptin: leptin at 20: an overview. *J Endocrinol* 2014;223:T1–8. <https://doi.org/10.1530/JOE-14-0405>.
- Friedman-Einat M, Cogburn LA, Yosefi S, Hen G, Shinder D, Shirak A, et al. Discovery and characterization of the first genuine avian leptin gene in the Rock dove (*Columba livia*). *Endocrinology* 2014;155:3376–84. <https://doi.org/10.1210/en.2014-1273>.
- Friedman-Einat M, Seroussi E. Avian leptin: bird's-eye view of the evolution of vertebrate energy-balance control. *Trends Endocrinol Metab* 2019;30:819–32. <https://doi.org/10.1016/j.tem.2019.07.007>.
- Furuse M, Tachibana T, Ohgushi A, Ando R, Yoshimatsu T, Denbow DM. Intra-cerebroventricular injection of ghrelin and growth hormone releasing factor inhibits food intake in neonatal chicks. *Neurosci Lett* 2001;301:123–6. [https://doi.org/10.1016/S0304-3940\(01\)01621-4](https://doi.org/10.1016/S0304-3940(01)01621-4).
- Gali Ramamoorthy T, Begum G, Harno E, White A. Developmental programming of hypothalamic neuronal circuits: impact on energy balance control. *Front Neurosci* 2015;9. <https://doi.org/10.3389/fnins.2015.00126>.
- Geelissen SME, Swennen Q, Geyten SVD, Kühn ER, Kaiya H, Kangawa K, et al. Peripheral ghrelin reduces food intake and respiratory quotient in chicken. *Domest Anim Endocrinol* 2006;30:108–16. <https://doi.org/10.1016/j.domaniend.2005.06.005>.
- Gerets HHJ, Peeters K, Arckens L, Vandesande F, Berghman LR. Sequence and distribution of pro-opiomelanocortin in the pituitary and the brain of the chicken (*Gallus gallus*). *J Comp Neurol* 2000;417:250–62. [https://doi.org/10.1002/\(SICI\)1096-9861\(20000207\)417:2<250::AID-CNE9>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1096-9861(20000207)417:2<250::AID-CNE9>3.0.CO;2-Z).
- Greene ES, Abdelli N, Dridi JS, Dridi S. Avian neuropeptide Y: beyond feed intake regulation. *Vet Sci* 2022;171:9. <https://doi.org/10.3390/vetsci9040171>.
- Gual P, Le Marchand-Brustel Y, Tanti J-F. Positive and negative regulation of insulin signaling through IRS-1 phosphorylation. *Biochimie* 2005;87:99–109. <https://doi.org/10.1016/j.biochi.2004.10.019>.
- Hardie DG. AMP-activated protein kinase: maintaining energy homeostasis at the cellular and whole-body levels. *Annu Rev Nutr* 2014;34:31–55. <https://doi.org/10.1146/annurev-nutr-071812-161148>.
- Hay N, Sonenberg N. Upstream and downstream of mTOR. *Genes Dev* 2004;18:1926–45. <https://doi.org/10.1101/gad.1212704>.
- Hen G, Yosefi S, Simchaev V, Shinder D, Hruby VJ, Friedman-Einat M. The melanocortin circuit in obese and lean strains of chicks. *J Endocrinol* 2006;190:527–35. <https://doi.org/10.1677/joe.1.06783>.
- Higgins SC, Gueorguiev M, Korbonits M. Ghrelin, the peripheral hunger hormone. *Ann Med* 2007;39:116–36. <https://doi.org/10.1080/07853890601149179>.
- Higgins SE, Ellestad LE, Trakooljil N, McCarthy F, Saliba J, Cogburn LA, et al. Transcriptional and pathway analysis in the hypothalamus of newly hatched chicks during fasting and delayed feeding. *BMC Genom* 2010;162:11. <https://doi.org/10.1186/1471-2164-11-162>.
- Honda K. Peripheral regulation of food intake in chickens: adiposity signals, satiety signals and others. *World's Poultry Sci J* 2021;77:301–12. <https://doi.org/10.1080/00439339.2021.1898296>.
- Honda K, Kamisoyama H, Saneyasu T, Sugahara K, Hasegawa S. Central administration of insulin suppresses food intake in chicks. *Neurosci Lett* 2007;423:153–7. <https://doi.org/10.1016/j.neulet.2007.07.004>.
- Honda K, Saneyasu T, Aoki K, Shimatani T, Yamaguchi T, Kamisoyama H. Correlation analysis of hypothalamic mRNA levels of appetite regulatory neuropeptides and several metabolic parameters in 28-d-old layer chickens: hypothalamic Neuropeptides in Chickens. *Anim Sci J* 2015;86:517–22. <https://doi.org/10.1111/asj.12320>.
- Honda K, Saneyasu T, Hasegawa S, Kamisoyama H. A comparative study of the central effects of melanocortin peptides on food intake in broiler and layer chicks. *Peptides* 2012;37:13–7. <https://doi.org/10.1016/j.peptides.2012.06.015>.
- Hu F, Xu Y, Liu F. Hypothalamic roles of mTOR complex I: integration of nutrient and hormone signals to regulate energy homeostasis. *Am J Physiol Endocrinol Metab* 2016;310:E994–1002. <https://doi.org/10.1152/ajpendo.00121.2016>.
- Hu X, Kong L, Xiao C, Zhu Q, Song Z. The AMPK-mTOR signaling pathway is involved in regulation of food intake in the hypothalamus of stressed chickens. *Comp Biochem Physiol Mol Integr Physiol* 2021;110979:258. <https://doi.org/10.1016/j.cbpa.2021.110979>.
- Hu X, Wang Y, Sheikahmadi A, Li X, Buyse J, Lin H, et al. Effects of dietary energy level on appetite and central adenosine monophosphate-activated protein kinase (AMPK) in broilers. *J Anim Sci* 2019;97:4488–95. <https://doi.org/10.1093/jas/skz2312>.
- Huang G, Li J, Wang H, Lan X, Wang Y. Discovery of a novel functional leptin protein (LEP) in zebra finches: evidence for the existence of an authentic avian leptin gene predominantly expressed in the brain and pituitary. *Endocrinology* 2014;155:3385–96. <https://doi.org/10.1210/en.2014-1084>.
- Inoki K, Li Y, Zhu T, Wu J, Guan K-L. TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. *Nat Cell Biol* 2002;4:648–57. <https://doi.org/10.1038/ncb839>.
- Jewell JL, Guan KL. Nutrient signaling to mTOR and cell growth. *Trends Biochem Sci* 2013;38(5):233–42. <https://doi.org/10.1016/j.tibs.2013.01.004>.
- Kageyama H, Takenoya F, Shiba K, Shioda S. Neuronal circuits involving ghrelin in the hypothalamus-mediated regulation of feeding. *Neuropeptides* 2010;44:133–8. <https://doi.org/10.1016/j.npep.2009.11.010>.
- Kaiya H, Kangawa K, Miyazato M. Update on ghrelin biology in birds. *Gen Comp Endocrinol* 2013;190:170–5. <https://doi.org/10.1016/j.ygcen.2013.04.014>.
- Kaiya H, Saito E-S, Tachibana T, Furuse M, Kangawa K. Changes in ghrelin levels of plasma and proventriculus and ghrelin mRNA of proventriculus in fasted and re-fed layer chicks. *Domest Anim Endocrinol* 2007;32:247–59. <https://doi.org/10.1016/j.domaniend.2006.03.006>.
- Kaiya H, Van Der Geyten S, Kojima M, Hosoda H, Kitajima Y, Matsumoto M, et al. Chicken ghrelin: purification, cDNA cloning, and biological activity. *Endocrinology* 2002;143:3454–63. <https://doi.org/10.1210/en.2002-220255>.
- Kameda Y, Miura M, Nishimaki T. Localization of neuropeptide Y mRNA and peptide in the chicken hypothalamus and their alterations after food deprivation, dehydration, and castration. *J Comp Neurol* 2001;436:376–88. <https://doi.org/10.1002/cne.1074>.
- Kawakami S, Bungo T, Ando R, Ohgushi A, Shimojo M, Masuda Y, et al. Central administration of α -melanocyte stimulating hormone inhibits fasting- and neuropeptide Y-induced feeding in neonatal chicks. *Eur J Pharmacol* 2000;398:361–4. [https://doi.org/10.1016/S0014-2999\(00\)00344-7](https://doi.org/10.1016/S0014-2999(00)00344-7).
- Kehinde A, Muyideen I, Monsurat G. Effects of the bioactive amino acids leucine and tryptophan on feed intake in layer chicks. *Int J Fauna Biol Stud* 2022;9:13–7. <https://doi.org/10.22271/23940522.2022.v9.i6a.937>.
- Kewan A, Saneyasu T, Kamisoyama H, Honda K. Effects of fasting and re-feeding on the expression of CCK, PYY, hypothalamic neuropeptides, and IGF-related genes in layer and broiler chicks. *Comp Biochem Physiol Mol Integr Physiol* 2021;110940:257. <https://doi.org/10.1016/j.cbpa.2021.110940>.
- Kim M-S, Pak YK, Jang P-G, Namkoong C, Choi Y-S, Won J-C, et al. Role of hypothalamic Foxo 1 in the regulation of food intake and energy homeostasis. *Nat Neurosci* 2006;9:901–6. <https://doi.org/10.1038/nn1731>.
- Kodani N, Nakae J. Tissue-specific metabolic regulation of FOXO-binding protein: FOXO does not act alone. *Cells* 2020;702:9. <https://doi.org/10.3390/cells9030702>.
- Kohno D, Sone H, Minokoshi Y, Yada T. Ghrelin raises [Ca²⁺]_i via AMPK in hypothalamic arcuate nucleus NPY neurons. *Biochem Biophys Res Commun* 2008;366:388–92. <https://doi.org/10.1016/j.bbrc.2007.11.166>.
- Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 1999;402:656–60. <https://doi.org/10.1038/45230>.
- Kuenzel WJ, Beck MM, Teruyama R. Neural sites and pathways regulating food intake in birds: a comparative analysis to mammalian systems. *J Exp Zool* 1999;283:348–64. [https://doi.org/10.1002/\(SICI\)1097-010X\(19990301\)283:4/5<348::AID-JEZ5>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1097-010X(19990301)283:4/5<348::AID-JEZ5>3.0.CO;2-5).
- Kuo AY, Cline MA, Werner E, Siegel PB, Denbow DM. Leptin effects on food and water intake in lines of chickens selected for high or low body weight. *Physiol Behav* 2005;84:459–64. <https://doi.org/10.1016/j.physbeh.2005.01.014>.
- Kwon O, Kim KW, Kim M-S. Leptin signalling pathways in hypothalamic neurons. *Cell Mol Life Sci* 2016;73:1457–77. <https://doi.org/10.1007/s00018-016-2133-1>.
- Lei MM, Wu SQ, Shao XB, Li XW, Chen Z, Ying SJ, et al. Creating leptin-like bio-functions by active immunization against chicken leptin receptor in growing chickens. *Domest Anim Endocrinol* 2015;50:55–64. <https://doi.org/10.1016/j.domaniend.2014.09.001>.
- Liu H, Du T, Li C, Yang G. STAT3 phosphorylation in central leptin resistance. *Nutr Metab* 2021;39:18. <https://doi.org/10.1186/s12986-021-00569-w>.
- Liu L, Xu S, Wang X, Jiao H, Lin H. Peripheral insulin doesn't alter appetite of broiler chicks. *Asian-Australas J Anim Sci* 2016;29:1294–9. <https://doi.org/10.5713/ajas.15.0674>.
- Liu L, Zhu L. Effect of 24 h fasting on gene expression of AMPK, appetite regulation peptides and lipometabolism related factors in the hypothalamus of broiler chicks. *Asian-Australas J Anim Sci* 2012;25:1300–8. <https://doi.org/10.5713/ajas.2012.12153>.
- Liu X, Dunn IC, Sharp PJ, Boswell T. Molecular cloning and tissue distribution of a short form chicken leptin receptor mRNA. *Domest Anim Endocrinol* 2007;32:155–66. <https://doi.org/10.1016/j.domaniend.2006.02.001>.
- Manning BD, Cantley LC. AKT/PKB signaling: navigating downstream. *Cell* 2007;129:1261–74. <https://doi.org/10.1016/j.cell.2007.06.009>.
- Mengie Ayele T, Tilahun Muche Z, Behaile Teklemariam A, Bogale A, Chekol Abebe E. Role of JAK2/STAT3 signaling pathway in the tumorigenesis, chemotherapy resistance, and treatment of solid tumors: a systemic review. *J Inflamm Res* 2022;ume 15:1349–64. <https://doi.org/10.2147/JIR.S353489>.
- Murugesan S, Nidamanuri AL. Role of leptin and ghrelin in regulation of physiological functions of chicken. *World's Poultry Sci J* 2022;78:1021–36. <https://doi.org/10.1080/00439339.2022.2119917>.

- Neglia S, Arcamone N, Esposito V, Gargiulo G. Ghrelin in the gastroenteric tract of birds: immunoreactivity expression. *Vet Res Commun* 2004;28:213–5. <https://doi.org/10.1023/B:VERC.0000045409.48367.c0>.
- Octoń E, Pietras M. Peripheral ghrelin inhibits feed intake through hypothalamo-pituitary-adrenal axis-dependent mechanism in chicken. *J Anim Feed Sci* 2011;20:118–30. <https://doi.org/10.22358/jafs/66163/2011>.
- Ottensmeyer FP, Beniac DR, Luo RZ-T, Yip CC. Mechanism of transmembrane signaling: insulin binding and the insulin receptor. *Biochemistry* 2000;39:12103–12. <https://doi.org/10.1021/bi001592t>.
- Piekarski A, Nagarajan G, Ishola P, Flees J, Greene ES, Kuenzel WJ, et al. AMP-activated protein kinase mediates the effect of leptin on avian autophagy in a tissue-specific manner. *Front Physiol* 2018;541:9. <https://doi.org/10.3389/fphys.2018.00541>.
- Proszkowiec-Weglarczyk M, Richards MP, Ramachandran R, McMurtry JP. Characterization of the AMP-activated protein kinase pathway in chickens. *Comp Biochem Physiol B Biochem Mol Biol* 2006;143:92–106. <https://doi.org/10.1016/j.cbpb.2005.10.009>.
- Rau AR, Hentges ST. GABAergic inputs to POMC neurons originating from the dorsomedial hypothalamus are regulated by energy state. *J Neurosci* 2019;39:6449–59. <https://doi.org/10.1523/JNEUROSCI.3193-18.2019>.
- Resnyk CW, Chen C, Huang H, Wu CH, Simon J, Le Bihan-Duval E, et al. RNA-seq analysis of abdominal fat in genetically fat and lean chickens highlights a divergence in expression of genes controlling adiposity, hemostasis, and lipid metabolism. *PLoS One* 2015;10:e0139549. <https://doi.org/10.1371/journal.pone.0139549>.
- Richards MP, Poch SM, McMurtry JP. Characterization of Turkey and chicken ghrelin genes, and regulation of ghrelin and ghrelin receptor mRNA levels in broiler chickens. *Gen Comp Endocrinol* 2006;145:298–310. <https://doi.org/10.1016/j.ygcen.2005.09.013>.
- Richards MP, Proszkowiec-Weglarczyk M. Mechanisms regulating feed intake, energy expenditure, and body weight in poultry. *Poultry Sci* 2007;86:1478–90. <https://doi.org/10.1093/ps/86.7.1478>.
- Ronnett GV, Ramamurthy S, Kleman AM, Landree LE, Aja S. AMPK in the brain: its roles in energy balance and neuroprotection. *J Neurochem* 2009;109:17–23. <https://doi.org/10.1111/j.1471-4159.2009.05916.x>.
- Saito E, Kaiya H, Takagi T, Yamasaki I, Denbow DM, Kangawa K, et al. Chicken ghrelin and growth hormone-releasing peptide-2 inhibit food intake of neonatal chicks. *Eur J Pharmacol* 2002a;453:75–9. [https://doi.org/10.1016/S0014-2999\(02\)02393-2](https://doi.org/10.1016/S0014-2999(02)02393-2).
- Saito E, Takagi T, Nakanishi T, Sashihara K, Furuse M. Ghrelin activates behavior of neonatal chicks in a short period of post-intracerebroventricular injection. *J Appl Anim Res* 2002b;22:33–41. <https://doi.org/10.1080/09712119.2002.9706376>.
- Saito E-S, Kaiya H, Tachibana T, Tomonaga S, Denbow DM, Kangawa K, et al. Inhibitory effect of ghrelin on food intake is mediated by the corticotropin-releasing factor system in neonatal chicks. *Regul Pept* 2005;125:201–8. <https://doi.org/10.1016/j.regpep.2004.09.003>.
- Sancak Y, Peterson TR, Shaull YD, Lindquist RA, Thoreen CC, Bar-Peled L, et al. The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. *Science* 2008;320:1496–501. <https://doi.org/10.1126/science.1157535>.
- Saneyasu T, Fujita S, Kitashiro A, Fukuzo S, Honda K, Kamisoyama H. Hypothalamic Akt-mediated signaling regulates food intake in chicks. *Neurosci Lett* 2018;670:48–52. <https://doi.org/10.1016/j.neulet.2018.01.032>.
- Saneyasu T, Honda K, Kamisoyama H, Nakayama Y, Ikegami K, Hasegawa S. Alpha-melanocyte stimulating hormone plays an important role in the regulation of food intake by the central melanocortin system in chicks. *Peptides* 2011a;32:996–1000. <https://doi.org/10.1016/j.peptides.2011.03.006>.
- Saneyasu T, Honda K, Kamisoyama H, Ikura A, Nakayama Y, Hasegawa S. Neuropeptide Y effect on food intake in broiler and layer chicks. *Comp Biochem Physiol Mol Integr Physiol* 2011b;159:422–6. <https://doi.org/10.1016/j.cbpa.2011.04.008>.
- Saneyasu T, Nakanishi K, Atsuta H, Ikura A, Kamisoyama H, Hasegawa S, et al. Age-dependent changes in the mRNA levels of neuropeptide Y, proopiomelanocortin, and corticotropin-releasing factor in the hypothalamus in growing broiler chicks. *J Poult Sci* 2013;50:364–9. <https://doi.org/10.2141/jpsa.0120188>.
- Saneyasu T, Fukuzo S, Kitashiro A, Nagata K, Honda K, Kamisoyama H. Central administration of insulin and refeeding lead to the phosphorylation of AKT, but not FOXO1, in the hypothalamus of broiler chicks. *Physiol Behav* 2019;112644:210. <https://doi.org/10.1016/j.physbeh.2019.112644>.
- Seki Y, Sato K, Kono T, Abe H, Akiba Y. Broiler chickens (Ross strain) lack insulin-responsive glucose transporter GLUT4 and have GLUT8 cDNA. *Gen Comp Endocrinol* 2003;133:80–7. [https://doi.org/10.1016/S0016-6480\(03\)00145-X](https://doi.org/10.1016/S0016-6480(03)00145-X).
- Seroussi E, Cinnamon Y, Yosefi S, Genin O, Smith JG, Rafati N, et al. Identification of the long-sought leptin in chicken and duck: expression pattern of the highly GC-rich avian leptin fits an autocrine/paracrine rather than endocrine function. *Endocrinology* 2016;157:737–51. <https://doi.org/10.1210/en.2015-1634>.
- Seroussi E, Knytl M, Pitel F, Elleder D, Krylov V, Leroux S, et al. Avian expression patterns and genomic mapping implicate leptin in digestion and TNF signaling, suggesting that their interacting adipokine role is unique to mammals. *Int J Mol Sci* 2019;4489:20. <https://doi.org/10.3390/ijms20184489>.
- Shiraishi J, Tanizawa H, Fujita M, Kawakami S-I, Bungo T. Localization of hypothalamic insulin receptor in neonatal chicks: evidence for insulinergic system control of feeding behavior. *Neurosci Lett* 2011a;491:177–80. <https://doi.org/10.1016/j.neulet.2011.01.031>.
- Shiraishi J, Yanagita K, Fujita M, Bungo T. Central insulin suppresses feeding behavior via melanocortins in chicks. *Domest Anim Endocrinol* 2008;34:223–8. <https://doi.org/10.1016/j.domaniend.2007.05.002>.
- Shiraishi J, Yanagita K, Fukumori R, Sugino T, Fujita M, Kawakami S-I, et al. Comparisons of insulin related parameters in commercial-type chicks: evidence for insulin resistance in broiler chicks. *Physiol Behav* 2011b;103:233–9. <https://doi.org/10.1016/j.physbeh.2011.02.008>.
- Shiraishi J, Yanagita K, Nishikawa F, Tahara Y, Fujita M, McMurtry JP, et al. A comparison of the anorexic effects of chicken, porcine, human and bovine insulin on the central nervous system of chicks. *J Poult Sci* 2009;46:144–8. <https://doi.org/10.2141/jpsa.46.144>.
- Simon J, Leroith D. Insulin receptors of chicken liver and brain. Characterization of alpha and beta subunit properties. *Eur J Biochem* 1986;158:125–32. <https://doi.org/10.1111/j.1432-1033.1986.tb09729.x>.
- Sims W, Yi J, Cline MA, Gilbert ER. Central injection of a synthetic chicken partial leptin peptide does not affect food intake in chicks. *Neurosci Lett* 2017;656:165–8. <https://doi.org/10.1016/j.neulet.2017.07.038>.
- Sirotkin AV, Pavlova S, Tena-Sempere M, Grossmann R, Jiménez MR, Rodríguez JMC, et al. Food restriction, ghrelin, its antagonist and obestatin control expression of ghrelin and its receptor in chicken hypothalamus and ovary. *Comp Biochem Physiol Mol Integr Physiol* 2013;164:141–53. <https://doi.org/10.1016/j.cbpa.2012.07.010>.
- Smith ML, Prall B, Nandar W, Cline MA. β -Melanocyte-Stimulating hormone potentially reduces appetite via the hypothalamus in chicks: β -MSH potentially reduces appetite. *J Neuroendocrinol* 2007;20:220–6. <https://doi.org/10.1111/j.1365-2826.2007.01639.x>.
- Song X, Jiao H, Zhao J, Wang X, Lin H. Dexamethasone and insulin stimulate ghrelin secretion of broilers in a different way. *Gen Comp Endocrinol* 2018;268:14–21. <https://doi.org/10.1016/j.ygcen.2018.07.009>.
- Song Z, Liu L, Yue Y, Jiao H, Lin H, Sheikahmadi A, et al. Fasting alters protein expression of AMP-activated protein kinase in the hypothalamus of broiler chicks (*Gallus gallus domesticus*). *Gen Comp Endocrinol* 2012;178:546–55. <https://doi.org/10.1016/j.ygcen.2012.06.026>.
- Swennen Q, Decuyper E, Buyse J. Implications of dietary macronutrients for growth and metabolism in broiler chickens. *World's Poult Sci J* 2007;63:541–56. <https://doi.org/10.1017/S0043933907001602>.
- Taati M, Nayebzadeh H, Khosravinia H, Cheraghi J. The role of the histaminergic system on the inhibitory effect of ghrelin on feed intake in broiler chickens. *Iran J Vet Res* 2010;11. <https://doi.org/10.22099/ijvr.2010.173>.
- Tachibana T, Oikawa D, Takahashi H, Boswell T, Furuse M. The anorexic effect of alpha-melanocyte-stimulating hormone is mediated by corticotropin-releasing factor in chicks. *Comp Biochem Physiol Mol Integr Physiol* 2007;147:173–8. <https://doi.org/10.1016/j.cbpa.2006.12.044>.
- Tachibana T, Sato M, Oikawa D, Takahashi H, Boswell T, Furuse M. Intracerebroventricular injection of neuropeptide Y modifies carbohydrate and lipid metabolism in chicks. *Regul Pept* 2006;136:1–8. <https://doi.org/10.1016/j.regpep.2006.04.005>.
- Tachibana T, Sugahara K, Ohgushi A, Ando R, Kawakami S-I, Yoshimatsu T, et al. Intracerebroventricular injection of agouti-related protein attenuates the anorexic effect of alpha-melanocyte stimulating hormone in neonatal chicks. *Neurosci Lett* 2001;305:131–4. [https://doi.org/10.1016/S0304-3940\(01\)01827-4](https://doi.org/10.1016/S0304-3940(01)01827-4).
- Tachibana T, Tanaka M, Kaiya H. Central injection of des-acyl chicken ghrelin does not affect food intake in chicks. *Gen Comp Endocrinol* 2011;171:183–8. <https://doi.org/10.1016/j.ygcen.2011.01.008>.
- Tachibana T, Tsutsui K. Neuropeptide control of feeding behavior in birds and its difference with mammals. *Front Neurosci* 2016;10. <https://doi.org/10.3389/fnins.2016.00485>.
- Tanaka M, Miyazaki T, Yamamoto I, Nakai N, Ohta Y, Tsushima N, et al. Molecular characterization of chicken growth hormone secretagogue receptor gene. *Gen Comp Endocrinol* 2003;134:198–202. [https://doi.org/10.1016/S0016-6480\(03\)00247-8](https://doi.org/10.1016/S0016-6480(03)00247-8).
- Taouis M, Derouet M, Simon J, Taylor SI, Reitman M. P-52: characterisation of insulin signalling in chicken. *Exp Clin Endocrinol Diabetes* 2009;104:118–9. <https://doi.org/10.1055/s-0029-1211595>.
- Van Swieten MMH, Pandit R, Adan RAH, Van Der Plasse G. The neuroanatomical function of leptin in the hypothalamus. *J Chem Neuroanat* 2014;61–62:207–20. <https://doi.org/10.1016/j.jchemneu.2014.05.004>.
- Wada R, Sakata I, Kaiya H, Nakamura K, Hayashi Y, Kangawa K, et al. Existence of ghrelin-immunopositive and -expressing cells in the proventriculus of the hatching and adult chicken. *Regul Pept* 2003;111:123–8. [https://doi.org/10.1016/S0167-0115\(02\)00265-3](https://doi.org/10.1016/S0167-0115(02)00265-3).
- Wang S, Khondowe P, Chen S, Yu J, Shu G, Zhu X, et al. Effects of “Bioactive” amino acids leucine, glutamate, arginine and tryptophan on feed intake and mRNA expression of relative neuropeptides in broiler chicks. *J Anim Sci Biotechnol* 2012;27:3. <https://doi.org/10.1186/2049-1891-3-27>.
- Wang X, Day JR, Vasilatos-Young R. The distribution of neuropeptide Y gene expression in the chicken brain. *Mol Cell Endocrinol* 2001;174:129–36. [https://doi.org/10.1016/S0303-7207\(00\)00436-6](https://doi.org/10.1016/S0303-7207(00)00436-6).
- Woods S, Seeley R. Insulin as an adiposity signal. *Int J Obes* 2001;25:S35–8. <https://doi.org/10.1038/sj.ijo.0801909>.
- Wynne K, Stanley S, McGowan B, Bloom S. Appetite control. *J Endocrinol* 2005;184:291–318. <https://doi.org/10.1677/joe.1.05866>.

- Xu J, Ji J, Yan X-H. Cross-talk between AMPK and mTOR in regulating energy balance. *Crit Rev Food Sci Nutr* 2012;52:373–81. <https://doi.org/10.1080/10408398.2010.500245>.
- Xu P, Siegel PB, Denbow DM. Genetic selection for body weight in chickens has altered responses of the brain's AMPK system to food intake regulation effect of ghrelin, but not obestatin. *Behav Brain Res* 2011;221:216–26. <https://doi.org/10.1016/j.bbr.2011.02.034>.
- Yang G, Lim C-Y, Chao Li, Xiao X, Radda GK, Cai Li, et al. FoxO 1 inhibits leptin regulation of pro-opiomelanocortin promoter activity by blocking STAT3 interaction with specificity protein 1. *J Biol Chem* 2009;284:3719–27. <https://doi.org/10.1074/jbc.M804965200>.
- Yang G, Murashige DS, Humphrey SJ, James DE. A positive feedback loop between Akt and mTORC2 via SIN1 phosphorylation. *Cell Rep* 2015;12:937–43. <https://doi.org/10.1016/j.celrep.2015.07.016>.
- Yousefvand S, Hamidi F, Zendeheel M, Parham A. Survey the effect of insulin on modulating feed intake via NPY receptors in 5-D-old chickens. *Int J Pept Res Therapeut* 2020;26:467–76. <https://doi.org/10.1007/s10989-019-09852-0>.
- Yousefvand S, Hamidi F, Zendeheel M, Parham A. Hypophagic effects of insulin are mediated via NPY 1/NPY 2 receptors in broiler cockerels. *Can J Physiol Pharmacol* 2018;96:1301–7. <https://doi.org/10.1139/cjpp-2018-0470>.
- Zhang J, Li X, Zhou Y, Cui L, Jing Li, Wu C, et al. The interaction of MC3R and MC4R with MRAP2, ACTH, α -MSH and AgRP in chickens. *J Endocrinol* 2017;234:155–74. <https://doi.org/10.1530/JOE-17-0131>.
- Zhang K, Shi Y, Xu P, Huang C, Zhou C, Liu P, et al. Cloning and prokaryotic expression of the chicken liver kinase B1 (LKB1) and its localization in liver, heart and hypothalamus. *Int J Biol Macromol* 2021;169:513–20. <https://doi.org/10.1016/j.ijbiomac.2020.12.195>.