

# Antioxidation and Antiapoptosis Characteristics of Heme Oxygenase-1 Enhance Tumorigenesis of Human Prostate Carcinoma Cells



Kun-Chun Chiang<sup>\*,1</sup>, Ke-Hung Tsui<sup>†,1</sup>, Yu-Hsiang Lin<sup>†,‡</sup>, Chen-Pang Hou<sup>†,‡</sup>, Kang-Shuo Chang<sup>§,||</sup>, Hsin-Han Tsai<sup>§,||</sup>, Yi-Syuan Shin<sup>¶</sup>, Chiu-Chun Chen<sup>¶</sup>, Tsui-Hsia Feng<sup>#</sup> and Horng-Heng Juang<sup>†,§,||</sup>

\*Department of General Surgery, Min-Sheng General Hospital, Tao-Yuan, Taiwan; <sup>†</sup>Department of Urology, Chang Gung Memorial Hospital-Linkou, Kwei-Shan, Tao-Yuan, Taiwan; <sup>‡</sup>Graduate Institute of Clinical Medical Science, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan; <sup>§</sup>Department of Anatomy, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan; <sup>||</sup>Institute of Biomedicine, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan; <sup>¶</sup>Department of Medicine, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan; <sup>#</sup>School of Nursing, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan

## Abstract

Heme oxygenase-1 (HO-1) has antiinflammatory and antioxidant properties and is deemed as a tissue protector. However, effects of HO-1 in prostate cancer remain in controversy. We evaluated the role of HO-1 in prostate carcinoma *in vitro* and *in vivo*.

Overexpression of HO-1 did not affect prostate cell proliferation in the normal condition but enhanced cell proliferation under serum starvation. HO-1 overexpression enhanced cell invasion of PC-3 cells through epithelial–mesenchymal transition (EMT) induction, which was supported by increased Slug, N-cadherin, and vimentin expressions. In the xenograft animal study, HO-1 overexpression enhanced PC-3 cell tumor growth *in vivo*. HO-1 attenuated reactive oxygen species induced by H<sub>2</sub>O<sub>2</sub> or pyocyanin treatment in PC-3 and DU145 cells. HO-1 further reduced PC-3 and DU145 cell apoptosis induced by H<sub>2</sub>O<sub>2</sub> or serum starvation. Our results suggested that HO-1 was able to increase prostate carcinoma cell invasion *in vitro* and tumor growth *in vivo*. The EMT induction and antioxidant and antiapoptotic effects of HO-1 in the prostate carcinoma cells may be responsible for these findings.

*Translational Oncology* (2020) 13, 102–112

Address all correspondence to: Horng-Heng Juang, Department of Urology, Chang Gung Memorial Hospital-Linkou, Kwei-Shan, Tao-Yuan, Taiwan. E-mail: [hhj143@mil.cgu.edu.tw](mailto:hhj143@mil.cgu.edu.tw).

<sup>†</sup>The authors contributed equally to this work.

Received 6 August 2019; Revised 14 October 2019; Accepted 15 October 2019

© 2019 The Authors. Published by Elsevier Inc. on behalf of Neoplasia Press, Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1936-5233/19

<https://doi.org/10.1016/j.tranonc.2019.10.008>

## Introduction

Heme degradation in human is catalyzed by heme oxygenase (HO) to produce carbon monoxide (CO), biliverdin, and iron [1–3]. HO includes two isozymes, which are HO-1 and HO-2, each has distinct expression pattern in the body. It has been shown that HO-2 represents as a constitutive and noninducible protein with mainly expressing in testis and brain [4]. To the contrary, HO-1 is a highly inducible protein with only very low expression in the cell before

numerous stimulations causing oxidative stress, including heavy metals, cytokines, heme, etc. [5] As a result of the antioxidative stress characteristic, HO-1 is deemed as a tissue protector, especially for kidney, heart, and liver [6]. However, this protective effect, while applying in cancer cells, may lead to increased tumor growth due to the antiapoptosis effect derived from the antioxidative stress of HO-1.

It is reported that the emerging role of HO-1 in tumor biology is multifaceted and tissue-specific [7]. Cytoprotective and antiapoptotic properties of HO-1 in tumor by increasing survival of cancer cells and resistance to anticancer drugs or therapies have been found in several cancer types [8,9]. The possible mechanisms by which HO-1 affects apoptosis of cancer cells may be through the modulation of caspase-3, B-cell lymphoma-2, B-cell lymphoma-extra large, vascular endothelial growth factor, and stromal cell-derived factor-1 [10–13]. Furthermore, HO-1 has also been shown to be the downstream gene of some oncogenes [9,11].

Thus, it is not surprised that HO-1 has been implied to induce tumor progression in renal cancer, glioma, lung cancer, etc. [14]. However, high HO-1 expression has also been shown to link with slower cancer growth in breast cancer and non-small-cell lung cancer [15,16]. For prostate cancer, the role of HO-1 is still controversial. HO-1 has been shown to increase tumor growth and metastasis in prostate cancer [17]; however, Gureon et al. demonstrated the antitumor effect of HO-1 on prostate cancer [18].

In this study, we aimed to further investigate the role of HO-1 in prostatic cancer. HO-1 was overexpressed in PC-3 cells to evaluate the effect of HO-1 on cell proliferation and metastasis *in vitro* and tumor growth *in vivo*. The related mechanisms would also be investigated.

## Material and Methods

### Cell Culture and Chemicals

The PC-3 and DU145 cells were obtained from the Bioresource Collection and Research Center (Taiwan) and were cultured in the RPMI-1640 medium (Life Technologies, Rockville, MD) with 10% fetal calf serum (FCS; HyClone, UT) as described previously [19]. DAPI (4,6-diamino-2-phenylindole), propidium iodide, H<sub>2</sub>O<sub>2</sub>, and H<sub>2</sub>DCFDA (2',7'-dichlorofluorescein diacetate) were purchased from Sigma-Aldrich Co. (St. Louis, MO). Pyocyanin was purchased from Enzo Life Sciences Inc. (Farmingdale, NY, USA).

### HO-1 Expression Vector and Stable Transfection

The clone of human HO-1 cDNA in pOTB7 vector (MGC:1723) was purchased from Invitrogen (Carlsbad, CA). The human HO-1 expression vector was constructed by cloning a full length of HO-1 cDNA into the pcDNA3.1/zeo expression vector (Invitrogen) with *Eco RI* sites. Proper ligation and orientation was confirmed by extensive restriction mapping and sequencing. Electroporation was conducted using an ECM 830 Square Wave Electroporation System (BTX, San Diego, CA). Transfected PC-3 (PC-HO1-1 and PC-HO1-2) cells were selected by 100 µg/mL of Zeocin (Invitrogen). The DU145 cells were transiently transfected with HO-1 expression vector (DU-HO1) for 72 hours without Zeocin selected. For construction of the mock-transfected cells (PC-DNA and DU-DNA), cells were transfected with a controlled pcDNA3.1 expression vector and clonally selected in the same manner as described previously [20].

### Cell Proliferation and Viability Assay

The cell proliferations were measured using 3H-thymidine incorporation, CyQUANT cell proliferation assay (Invitrogen), and

WST-1 cell proliferation assay (abcam, Cambridge, UK), as described previously [21,22]. Cell viability was analyzed using the MTS assay (Promega Biosciences, Madison, WI).

### EdU Staining Proliferation Assay

The EdU fluorescence of cells was detected using Attune NxT acoustic focusing cytometer (Thermo Fisher Scientific Inc., Waltham, MA) as described previously [23]. Briefly, PC-DNA, PC-HO1-1, and PC-HO1-2 cells ( $5 \times 10^5$ ) were cultured in serum-free medium for 24 hours. After another 48 hours incubated with 10% serum medium, the cells were incubated with EdU (5-ethynyl-2'-deoxyuridine; 10 µM) for further 2 hours. Then, the cells were collected and analyzed using Click-iT EdU Flow Cytometry Assay Kits (Thermo Fisher Scientific Inc.).

### ROS Analysis

Cells were cultured in RPMI-1640 medium with 10% FCS for 48 hours, and then the cells were harvested with trypsin and washed twice with PBS. 20 µl of H<sub>2</sub>DCFDA added to the cell pellet and incubated at 37 °C and 5% CO<sub>2</sub> incubator for 30 min. After adding reactive oxygen species (ROS) inducer (20 µM of pyocyanin or 125 µM of H<sub>2</sub>O<sub>2</sub>) as indicated for 1 hour, cells were pelleted and then suspended in 500 µl of PBS. The ROS was analyzed using the FACS-Calibur Cytometer (BD Biosciences, Franklin Lakes, NJ). We also analyzed the total ROS induced by H<sub>2</sub>O<sub>2</sub> using immunofluorescence reader. Briefly, cells ( $3 \times 10^3$ /per well) were cultured in a 96-well plate for 48 hours and then washed twice with PBS. 200 µl of H<sub>2</sub>DCFDA (20 µM in RPMI 1640 medium with 2% FCS) were added to each well and then incubated for 30 min in incubator with 37 °C and 5% CO<sub>2</sub>. The 0, 125, and 50 M of H<sub>2</sub>O<sub>2</sub>, respectively, in RPMI 1640 medium with 10% FCS were added for 1 hour after cells were washed twice with PBS. The intensity of DCF-DA fluorescence was detected and quantified with the Chameleon Fluoro-Lumino-Photometer (Turku, Finland).

### Sub-G1 Cycle Analysis

Cells were treated with 125 µM of H<sub>2</sub>O<sub>2</sub> for 16 hours or serum starvation for 5 days to induce cell death. Cell cycle of sub-G1 analysis was performed and quantified using the FACS-Calibur E6147 Cytometer and CellQuest Pro 4.02 software (BD Biosciences) as described previously [21].

### Annexin V-FITC Apoptosis Detection

The cell pellets were harvested after treated with H<sub>2</sub>O<sub>2</sub> (500 µM) for 12 hours. The detection and quantification of cell apoptosis were performed after treated with Annexin V-FITC (BioVision Inc, Milpitas, CA) using the FACS-Calibur E6147 Cytometer (BD Biosciences) as described previously [22].

### Nuclear and Cytoplasmic Extraction

Cells were harvested with trypsin and washed twice with PBS. Nuclear and cytoplasmic fractions were separated using the NE-PER Nuclear and cytoplasmic extraction kit (Thermo, Rockford, NJ) as described previously [24].

### Immunoblot Assay

Equal quantities of cell extracts which was measured by BCA protein assay kit were separated onto a 10% SDS-PAGE gel, transferred and analyzed by the Western lightning plus-ECL detection system (Perkin Elmer, Inc., Waltham, MA). Antibodies

against HO (HO-1; Hsp32, Stressgen, Victoria, BC, Canada), PARP, cleaved PARP (BD Biosciences), N-cadherin, Vimentin (Abgent, San Diego, CA), Lamin B1 (Santa Cruz Biotechnology, Santa Cruz, CA), Slug, and  $\beta$ -actin (Millipore, Temecula, CA) were used.

### Immunofluorescence

Cells were seeded for 24 hours on sterile glass coverslips. The processes of fixation, permeabilization, and block were performed as described previously [25].

### F-actin Staining

Cells were seeded onto glass bottoms of the culture dishes (MatTek, Ashland, MD), then, precoated with fibronectin, and allowed to attach overnight. The F-actin protein expression was revealed by incubation with Texas Red X-Phalloidin and mounted with ProLongR Gold reagent (Invitrogen) as described previously [26].

### Real-time Reverse Transcription—polymerase Chain Reaction

Total RNA from cells was isolated using Trizol reagent. The cDNA was synthesized, and real-time polymerase chain reaction (qPCR) was performed as described previously [27]. The mRNA expressions of genes were assayed using the FAM dye-labeled TaqMan MGB probes for HO-1 (Hs00157965\_m1) and  $\beta$ -actin (Hs01060665\_g1), purchased from Applied Biosystems (Foster City, CA).

### Matrigel Invasion Assay

Cells ( $1 \times 10^5$ ) migrated to the matrigel-coated transmembrane for 24 hours. The images were captured using a digital camera connected to an inverted microscope (IX71, Olympus, Tokyo, Japan) with PAX-it Digital Image Management & Image Analysis and standardized for light intensity [28].

### Xenograft Animal Study

All animal experiments met the Guide for Laboratory Animal Facilities and Care as promulgated by Council of Agriculture Executive Yuan, Taiwan. The protocol was approved by the Chang Gung University Animal Research Committee (Permit Number: CGU15154). All methods were performed in accordance with the “Animal Welfare Law and Policy” (LAW3ANI). The male nude mice (BALB/cAnN-Foxn1, 4 weeks old) were randomized into two groups: PC-DNA ( $n = 6$ ) and PC-HO-1 ( $n = 6$ ). Animal were anesthetized intraperitoneally and equal volumes of cells ( $4 \times 10^6/100 \mu\text{l}$ ) were injected into the back of mice. Tumor volume was measured using vernier calipers and calculated as  $\pi/6 \times \text{larger diameter} \times (\text{smaller diameter})^2$  as described previously [28].

### Statistical Analysis

All the results are expressed as the mean  $\pm$  standard error (SE). Statistical analyses were performed using SigmaStat program for Windows, version 2.03 (SPSS Inc. Chicago, IL). The significance of difference was determined by Student's *t*-test or one way ANOVA with a *P* value less than 0.05 ( $*P < 0.05$ ) or 0.01 ( $**P < 0.01$ ).

## Results

### HO-1 Expression is Predominant in the Cytoplasm of PC-3 Cells With HO-1 Ectopic Overexpression

The expression vector containing human HO-1 cDNA was ectopically stably transfected into PC-3 cells to evaluate the

location of HO-1 protein in cells. Expressions of HO-1 in the selected clones (PC-HO1-1 and PC-HO1-2) were determined by immunoblot (Figure 1A) and RT-PCR (Figure 1B) assays. Further immunoblot assays with subcellular extraction revealed clearly that cytoplasmic and nuclear positive HO-1 was detected PC-3 cells; however, the majority of HO-1 expression in PC-HO1-1 and PC-HO1-2 cells located in the cytoplasm in comparison with PC-DNA cells (mock-transfected PC-3 cells) (Figure 1C). Similar results were found in the immunofluorescence assays in which HO-1 expressed in both nuclear and cytosol area, but predominantly in the cytoplasm in the HO-1-overexpressed PC-3 cells (Figure 1D).

### HO-1 Overexpression has No Obvious Effect on Cell Proliferation but Enhances Cell Invasion of PC-3 Cells In Vitro Via Modulation of EMT


The CyQUANT cell proliferation (Figure 2A) and WST-1 cell proliferation (Figure 2B) assays revealed that ectopic HO-1-overexpressed PC3 cells did not affect cell proliferation. Further study using the EdU staining proliferation assays also confirmed that overexpression of HO-1 did not affect cell proliferation assays *in vitro* (Figure 2C). The matrigel invasion assays indicated that PC-HO1-1 and PC-HO1-2 cells enhanced cell invasion to 1.8- and 2.3-folds, respectively, as compared with PC-DNA cells (Figure 2D). The immunoblot assays showed that ectopic overexpression of HO-1 upregulated N-cadherin, Vimentin, and Slug expressions in PC-3 cells (Figure 2E). Results of F-actin staining further suggested that modulation of epithelial—mesenchymal transition (EMT) altered the organization reminiscent of actin cytoskeletal as HO-1 overexpressed in PC-3 cells (Figure 2F).

### HO-1 Enhances Tumorigenesis of PC-3 Cells In Vivo

PC-DNA and PC-HO-1 cells were injected subcutaneously into the back of nude mice to determine the effects of HO-1 on tumorigenesis. After 34 days of growth, the tumor volume of tumors derived from PC-HO1-1 cells was more than 7 times the size of those from the PC-DNA cells ( $40.90 \pm 10.85 \text{ mm}^3$  vs.  $285.74 \pm 60.90 \text{ mm}^3$ ) (Figure 3A and B). There was no significant difference in the mean body weight of animals between two groups (Figure 3C). The weight of tumors derived from PC-HO1-1 cells was about 4 times the weight of tumors from the PC-DNA cells' group (Figure 3D). We randomly selected four tissues from each groups, and results of immunoblot assays confirmed that HO-1 was overexpressed in the xenograft tumors derived from PC-HO1-1 cells (Figure 3E).

### HO-1 Alleviates ROS Generation in PC-3 Cells

Treatments of 125  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  induced 24.8-fold ROS immunofluorescence intensity in PC-DNA cells, while the same dosage of  $\text{H}_2\text{O}_2$  induced only 15.1-fold and 17.5, respectively, ROS immunofluorescence intensity in PC-HO-1 and PC-HO1-2 cells as compared with the vehicle treatment (Figure 4A). Further flow cytometry assays showed similar results. Ectopic HO-1 overexpression blocked the ROS generation stimulated by 125  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  or 20  $\mu\text{M}$  of pyocyanin (a ROS inducer) treatments in PC-HO1-1 (Figure 4B) and PC-HO1-2 (Figure 4C) cells as compared with PC-DNA cells.




**Figure 1. Ectopic HO-1 overexpression is predominant in the cytoplasm areas of prostate carcinoma cells.** PC3 cells were transfected stably with HO-1 expression vector (PC-HO1-1 and PC-HO1-2) or pcDNA3.1 (PC-DNA). Expressions of HO-1 were analyzed using (A) immunoblot and (B) RT-qPCR assays. (C) HO-1 exhibition of ectopic overexpression was determined by subcellular extraction and immunoblotting. (D) The location patterns of HO-1 in PC-DNA, PC-HO1-1, and PC-HO1-2 cells were determined by immunofluorescence staining. The blue-stained nuclei of PC-DNA, PC-HO1-1, and PC-HO1-2 cells were stained by DAPI. All images were observed and recorded under the same settings of a fluorescence microscope with magnification of  $\times 400$ .


#### *HO-1 Prevents Cell Death Induced by Serum Starvation or H<sub>2</sub>O<sub>2</sub> Treatment in PC-3 Cells*

Although ectopic HO-1 overexpression did not enhance cell proliferation in normal condition as shown in Figure 2, which is contrary to the finding that HO-1 overexpression strengthened tumorigenesis shown in Figure 3, we discovered another possible mechanism for the increased *in vivo* tumorigenesis. Results of CyQUANT cell proliferation assays showed PC-HO1-1 and PC-HO1-2 cells, after 5 days of serum starvation, had higher cell proliferative rate as compared with PC-DNA cells (Figure 5A). Further immunoblot assays indicated that PC-HO1-1 (Figure 5B, top) and PC-HO1-2 (Figure 5B, bottom) cells had less cleaved form

of PARP protein as compared with PC-DNA cells after serum starvation, suggesting that ectopic HO-1 overexpression rendered cells more resistant to apoptosis after serum starvation for 3 or 5 days. This finding was supported by flow cytometry assays showing the sub-G1 cell percentage of PC-DNA cells was much higher than that of PC-HO1-1 and PC-HO1-2 cells after 5 days of serum starvation (Figure 5C, top). Quantitative analysis showed that PC-DNA, PC-HO1-1, and PC-HO1-2 cells induced 11.7%, 5.8%, and 9.5%, respectively, of cell apoptosis after serum starvation (Figure 5C, bottom). Flow cytometry assays revealed that PC-HO1-1 and PC-HO1-2 cells had lower percentage of cells in sub-G1 phase in comparison with PC-DNA cells after 250 μM of H<sub>2</sub>O<sub>2</sub> treatment for



**Figure 2. Ectopic overexpression of HO-1 modulates epithelial–mesenchymal transition and F-actin distribution in prostate carcinoma cells.** The cell proliferation of PC-DNA (●), PC-HO1-1 (○), and PC-HO1-2 (▼) detected by CyQUANT (A), WST-1 (B), and EdU (C) cell proliferation assay kit. The cell invasion of PC-DNA, PC-HO1-1, and PC-HO1-2 measured using the matrigel invasion assay for 24 hours of incubation. The scale bar is 150  $\mu$ m. (D). (E) The expressions of N-cadherin, Vimentin, and Slug in PC-DNA, PC-HO1-1, and PC-HO1-2 cells were determined by immunoblot assays. (F) Immunofluorescence staining of F-actin (red) and DAPI (blue) among PC-DNA, PC-HO1-1, and PC-HO1-2 cells. The scale bar is 10  $\mu$ m. \*  $p < 0.05$ , \*\* $p < 0.01$ .



**Figure 3. Effects of ectopic HO-1 overexpression on the tumorigenesis of prostate carcinoma cells.** (A) Four-week-old male athymic nude (nu/nu) mice were randomized into two groups: PC-DNA ( $n = 6$ ) and PC-HO-1 ( $n = 6$ ). PC-DNA cells and PC-HO-1 cells ( $1 \times 10^6$ ) were injected s.c. in the dorsal area, respectively. Tumor growth rates (B) and animal body weight (C) were measured per 3 days starting at 1 week of growth point (day 1) in which the tumors became perceptible under the skin after inoculation. Subsequently, tumor volumes were measured as indicated days. The tumor weight (D) and levels of HO-1 protein expression (E) were determined after animals were sacrificed. \*  $p < 0.05$ , \*\*  $p < 0.01$ .

16 hours (Figure 5D). Results of MTS assays also indicated that  $H_2O_2$  treatment could decrease cell viability dose dependently with PC-HO1-1 and PC-HO1-2 cells much more resistant to  $H_2O_2$  treatment (Figure 5E). Results of flow cytometry with double stains for PI and Annexin V-FITC also showed that after 12 hours of 500  $\mu M$  of  $H_2O_2$ , PC-HO1-1 and PC-HO1-2 cells had less apoptotic cells (Figure 5F).


#### HO-1 Alleviates ROS Generation and Cell Death Induced by Serum Starvation or $H_2O_2$ Treatment in DU145 Cells

To confirm the antioxidant and antiapoptotic characteristics of HO-1 in prostate carcinoma cells, we repeated the experiments in Figure 5 using another prostate carcinoma cells, DU145. Immunoblot (Figure 6A, top) and RT-qPCR (Figure 6A, bottom) assays were applied to determine expressions of HO-1 after transient overexpression of HO-1 in DU145 (DU-HO1) cells. Results of MTS assays indicated transient overexpression of HO-1 (DU-HO1) could increase cell viability after serum starvation for 3 days as compared with mock-transfected DU145 (DU-DNA) cells (DU-DNA) (Figure 6B). The immunoblot assays also indicated

DU-HO1 cells had less cleaved form of PARP protein after serum starvation for 3 days (Figure 6C). Flow cytometry assays revealed that DU-HO1 cells had lower percentage of cells in sub-G1 phase in comparison with DU-DNA cells after serum starvation for 3 days (Figure 6D). Flow cytometry assays with PI and annexin V-FITC double stains showed transient HO-1 overexpression had less apoptotic cells after 12 hours of 500  $\mu M$  of  $H_2O_2$  treatment (Figure 6E). Further flow cytometry assays showed ectopic HO-1 overexpression in DU145 cells decreased ROS generation stimulated by 125  $\mu M$  of  $H_2O_2$  (Figure 6F, top) or 20  $\mu M$  of pyocyanin (Figure 6F, bottom) treatments.

#### Discussion

Heme could induce oxidative stress in tissues. Therefore, heme-degradation pathway is deemed as a tissue protection pathway. Accordingly, HO-1 activity, which is responsible for heme degradation and induced by numerous stresses, is vital for cell survival after stress. HO-1 converts prooxidant heme into CO, biliverdin, and iron [1–3]. Iron could further induce ferritin synthesis, which has cytoprotective effect [6]. Biliverdin would be reduced by biliverdin




**Figure 4. HO-1 alleviates ROS generation in prostate carcinoma PC-3 cells.** (A) Fold change of ROS generation through treatments of 125  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  on PC-DNA, PC-HO1-1, and PC-HO1-2 cells detected by DCF-DA dye determined by immunofluorescence reader. The values shown in each bar chart represent the fold induction  $\pm$  SE ( $n = 6$ ) of intense fluorescence of cells induced by  $\text{H}_2\text{O}_2$  relative to that of solvent-treated group (\*\* $P < 0.01$ ) or relative to that of indicated (## $P < 0.05$ ; ### $P < 0.01$ ). Ectopic overexpression of HO-1 blocked ROS generation stimulated by 125  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  or 20  $\mu\text{M}$  of pyrocyanin determined by flow cytometry in PC-HO1-1 (B) and PC-HO1-2 (C) cells.

reductase to form bilirubin. Both biliverdin and bilirubin are capable of eliminating ROS, thus having antioxidant effect [6]. CO could increase cGMP production and affect numerous signaling pathways. In general, the stress-induced HO-1 activity leads to antiinflammatory, antiapoptotic, and antioxidant effects [6,29].

To study HO-1 effect on PC-3 cells, HO-1 was overexpressed in PC-3 cells and PC-HO1-1 and PC-HO1-2 cells were obtained (Figure 1A and B). As shown in Figure 1C and D, HO-1 expressed predominantly in the cytoplasm of PC-3 cells by immunoblotting and immunofluorescence assays. Our results are in agreement with general concept that HO-1 localizes mainly in the cytoplasm where it is associated with endoplasmic reticulum, caveolin, or mitochondria [2]. However, this result is different from other studies showing


hemin-treated prostate carcinoma cells and the HO-1-overexpressed PC3 cells in xenograft expressed HO-1 mainly located in the nuclei [30,31]. One important aspect of HO-1 effect on tumor biology might be related to its cellular localization, and the subcellular location of HO-1 in carcinoma cells which may account for the divergent effect on the different types of cancer has been discussed [29].

Results of previous studies investigating the functions and expression of HO-1 in prostate carcinoma cells *in vitro* and *in vivo* are still not consistent. For example, a significant elevation of HO-1 expression in malignant prostate cancer epithelial cells has been shown and, thus, HO-1 inhibitor is deemed as a potential therapeutic agent for prostate cancer [32,33]; however, results from other studies



**Figure 5. HO-1 attenuates serum starvation and H<sub>2</sub>O<sub>2</sub> treatment induced cell apoptosis in PC-3 cells.** (A) Cell proliferation ratio of PC-DNA, PC-HO1-1, and PC-HO1-2 cells after 5 days of serum starvation using CyQUANT cell proliferation assay kit. The proliferation ratio of cells incubated in full medium was regarded as 1. (B) PARP protein expressions of PC-DNA, PC-HO1-1, and PC-HO1-2 cells after 3 and 5 days of serum starvation utilizing immunoblot assays. (C) Fluorescence intensity of sub-G1 phase detected PI staining using flow cytometry after 5 days of serum starvation. (D) Fluorescence intensity of sub-G1 phase after 250 μM of H<sub>2</sub>O<sub>2</sub> treatment for 16 hours detected by PI dye using flow cytometry. (E) Cell viability with different concentrations of H<sub>2</sub>O<sub>2</sub> treated PC-DNA, PC-HO1-2, and PC-HO1-1 cells utilizing MTS assays. (F) Cell apoptosis was determined by Annexin V-FITC in conjunction with PI staining. Fluorescence intensity for Annexin V-FITC is plotted on the x-axis, and PI is plotted on the y-axis. \* p < 0.05, \*\* p < 0.01.





**Figure 6. HO-1 attenuates oxidation and apoptosis under serum starvation and H<sub>2</sub>O<sub>2</sub> treatment in prostate carcinoma DU145 cells.** DU145 cells were transfected transiently with HO-1 expression vector (DU-HO1) or pcDNA3.1 (DU-DNA) for 72 hours. (A) Expressions of HO-1 were analyzed by immunoblot (top) and RT-qPCR (bottom) assays. (B) Cell viability of DU-DNA and DU-HO1 cells after 3 days of serum starvation utilizing MTS assays. (C) PARP protein expressions of DU-DNA and DU-HO1 cells after 3 and 5 days of serum starvation utilizing immunoblot assays. (D) Percentage of sub-G1 phase after 3 days of serum starvation detected PI staining using flow cytometry. (E) Cell apoptosis of DU-DNA and DU-HO1 cells induced by 500 μM of H<sub>2</sub>O<sub>2</sub> for 12 hours were determined by Annexin V-FITC in conjunction with PI staining. (F) ROS generation of DU-DNA and DU-HO1 cells stimulated by 125 μM of H<sub>2</sub>O<sub>2</sub> (upper) or 20 μM of pyrocyanin (bottom) treatments determining by flow cytometry. \* p< 0.05, \*\*p<0.01.

indicated HO-1 reduced cell proliferation and metastasis in prostate cancer cells, thus implying stimulation of HO-1 expression may be a therapeutic direction for prostate cancer [18,30,31]. In this study, our result showed that the cell proliferation rate of PC-HO1-1 and PC-HO1-2 cells was not changed as compared with PC-DNA cells *in vitro* (Figure 2A–C). Moreover, PC-HO1 cells had higher invasive potential than PC-DNA cells (Figure 2D).

EMT is the process during which epithelial cells differentiate into mesenchymal cells. EMT is an important process for normal development and wound healing. However, EMT also plays a crucial role in cancer progression. For example, after EMT, cancer cells get more stem-cell like characteristics, which makes cancer cells more resistant to chemotherapy [34–36]. The increased invasiveness of cancer cells after EMT further worsens the clinical outcome [37]. EMT is under delicate control. So far, at least three transcriptional families have been shown to involve in EMT regulations, which include Snail/Slug, ZEB1/2, and Twist families [38]. After EMT, the mesenchymal cell markers, such as vimentin or N-cadherin, would also increase [39]. Figure 2E showed that HO-1 overexpression would increase Slug, N-cadherin, and vimentin expression in PC-3 cells, which demonstrated HO-1 overexpression would induce EMT in PC-3 cells to enhance cell invasiveness shown in Figure 2D. Apart from EMT, HO-1 overexpression also increased F-actin synthesis, thus further increasing cell motility (Figure 2F).

Although the cell proliferation rate did not change after HO-1 overexpression in PC-3 cells *in vitro*, the tumorigenesis ability increased. As shown in Figure 3A, B, and D, the xenograft animal study revealed that tumors from PC-HO1-1 had higher tumor volume as compared with PC-DNA cells. To investigate the reason behind this, we applied H<sub>2</sub>O<sub>2</sub> or pyocyanin to treat PC-DNA and PC-HO1-1 cells. Figure 4 showed that H<sub>2</sub>O<sub>2</sub> increased ROS in both PC-DNA cells and PC-HO1 cells with the latter having much less amount of ROS, further demonstrating the scavenger role of HO-1 for ROS. The role of ROS in tumorigenesis has been discussed. Because ROS plays a causal role in tumor development but high ROS level has the potential to actually block tumorigenesis [40,41], our results are in agreement with previous studies demonstrating HO-1 can reduce H<sub>2</sub>O<sub>2</sub> or pyocyanin-induced oxidative damage [42,43], which may account for the higher tumorigenesis of PC-HO1-1 cells noted *in vivo* shown in Figure 3.

Figure 5A showed that after 5 days of starvation, the cell proliferation rate of PC-HO1-1 and PC-HO1-2 cells was higher than PC-DNA cells. The western blot result further revealed that PC-HO1-1 had lower cleaved PARP than PC-DNA cells (Figure 5B), indicating the lower apoptosis rate of PC-HO1-1 cells. The result was also supported by the flow cytometry data showing in normal medium, the percentage of apoptotic cells was similar in PC-DNA and PC-HO1-1 cells; however, after serum starvation, the apoptotic cell percentage was much higher in PC-DNA cells (Figure 5C). The MTS test showed that after 250–500 μM H<sub>2</sub>O<sub>2</sub> treatment, PC-HO1-1 had higher cell proliferative rate than PC-DNA cells (Figure 5D), which is supported by the flow cytometry result showing after H<sub>2</sub>O<sub>2</sub> treatment, PC-HO1-1 had lower percentage of apoptotic cells (Figure 5E and F). These results are in agreement with previous studies which indicated HO-1 had antiapoptotic effect [29,44].

The antioxidant and antiapoptotic effects of HO-1 were also evaluated in DU145 cells (Figure 6). Overexpression of HO-1 in DU145 cells decreased cell apoptosis induced by serum starvation or H<sub>2</sub>O<sub>2</sub> treatment (Figure 6A–D). The antioxidant effect of HO-1 in

DU145 cells was also observed (Figure 6E). Taken together, the result from Figures 4–6 suggested that HO-1 overexpression rendered PC-3 cells more resistant to serum starvation or ROS-induced treatment, which may contribute to the higher tumorigenesis of PC-HO1-1 cells noted *in vivo* (Figure 3).

## Conclusion

Our results suggested that although ectopic HO-1 overexpression did not increase cell growth *in vitro*, it did enhance cell proliferation under the stress of serum starvation *in vitro* and tumorigenesis *in vivo*. HO-1 overexpression further enhanced prostate cancer cell invasion through induction of EMT. The antioxidant and antiapoptotic effects of HO-1 shown in our study may attribute to the higher tumorigenesis found in the xenograft animal model.

## Authors' Contributions

K.-C.C. and K.-H.T. wrote the manuscript and designed this experiment and contributed equally to this work; Y.-H.L., C.-P.H., K.-S.C., H.-H.T., T.-H.F., Y.-S.S., and C.-C.C. helped conduct the experiment and analysis data; H.-H.J. was in charge of the whole experiment conduction and paper writing.

## Conflicts of Interest

The authors declare that no conflict of interest exists.

## Acknowledgment

This research was supported by grants from the Taiwan Ministry of Science and Technology (MOST-107-2314-B-182A-017-MY3 and MOST-105-2320-B-182-020-MY3) and Chang Gung Memorial Hospital (CRRPD1F0041-3, CMRPD1F0141-3, CMRPD1I0111-3, CMRPG3G0821, CMRPG3F0810-3, and CMRP-G3H1321-3).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.tranon.2019.10.008>.

## References

- [1] Tenhunen R, Marver HS and Schmid R (1968). The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc Natl Acad Sci USA* **61**(2), 748–755.
- [2] Ryter SW, Alam J and Choi AM (2006). Heme oxygenase-1/carbon monoxide: from basic science to therapeutic applications. *Physiol Rev* **86**(2), 583–650.
- [3] Maines MD (1997). The heme oxygenase system: a regulator of second messenger gases. *Annu Rev Pharmacol Toxicol* **37**, 517–554.
- [4] Trakshel GM, Kutty RK and Maines MD (1986). Purification and characterization of the major constitutive form of testicular heme oxygenase. The noninducible isoform. *J Biol Chem* **261**(24), 11131–11137.
- [5] Grochot-Przeczek A, Kotlinowski J, Kozakowska M, Starowicz K, Jagodzinska J and Stachurska A, et al (2008). Heme oxygenase-1 and the vascular bed: from molecular mechanisms to therapeutic opportunities. *Antioxidants Redox Signal* **10**(10), 1767–1812.
- [6] Loboda A, Jozkowicz A and Dulak J (2015). HO-1/CO system in tumor growth, angiogenesis and metabolism - targeting HO-1 as an anti-tumor therapy. *Vasc Pharmacol* **74**, 11–22.
- [7] Podkalicka P, Mucha O, Józkowicz A, Dulak J and Łoboda A (2018). Heme oxygenase inhibition in cancers: possible tools and targets. *Contemp Oncol* **22**(1A), 23–32.
- [8] Owis D, Legat M, Grzela T, Niderla J, Wilczek E and Wilczynski GM, et al (2006). Heme oxygenase-1 protects tumor cells against photodynamic therapy-mediated cytotoxicity. *Oncogene* **25**(24), 3365–3374.
- [9] Na HK and Surh YJ (2014 Feb). Oncogenic potential of Nrf2 and its principal target protein heme oxygenase-1. *Free Radical Biol Med* **67**, 353–365.

- [10] Deshane J, Chen S, Caballero S, Grochot-Przeczek A, Was H and Li Calzi S, et al (2007). Stromal cell-derived factor 1 promotes angiogenesis via a heme oxygenase 1-dependent mechanism. *J Exp Med* **204**(3), 605–618.
- [11] Banerjee P, Basu A, Datta D, Gasser M, Waaga-Gasser AM and Pal S (2011). The heme oxygenase-1 protein is overexpressed in human renal cancer cells following activation of the Ras-Raf-ERK pathway and mediates anti-apoptotic signal. *J Biol Chem* **286**(38), 33580–33590.
- [12] Loboda A, Jazwa A, Wegiel B, Jozkowicz A and Dulak J (2005). Heme oxygenase-1-dependent and -independent regulation of angiogenic genes expression: effect of cobalt protoporphyrin and cobalt chloride on VEGF and IL-8 synthesis in human microvascular endothelial cells. *Cell Mol Biol* **51**(4), 347–355.
- [13] Birrane G, Li H, Yang S, Tachado SD and Seng S (2013). Cigarette smoke induces nuclear translocation of heme oxygenase (HO-1) in prostate cancer cells: nuclear HO-1 promotes vascular endothelial growth factor secretion. *Int J Oncol* **42**(6), 1919–1928.
- [14] Was H, Dulak J and Jozkowicz A (2010). Heme oxygenase-1 in tumor biology and therapy. *Curr Drug Targets* **11**(12), 1551–1570.
- [15] Hill M, Pereira V, Chauveau C, Zagani R, Remy S and Tesson L, et al (2005). Heme oxygenase-1 inhibits rat and human breast cancer cell proliferation: mutual cross inhibition with indoleamine 2,3-dioxygenase. *FASEB J* **19**(14), 1957–1968.
- [16] Skrzypek K, Tertilt M, Golda S, Ciesla M, Weglarczyk K and Collet G, et al (2013). Interplay between heme oxygenase-1 and miR-378 affects non-small cell lung carcinoma growth, vascularization, and metastasis. *Antioxidants Redox Signal* **19**(7), 644–660.
- [17] Halin Bergström S, Nilsson M, Adamo H, Thysell E, Jernberg E and Stattin P, et al (2016). Extratumoral heme oxygenase-1 (HO-1) expressing macrophages likely promote primary and metastatic prostate tumor growth. *PLoS One* **11**(6):e0157280.
- [18] Gueron G, Giudice J, Valacco P, Paez A, Elguero B and Toscani M, et al (2014). Heme-oxygenase-1 implications in cell morphology and the adhesive behavior of prostate cancer cells. *Oncotarget* **5**(12), 4087–4102.
- [19] Tsui KH, Chung LC, Feng TH, Lee TY, Chang PL and Chen WT, et al (2015). Divergent effect of liver X receptor agonists on prostate-specific antigen expression is dependent on androgen receptor in prostate carcinoma cells. *Prostate* **75**(6), 603–615.
- [20] Tsui KH, Hsu SY, Chung LC, Lin YH, Feng TH and Lee TY, et al (2015). Growth differentiation factor-15: a p53- and demethylation-upregulating gene represses cell proliferation, invasion, and tumorigenesis in bladder carcinoma cells. *Sci Rep* **5**, 12870.
- [21] Tsui KH, Lin YH, Chung LC, Chuang ST, Feng TH and Chiang KC, et al (2016). Prostate-derived ets factor represses tumorigenesis and modulates epithelial-to-mesenchymal transition in bladder carcinoma cells. *Cancer Lett* **375**(1), 142–151.
- [22] Tsui KH, Chang YL, Yang PS, Hou CP, Lin YH and Lin BW, et al (2018). The inhibitory effects of capillarisin on cell proliferation and invasion of prostate carcinoma cells. *Cell Prolif* **51**(2):e12429.
- [23] Chang KS, Tsui KH, Lin YH, Hou CP, Feng TH and Juang HH (2019). Migration and invasion enhancer I is the NF- $\kappa$ B-inducing gene enhancing the cell proliferation and invasion ability of human prostate carcinoma in vitro and in vivo. *Cancers* **11**(10), 1486.
- [24] Tsui KH, Hou CP, Chang KS, Lin YH, Feng TH and Chen CC, et al (2019). Metallothionein 3 is a hypoxia-upregulated oncogene enhancing cell invasion and tumorigenesis in human bladder carcinoma cells. *Int J Mol Sci* **20**(4), 980.
- [25] Juang HH, Chung LC, Sung HC, Feng TH, Lee YH and Chang PL, et al (2013). Metallothionein 3: an androgen-upregulated gene enhances cell invasion and tumorigenesis of prostate carcinoma cells. *Prostate* **73**(14), 1495–1506.
- [26] Chiang KC, Yeh CN, Chung LC, Feng TH, Sun CC and Chen MF, et al (2015). WNT-1 inducible signaling pathway protein 1 enhances growth and tumorigenesis in human breast cancer. *Sci Rep* **5**, 8686.
- [27] Lee JC, Chung LC, Chen YJ, Feng TH and Juang HH (2014). N-myc downstream-regulated gene 1 downregulates cell proliferation, invasiveness, and tumorigenesis in human oral squamous cell carcinoma. *Cancer Lett* **355**(2), 242–252.
- [28] Chung LC, Chiang KC, Feng TH, Chang KS, Chuang ST and Chen YJ, et al (2017). Caffeic acid phenethyl ester upregulates N-myc downstream regulated gene 1 via ERK pathway to inhibit human oral cancer cell growth in vitro and in vivo. *Mol Nutr Food Res* **61**(9), 1600842.
- [29] Nititi M, Piras S, Marinari UM, Moretta L, Pronzato MA and Furfaro AL (2017). HO-1 induction in cancer progression: a matter of cell adaptation. *Antioxidants* **6**(2), 29.
- [30] Gueron G, De Siervi A, Ferrando M, Salierno M, De Luca P and Elguero B, et al (2009). Critical role of endogenous heme oxygenase 1 as a tuner of the invasive potential of prostate cancer cells. *Mol Cancer Res* **7**(11), 1745–1755.
- [31] Sacca P, Meiss R, Casas G, Mazza O, Calvo JC and Navone N, et al (2007). Nuclear translocation of haeme oxygenase-1 is associated to prostate cancer. *Br J Canc* **97**(12), 1683–1689.
- [32] Maines MD and Abrahamsson PA (1996). Expression of heme oxygenase-1 (HSP32) in human prostate: normal, hyperplastic, and tumor tissue distribution. *Urology* **47**(5), 727–733.
- [33] Alaoui-Jamali MA, Bismar TA, Gupta A, Szarek WA, Su J and Song W, et al (2009). A novel experimental heme oxygenase-1-targeted therapy for hormone-refractory prostate cancer. *Cancer Res* **69**(20), 8017–8024.
- [34] Thierry JP, Aclouque H, Huang RY and Nieto MA (2009). Epithelial-mesenchymal transitions in development and disease. *Cell* **139**(5), 871–890.
- [35] Scheel C, Eaton EN, Li SH, Chaffer CL, Reinhardt F and Kah KJ, et al (2011). Paracrine and autocrine signals induce and maintain mesenchymal and stem cell states in the breast. *Cell* **145**(6), 926–940.
- [36] Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A and Zhou AY, et al (2008). The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* **133**(4), 704–715.
- [37] Hanahan D and Weinberg RA (2011). Hallmarks of cancer: the next generation. *Cell* **144**(5), 646–674.
- [38] Peinado H, Olmeda D and Cano A (2007). Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat Rev Cancer* **7**(6), 415–428.
- [39] Christiansen JJ and Rajasekaran AK (2006). Reassessing epithelial to mesenchymal transition as a prerequisite for carcinoma invasion and metastasis. *Cancer Res* **66**(17), 8319–8326.
- [40] Sullivan LB and Chandel NS (2014). Mitochondrial reactive oxygen species and cancer. *Cancer Metabol* **2**, 17.
- [41] Chio IIC and Tuveson DA (2017). ROS in cancer: the burning question. *Trends Mol Med* **23**(5), 411–429.
- [42] Lang D, Reuter S, Buzescu T, August C and Heidenreich S (2005). Heme-induced heme oxygenase-1 (HO-1) in human monocytes inhibits apoptosis despite caspase-3 up-regulation. *Int Immunol* **17**(2), 155–165.
- [43] Tobiasch E, Gunther L and Bach FH (2001). Heme oxygenase-1 protects pancreatic beta cells from apoptosis caused by various stimuli. *J Investig Med* **49**(6), 566–571.
- [44] Jozkowicz A, Was H and Dulak J (2007). Heme oxygenase-1 in tumors: is it a false friend? *Antioxidants Redox Signal* **9**(12), 2099–2117.