

In-Flow Generation of Thionyl Fluoride (SO₂F₂) Enables the Rapid and Efficient Synthesis of Acyl Fluorides from Carboxylic Acids

Daniele Mazzarella,^{||} Jelena Stanić,^{||} Miguel Bernús, Arad Seyed Mehdi, Cassandra J. Henderson, Omar Boutureira, and Timothy Noël^{*}



Cite This: *JACS Au* 2024, 4, 2989–2994




Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information



ABSTRACT: Herein, we report an approach for generating thionyl fluoride (SO₂F₂) from the commodity chemicals thionyl chloride (SOCl₂) and potassium fluoride (KF). The methodology relies on a microfluidic device that can efficiently produce and dose this toxic gaseous reagent under extremely mild and safe conditions. Subsequently, the in situ-generated thionyl fluoride is reacted with an array of structurally and electronically differing carboxylic acids, leading to the direct and efficient synthesis of highly sought-after acyl fluorides. Importantly, our investigation also highlights the inherent modularity of this flow-based platform. We demonstrate the adaptability of this approach by not only synthesizing acyl fluorides but also directly converting carboxylic acids into a diverse array of valuable compounds such as esters, thioesters, amides, and ketones. This versatility showcases the potential of this approach for a wide range of synthetic applications, underscoring its significance in the realm of chemical synthesis.

KEYWORDS: fluorine chemistry, flow chemistry, gaseous reagents, on-demand synthesis, multistep synthesis

In contemporary synthetic laboratories, the utilization of toxic and hazardous gaseous compounds is subject to stringent regulations and control measures.¹ The elusive nature of gases makes handling and dosing of these reagents a formidable challenge when employing traditional batch equipment.² This has prompted the scientific community to devise engineered chemicals that can release *in situ* the desired gases via a chemical reaction.³ Alternatively, some solid reagents have been developed to serve as gas surrogates to perform the same types of transformations.⁴ However, these approaches often suffer from inefficiency due to the formation of stoichiometric byproducts. Furthermore, they frequently necessitate the initial use of the parent gases for the synthesis of the reagents, overall lowering the atom economy.

In our laboratory,⁵ we have recently made significant strides in leveraging the power of flow chemistry⁶ to directly harness and manage various gaseous reagents.⁷ Within this research framework, we have unveiled a modular flow platform capable of producing SO₂F₂ from readily available, bench-stable chemicals such as KF and SO₂Cl₂ (Scheme 1A) through a Cl–F exchange process.^{5a} The intrinsic containment properties of this flow system ensure the safe and controlled generation of the gaseous reagent while also facilitating precise dosing of the reactive gas.^{7,8} This breakthrough has enabled the execution of

a diverse range of SuFEx (sulfur(VI) fluoride exchange)⁹ ligations on a wide spectrum of compounds, including small molecules, biorelevant compounds, peptides, and proteins.


Recently, we wondered whether we could extend the range of gases generated through this microfluidic device to thionyl fluoride, SO₂F₂. This gas has long been overlooked as a deoxyfluorinating reagent due to safety concerns associated with its handling and toxicity. Only recently, work by Sammis and co-workers demonstrated how stock solutions of this gas could be produced through an *ex situ* approach.¹⁰ However, this strategy requires multiple manipulations due to the batch setup in order to avoid the presence of undesired F/Cl mixed species SOFCl and stoichiometric amounts of HCl, limiting its practicality and potential applications. Our interest in generating SO₂F₂ stemmed from its ability to swiftly convert abundant carboxylic acids into acyl fluorides, as detailed by the

Received: April 11, 2024

Revised: June 11, 2024

Accepted: June 11, 2024

Published: July 12, 2024



Scheme 1. (A) In-Flow Generation of Sulfuryl Fluoride from Sulfuryl Chloride Enables the Rapid and Direct Synthesis of Fluorosulfates and Sulfamoyl Fluorides; (B) Selection of Reagents Capable of Converting Carboxylic Acids into Acyl Fluorides; (C) In-Flow Generation of Thionyl Fluoride Allows the Rapid and Direct Synthesis of Acyl Fluorides





Table 1. Optimization of the Reaction Conditions^a



entry	flow rate 1st reactor (mL min ⁻¹)	flow rate 2nd reactor (mL min ⁻¹)	SOCl ₂ /acid ratio	yield (%) ^b
1	0.400	0.500	4:1	>95
2	0.333	0.500	2:1	>95
3	0.262	0.500	1.1:1	84
4 ^c	0.333	0.500	2:1	>95
5	0.600	1.00	1.5:1	94
6	1.20	2.00	1.5:1	>95
7	1.50	2.50	1.5:1	92

^aReactions performed on a 0.5 mmol scale of carboxylic acid **1**, using 2.5 equiv of triethylamine and 4–1.1 equiv of thionyl fluoride. ^bYields were determined by ¹⁹F NMR analysis, using 1,2-difluorobenzene as the internal standard. ^cReaction performed using 3 equiv of Et₃N.


group of Sammis.^{10b} This stands in stark contrast to the limitations of low atom economy, limited functional group tolerability, and high costs associated with the use of engineered reagents like DAST, cyanuric fluoride, Deoxo-Fluor, or Fluolead, among others (Scheme 1B).¹¹ Acyl fluorides are in fact an enticing class of carboxylic acid derivatives known for their synthetic utility as well as their enhanced stability and peculiar reactivity compared to their chloride analogues.¹² Acyl fluorides are isolable and exhibit increased stability with respect to other acyl halides and yet require mild reaction conditions to engage in diverse synthetic transformations.¹³ Thus, when reacted with nucleophiles, they offer a straightforward route to a wide range of valuable products such as esters, amides, or thioesters, almost regardless of the steric and electronic properties.¹⁴

Herein, we report the development of a strategy that harnesses the power of a microfluidic device to form thionyl fluoride and swiftly use it to convert carboxylic acids into acyl fluorides (Scheme 1C). Furthermore, we demonstrate the feasibility of a multistep flow approach¹⁵ where the carboxylic acids are directly converted into amides, esters, thioesters, and ketones.

Our investigation commenced with an exploration of the feasibility of SOF₂ generation by flowing a solution of SOCl₂ in CH₃CN through a packed bed reactor filled with a 1:1 mixture of KF and glass beads (see the Supporting Information for

details). We observed efficient and consistent formation of the coveted gaseous reagent regardless of the residence time of the solution within the packed bed reactor. Interestingly, the reactor cartridge itself has a reduced lifespan when decreasing the residence time of the solution within it (see the Supporting Information for details). We reasoned that higher flow rates might lead to the formation of preferred flow channels, which prevent SOCl₂ from reacting with the remaining KF present in the packed bed reactor.¹⁶ This issue can be effectively addressed by using relatively low flow rates (up to 15 mmol of SOF₂ was produced by a single cartridge at 0.1 mL/min; see the Supporting Information for further discussion). Subsequently, we coupled the thionyl fluoride generator to a stream of CH₃CN solution containing a mixture of model substrate 4-phenylbenzoic acid **1** and Et₃N and studied the influence of the stoichiometry and flow rates of the different feeding solutions (Table 1).

This process demonstrated exceptional efficiency, as the acyl fluoride formation occurred in high yields when employing 4, 2, and 1.1 equiv of thionyl chloride with respect to the carboxylic acid (entries 1–3). Increasing the amount of the Et₃N did not diminish the acyl fluoride formation (entry 4). Furthermore, the reaction displayed impressive speed, with optimal yields achieved even at reduced residence times in the second reactor of 90, 45, or as little as 36 s (entries 5–7). In the case of entry 4, the productivity of this process amounts to

Scheme 2. Array of Acyl Fluorides Synthesized by Means of Our Device^b


^aThe acyl fluoride was directly isolated. ^bTo facilitate isolation, the solution of the acyl fluoride was treated with 1.05 equiv of *N*-hydroxyphthalimide, and yields (%) refer to the corresponding esters unless otherwise indicated. The yields of the acyl fluorides, as measured by quantitative ¹⁹F NMR, have been reported in the [Supporting Information](#). *t*_{R1}, *t*_{R2}: residence times in the first and second reactors, respectively.

0.792 g h⁻¹, and the space-time yield amounts to 149 g L⁻¹ h⁻¹.

To showcase the robustness and versatility of this process, we selected the conditions detailed in entry 4 and varied the carboxylic acid partner (Scheme 2). We began by examining a wide array of aromatic carboxylic acids. The reaction is not very sensitive to the electronic nature of the substituents or their position on the aromatic ring. Indeed, electron-rich (2, 3), electron-neutral (4), and electron-poor derivatives (5, 6), as well as acids with substituents on the ortho (7) and meta (8) positions, were swiftly converted into the corresponding acyl fluorides (79–93% yields). As a limitation to this method, highly encumbered carboxylic acids did not afford the desired acyl fluoride (see the [Supporting Information](#) for further discussion). Moreover, the reaction also took place when

heteroaromatic (9–11) or ferrocenyl (12) derivatives were used (60–91% yields). Subsequently, we assessed the generality of the process in terms of alkyl carboxylic acids. Primary (13), secondary (14), and tertiary (15) carboxylic compounds all yielded the corresponding acyl fluorides in good-to-excellent yields (78–95% yields). Similar success was observed when an α,β -unsaturated carboxylic acid was used (16, 81% yield). Moreover, several Boc-protected α -amino acidic derivatives, such as glycine (17), alanine (18), leucine (19), proline (20), and phenylalanine (21), as well as non-natural structures such as cyclovaline (22) and azabicyclo[2.2.1]heptane-3-carboxylic acid (23), were smoothly converted into the targeted acyl fluoride derivatives (41–97% yield) with very little erosion of their enantiopurity (see the [Supporting Information](#)). Finally, we demonstrated

Scheme 3. Employing the Microfluidic Setup for the Synthesis of Various Acyl Derivatives by Fluorine Displacement



^a 3 equiv of Et₃N was used in the second module for entries **29** and **31**, and 2.5 equiv for entries **30** and **32**. t_{R1}, t_{R2}, t_{R3}: residence times in the first, second, and third reactors, respectively.

how the mild conditions of this approach make it suitable for the functionalization of structurally diverse biorelevant compounds decorated with various functional groups, such as ibuprofen (**24**), pinonic acid (**25**), biotin (**26**), dehydrocholic acid (**27**), and gibberellic acid (**28**) (55–94% yields). Crucially, this method enables the exclusive formation of the acyl fluoride even in the presence of a ketone functionality (see **25** and **27**), which would be rapidly converted into a difluorinated motif in the presence of other deoxyfluorinating reagents such as DAST or Deoxo-Fluor.¹⁷

After having found a set of optimal conditions to promote the formation of acyl fluorides and having assessed the generality of the scope, we embarked upon an endeavor to leverage the full potential of this class of compounds with this microfluidic device. As previously described, acyl fluorides are appealing intermediates to forge new carbon–carbon and carbon–heteroatom bonds. Exploiting the advantages offered by the modular flow chemistry approach, we envisioned a device comprising three sections (Scheme 3): the first module, in which SOCl₂ is converted to SOF₂ through Cl–F exchange; the second, in which the gaseous reagent is reacted with a carboxylic acid to yield the acyl fluoride; and the third, in which the acyl fluoride is finally mixed with a chosen nucleophile to furnish the target compound. This required reoptimization of the procedure (see the Supporting Information for details). Crucially, the equivalents of SOF₂ were lowered to 1.1 to avoid subsequent undesired reactions with the nucleophilic component within the third module. This adjustment, in turn, mandated an extension of the residence time within the second module to 6 min to ensure optimal acyl fluoride formation. It is worth noting that the productive use of 1.1 equiv of SOF₂ in batch conditions would be extremely challenging to achieve, as the gas would inevitably evolve toward the headspace of the reactor. Under these finely tuned

reaction conditions, our investigation was directed toward the exploration of potential coupling nucleophiles. As expected, a thiol, an amine, and an alcohol could all be employed to obtain the corresponding thioester (**29**, 61% yield), amide (**30**, 74% yield), and ester (**31**, 61% yield) derivatives of dehydrocholic acid upon acyl substitution. Furthermore, we evaluated the possibility of forging C–C bonds. After a brief optimization of the reaction conditions already present in literature¹⁸ (see the Supporting Information for details), we were able to obtain ketone **32** in 46% yield by employing 1,3-dimethoxybenzene as the nucleophile, TMSOTf as the additive, and ibuprofen as the acyl fluoride precursor in a Friedel–Crafts-type acylation reaction.

In conclusion, we have developed a microfluidic reactor capable of safely producing SOF₂, an overlooked reagent due to its gaseous and toxic nature, starting from the commodity chemicals SOCl₂ and KF. This gas was generated *in situ* and reacted with a wide variety of carboxylic acids, including aromatic, aliphatic, and α -amino acid derivatives, to yield the corresponding acyl fluorides in a telescoped fashion. Furthermore, this method was capable of converting biorelevant molecules with perfect selectivity and good chemical yields. Finally, as acyl fluorides are attractive intermediates for the synthesis of other acyl derivatives, we devised a streamlined three-module flow setup, where, after the generation of the gas and the acyl fluoride production, the latter is directly reacted with a nucleophile to forge C–S, C–N, C–O, and C–C bonds. Based on these findings, we believe that this flow approach makes thionyl fluoride a convenient reagent to convert carboxylic acids into acyl fluorides. Further applications of this strategy are ongoing in our laboratories.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacsau.4c00318>.

Experimental procedures, optimization tables, photographs of the setup and scope limitations, characterization data (^1H , ^{13}C , and ^{19}F NMR, HRMS) for synthesized compounds, and further references provided by the authors. (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Timothy Noël – Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, 1098 XH Amsterdam, The Netherlands; orcid.org/0000-0002-3107-6927; Email: t.noel@uva.nl

Authors

Daniele Mazzarella – Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, 1098 XH Amsterdam, The Netherlands; Department of Chemical Sciences, University of Padova, 35131 Padova, Italy

Jelena Stanić – Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, 1098 XH Amsterdam, The Netherlands; orcid.org/0000-0001-8781-5330

Miguel Bernús – Departament de Química Analítica i Química Orgànica, Universitat Rovira i Virgili, 43007 Tarragona, Spain; orcid.org/0000-0003-0302-0720

Arad Seyed Mehdi – Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, 1098 XH Amsterdam, The Netherlands

Cassandra J. Henderson – Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, 1098 XH Amsterdam, The Netherlands; orcid.org/0000-0001-7402-6375

Omar Boutureira – Departament de Química Analítica i Química Orgànica, Universitat Rovira i Virgili, 43007 Tarragona, Spain; orcid.org/0000-0002-0768-8309

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacsau.4c00318>

Author Contributions

^{||}D.M. and J.S. contributed equally. CRediT: **Jelena Stanić** conceptualization, data curation, formal analysis.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors acknowledge financial support from the European Union H2020 research and innovation program for an ERC CoG grant for T.N. (FlowHAT, No. 101044355) and a Marie S. Curie Grant fellowship for D.M. (ELECTROORGANO, No. 101022144). Financial support from NWO via ENW-M2 grant is greatly appreciated (PC-Label, No. 14952, T.N. and C.J.H.). The authors are also grateful to the Spanish Government-MCIN, the National Agency of Investigation-AEI/10.13039/501100011033, and the European Regional Development

Fund-ERDF for project PID2020-120584RB-I00 to O.B., and FPU Fellowship (FPU19/01969 and EST22/00303) to M.B.

■ REFERENCES

- (1) (a) National Research Council. Committee on Prudent Practices in the Laboratory. In *Prudent Practices in the Laboratory: Handling and Management of Chemical Hazards: Updated Version*; National Academies Press: Washington, DC, 2011. (b) Schaufele, M. E. Toxic and flammable gases in research laboratories: Considerations for controls and continuous leak detection. *J. Chem. Health Saf.* **2013**, *20*, 8.
- (2) Leonard, K.; Lygo, B.; Procter, G. *Advanced Practical Organic Chemistry*; CRC Press: Boca Raton, FL, 2017.
- (3) (a) Demaerel, J.; Veryser, C.; De Borggraeve, W. M. *Ex situ* gas generation for lab scale organic synthesis. *React. Chem. Eng.* **2020**, *5*, 615. (b) Konishi, H. Creation of Novel Toxic Gas Surrogates and the Development of Safe and Facile Catalytic Reactions. *Chem. Pharm. Bull.* **2018**, *66*, 1. (c) Emmett, E. J.; Willis, M. C. The Development and Application of Sulfur Dioxide Surrogates in Synthetic Organic Chemistry. *Asian J. Org. Chem.* **2015**, *4*, 602. (d) Friis, S. D.; Lindhart, A. T.; Skrydstrup, T. The Development and Application of Two-Chamber Reactors and Carbon Monoxide Precursors for Safe Carbonylation Reactions. *Acc. Chem. Res.* **2016**, *49*, 594. (e) Chrominski, M.; Ziemkiewicz, K.; Kowalska, J.; Jemielity, J. Introducing SuFNucs: Sulfamoyl-Fluoride-Functionalized Nucleosides That Undergo Sulfur Fluoride Exchange Reaction. *Org. Lett.* **2022**, *24*, 4977.
- (4) (a) Woolven, H.; González-Rodríguez, C.; Marco, I.; Thompson, A. L.; Willis, M. C. DABCO-Bis(sulfur dioxide), DABSO, as a Convenient Source of Sulfur Dioxide for Organic Synthesis: Utility in Sulfonamide and Sulfamide Preparation. *Org. Lett.* **2011**, *13*, 4876. (b) Guo, T.; Meng, G.; Zhan, X.; Yang, Q.; Ma, T.; Xu, L.; Sharpless, K. B.; Dong, J. A New Portal to SuFEx Click Chemistry: A Stable Fluorosulfonyl Imidazolium Salt Emerging as an "F-SO₂+" Donor of Unprecedented Reactivity, Selectivity, and Scope. *Angew. Chem., Int. Ed.* **2018**, *57*, 2605. (c) Zhou, H.; Mukherjee, P.; Liu, R.; Evrard, E.; Wang, D.; Humphrey, J. M.; Butler, T. W.; Hoth, L. R.; Sperry, J. B.; Sakata, S. K.; Helal, C. J.; am Ende, C. W. Introduction of a Crystalline, Shelf-Stable Reagent for the Synthesis of Sulfur(VI) Fluorides. *Org. Lett.* **2018**, *20*, 812. (d) Ueda, T.; Konishi, H.; Manabe, K. Palladium-Catalyzed Reductive Carbonylation of Aryl Halides with *N*-Formylsaccharin as a CO Source. *Angew. Chem., Int. Ed.* **2013**, *52*, 8611.
- (5) (a) Bernus, M.; Mazzarella, D.; Stanić, J.; Zhai, Z.; Yeste-Vazquez, A.; Boutureira, O.; Gargano, A. F. G.; Grossmann, T. N.; Noël, T. A modular flow platform for sulfur(VI) fluoride exchange ligation of small molecules, peptides and proteins. *Nat. Synth.* **2024**, *3*, 185. (b) Raymenants, F.; Masson, T. M.; Sanjosé-Orduna, J.; Noël, T. Efficient C(sp³)-H Carbonylation of Light and Heavy Hydrocarbons with Carbon Monoxide via Hydrogen Atom Transfer Photocatalysis in Flow. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202308563. (c) Dong, Z.; Zondag, S. D. A.; Schmid, M.; Wen, Z.; Noël, T. A *meso*-scale ultrasonic milli-reactor enables gas-liquid-solid photocatalytic reactions in flow. *Chem. Eng. J.* **2022**, *428*, No. 130968. (d) Laudadio, G.; Deng, Y.; Van Der Wal, K.; Ravelli, D.; Nuño, M.; Fagnoni, M.; Guthrie, D.; Sun, Y.; Noël, T. C(sp³)-H functionalizations of light hydrocarbons using decatungstate photocatalysis in flow. *Science* **2020**, *369*, No. eabb4688. (e) Laudadio, G.; Govaerts, S.; Wang, Y.; Ravelli, D.; Koolman, H. F.; Fagnoni, M.; Djuric, S. W.; Noël, T. Selective C(sp³)-H Aerobic Oxidation Enabled by Decatungstate Photocatalysis in Flow. *Angew. Chem., Int. Ed.* **2018**, *57*, 4078. (f) Casnati, A.; Gemoets, H. P. L.; Motti, E.; Della Ca', N.; Noël, T. Homogeneous and Gas-Liquid Catalytic-Type Reaction Enabled by Continuous-Flow Chemistry. *Chem. – Eur. J.* **2018**, *24*, 14079. (g) Straathof, N. J. W.; Su, Y.; Hessel, V.; Noël, T. Accelerated gas-liquid visible light photoredox catalysis with continuous-flow photochemical microreactors. *Nat. Protoc.* **2016**, *11*, 10.
- (6) (a) Capaldo, L.; Wen, Z.; Noël, T. A field guide to flow chemistry for synthetic organic chemists. *Chem. Sci.* **2023**, *14*, 4230.

- (b) Plutschack, M. B.; Pieber, B.; Gilmore, K.; Seeberger, P. H. The Hitchhiker's Guide to Flow Chemistry. *Chem. Rev.* **2017**, *117*, 11796.
- (c) Wegner, J.; Ceylan, S.; Kirschnig, A. Flow Chemistry – A Key Enabling Technology for (Multistep) Organic Synthesis. *Adv. Synth. Catal.* **2012**, *354*, 17. (d) Jensen, K. F. Flow Chemistry—Microreaction Technology Comes of Age. *AIChE J.* **2017**, *63*, 858.
- (7) (a) Mallia, C. J.; Baxendale, I. R. The Use of Gases in Flow Synthesis. *Org. Process Res. Dev.* **2016**, *20*, 327. (b) Sivo, A.; Galaverna, R.; Gomes, G. R.; Pastre, J. C.; Vilé, G. From Circular Synthesis to Material Manufacturing: Advances, Challenges, and Future Steps for Using Flow Chemistry in Novel Application Area. *React. Chem. Eng.* **2021**, *6*, 756.
- (8) O'Brien, M.; Polyzos, A. *Flow Chemistry in Organic Synthesis*; Jamison, T. F.; Koch, G., Eds.; Wiley, 2018; pp 243–272.
- (9) (a) Dong, J.; Krasnova, L.; Finn, M. G.; Sharpless, K. B. Sulfur(VI) Fluoride Exchange (SuFEx): Another Good Reaction for Click Chemistry. *Angew. Chem., Int. Ed.* **2014**, *53*, 9430. (b) Barrow, A. S.; Smedley, C. J.; Zheng, Q.; Li, S.; Dong, J.; Moses, J. E. The growing applications of SuFEx click chemistry. *Chem. Soc. Rev.* **2019**, *48*, 4731.
- (10) (a) Bolduc, T. G.; Lee, C.; Chappell, W. P.; Sammis, G. M. Thionyl Fluoride-Mediated One-Pot Substitutions and Reductions of Carboxylic Acids. *J. Org. Chem.* **2022**, *87*, 7308. (b) Lee, C.; Thomson, B. J.; Sammis, G. Rapid and column-free syntheses of acyl fluorides and peptides using ex situ generated thionyl fluoride. *Chem. Sci.* **2021**, *13*, 188. (c) Thomson, B. J.; Khasnavis, S. R.; Grigorian, E. C.; Krishnan, R.; Yassa, T. D.; Lee, K.; Sammis, G. M.; Ball, N. D. Facile Synthesis of Sulfonyl Fluorides from Sulfonic Acids. *Chem. Commun.* **2023**, *59*, 555.
- (11) For a review on the topic: (a) Gonay, M.; Batisse, C.; Paquin, J. F. Recent Advances in the Synthesis of Acyl Fluorides. *Synthesis* **2021**, *53*, 653. For selected examples: (b) Olah, G. A.; Nojima, M.; Kerekes, I. Synthetic Methods and Reactions; IV. Fluorination of Carboxylic Acids with Cyanuric Fluoride. *Synthesis* **1973**, *8*, 487. (c) Markovskij, L. N.; Pashinnik, V. E.; Kirsanov, A. V. Application of Dialkylaminosulfur Trifluorides in the Synthesis of Fluoroorganic Compounds. *Synthesis* **1973**, *1973*, 787. (d) Olah, G. A.; Nojima, M.; Kerekes, I. Synthetic methods and reactions. I. Selenium tetrafluoride and its pyridine complex. Convenient fluorinating agents for fluorination of ketones, aldehydes, amides, alcohols, carboxylic acids, and anhydrides. *J. Am. Chem. Soc.* **1974**, *96*, 925. (e) Hasek, W. R.; Smith, W. C.; Engelhardt, V. A. The Chemistry of Sulfur Tetrafluoride. II. The Fluorination of Organic Carbonyl Compounds. *J. Am. Chem. Soc.* **1960**, *82*, 543. (f) Lal, G. S.; Pez, G. P.; Pesaresi, R. J.; Prozonc, F. M. Bis(2-methoxyethyl)aminosulfur trifluoride: a new broad-spectrum deoxofluorinating agent with enhanced thermal stability. *Chem. Commun.* **1999**, 215. (g) Scattolin, T.; Deckers, K.; Schoenebeck, F. Direct Synthesis of Acyl Fluorides from Carboxylic Acids with the Bench-Stable Solid Reagent (Me₄N)SCF₃. *Org. Lett.* **2017**, *19*, 5740. (h) Munoz, S. B.; Dang, H.; Isipzua-Rodriguez, X.; Mathew, T.; Surya Prakash, G. K. Synthesis of Acyl Fluorides via DAST-Mediated Fluorinative C–C Bond Cleavage of Activated Ketones. *Org. Lett.* **2019**, *21*, 1659. (i) Liang, Y.; Taya, A.; Zhao, Z.; Saito, N.; Shibata, N. Deoxyfluorination of acyl fluorides to trifluoromethyl compounds by FLUOLEAD/Olah's reagent under solvent-free conditions. *Beilstein J. Org. Chem.* **2020**, *16*, 3052. (j) Gonay, M.; Batisse, C.; Paquin, J. F. Synthesis of Acyl Fluorides from Carboxylic Acids Using NaF-Assisted Deoxofluorination with XtalFluor-E. *J. Org. Chem.* **2020**, *85*, 10253. (k) Wang, X.; Wang, F.; Huang, F.; Ni, C.; Hu, J. Deoxyfluorination of Carboxylic Acids with CpFluor: Access to Acyl Fluorides and Amides. *Org. Lett.* **2021**, *23*, 1764. (l) Liang, Y.; Zhao, Z.; Taya, A.; Shibata, N. Acyl Fluorides from Carboxylic Acids, Aldehydes, or Alcohols under Oxidative Fluorination. *Org. Lett.* **2021**, *23*, 847.
- (12) Ogiwara, Y.; Sakai, N. Acyl Fluorides in Late-Transition-Metal Catalysis. *Angew. Chem., Int. Ed.* **2020**, *59*, 574.
- (13) Nagano, S.; Maruoka, K. Synthesis of Acyl Fluorides from Carboxylic Acids with KI/AgSCF₃ for Efficient Amide and Peptide Synthesis. *Adv. Synth. Catal.* **2023**, *365*, 295.
- (14) Trynieszewski, M.; Barbasiewicz, M. Gram-Scale Preparation of Acyl Fluorides and Their Reactions with Hindered Nucleophiles. *Synthesis* **2022**, *54*, 1446.
- (15) (a) Jiao, J.; Nie, W.; Yu, T.; Yang, F.; Zhang, Q.; Aihemaiti, F.; Yang, T.; Liu, X.; Wang, J.; Li, P. Multi-Step Continuous-Flow Organic Synthesis: Opportunities and Challenges. *Chem. – Eur. J.* **2021**, *27*, 4817. (b) Britton, J.; Raston, C. L. Multi-step continuous-flow synthesis. *Chem. Soc. Rev.* **2017**, *46*, 1250.
- (16) Jiang, L.; Wu, B.; Song, Y.; Yang, M.; Wang, D.; Liu, Y.; Xue, Z. Mass transfer coefficient measurement during brine flush in a CO₂-filled packed bed by X-ray CT scanning. *Int. J. Heat Mass Transfer* **2017**, *115*, 615.
- (17) Lal, G. S.; Pez, G. P.; Pesaresi, R. J.; Prozonc, F. M.; Cheng, H. Bis(2-methoxyethyl)aminosulfur Trifluoride: A New Broad-Spectrum Deoxofluorinating Agent with Enhanced Thermal Stability. *J. Org. Chem.* **1999**, *64*, 7048.
- (18) Raghavendra Rao, K.; Valleé, Y. Friedel–Crafts reactions of acyl trifluoromethanesulfonates and cyclic acylsulfonium cations generated from acyl fluorides. *Tetrahedron* **2016**, *72*, 4442.