

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Diimidazolium phthalate monohydrate

Chua-Hua Yu

Ordered Matter Science Research Center, Southeast University, Nanjing 211189, People's Republic of China

Correspondence e-mail: jxyuchunhua@163.com


Received 29 May 2012; accepted 24 June 2012

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.058; wR factor = 0.141; data-to-parameter ratio = 17.1.

In the title compound, $2\text{C}_3\text{H}_5\text{N}_2^+ \cdot \text{C}_8\text{H}_4\text{O}_4^{2-} \cdot \text{H}_2\text{O}$, the cations, anion and water molecule are connected by $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds, forming a three-dimensional network.

Related literature

The title compound was synthesized during a search for ferroelectric materials. For background to ferroelectric organic materials with framework structures, see: Zhang *et al.* (2009, 2010); Zhang & Xiong (2012). For related structures, see: Yu & Zhu (2012); Zhu & Yu (2011).



Experimental

Crystal data

 $2\text{C}_3\text{H}_5\text{N}_2^+ \cdot \text{C}_8\text{H}_4\text{O}_4^{2-} \cdot \text{H}_2\text{O}$
 $M_r = 320.31$

 Monoclinic, $P2_1/n$
 $a = 9.1250$ (18) Å

 $b = 12.979$ (3) Å

 $c = 13.549$ (3) Å

 $\beta = 103.85$ (3)°

 $V = 1558.0$ (6) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.11$ mm⁻¹
 $T = 293$ K

 $0.32 \times 0.28 \times 0.26$ mm

Data collection

Rigaku SCXmini diffractometer

Absorption correction: multi-scan

 (*CrystalClear*; Rigaku, 2005)

 $T_{\min} = 0.967$, $T_{\max} = 0.973$

15753 measured reflections

3567 independent reflections

 2758 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

3 standard reflections every 180 reflections

intensity decay: none

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.141$
 $S = 1.10$

3567 reflections

208 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.32$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.50$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1A} \cdots \text{O1}$	0.96	1.71	2.665 (2)	170
$\text{N2}-\text{H2A} \cdots \text{O2}^{\text{i}}$	0.91	1.75	2.655 (2)	172
$\text{N3}-\text{H3A} \cdots \text{O3}$	0.95	1.80	2.747 (2)	177
$\text{N3}-\text{H3A} \cdots \text{O4}$	0.95	2.56	3.191 (2)	124
$\text{N4}-\text{H4A} \cdots \text{O3}^{\text{ii}}$	0.86	1.93	2.774 (2)	168
$\text{O1W}-\text{H1WA} \cdots \text{O4}$	0.86	1.98	2.834 (2)	170
$\text{O1W}-\text{H1WB} \cdots \text{O2}^{\text{iii}}$	0.90	1.94	2.8326 (19)	172

 Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005); software used to prepare material for publication: *SHELXL97*.

The author thanks the Ordered Matter Science Research Center, Southeast University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2058).

References

- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yu, C.-H. & Zhu, R.-Q. (2012). *Acta Cryst.* **E68**, o1911.
- Zhang, W., Chen, L.-Z., Xiong, R.-G., Nakamura, T. & Huang, S.-P. (2009). *J. Am. Chem. Soc.* **131**, 12544–12545.
- Zhang, W. & Xiong, R.-G. (2012). *Chem. Rev.* **112**, 1163–1195.
- Zhang, W., Ye, H.-Y., Cai, H.-L., Ge, J.-Z., Xiong, R.-G. & Huang, S.-P. D. (2010). *J. Am. Chem. Soc.* **132**, 7300–7302.
- Zhu, R.-Q. & Yu, C.-H. (2011). *Acta Cryst.* **E67**, o2746.

supplementary materials

Acta Cryst. (2012). E68, o2295 [doi:10.1107/S1600536812028619]

Diimidazolium phthalate monohydrate

Chua-Hua Yu

Comment

In our search for potential ferroelectric phase change materials, the title compound was synthesized. This search is carried out by measurement of the dielectric constant of compounds on the basis of temperature, for example, (Zhang, Chen *et al.*, 2009; Zhang, Ye *et al.*, 2010; Zhang & Xiong, 2012), this has been carried out for $C_3H_5N_2^+.C_2HO_4^-$ (Yu & Zhu, 2012) and $C_3H_9N_2^+.C_8H_5O_4^-$ (Zhu & Yu, 2011). In the case of the title compound no dielectric anomaly was observed ranging from 130 K to 375 K.

The crystal structure of the title compound contains two protonated imidazolium cations, one phthalate anion, losing two H atoms, and one molecule of water molecule is shown in Fig. 1. The asymmetric unit was selected with the cations, anion and water molecule connected by the intramolecular hydrogen bonds, N1–1A \cdots O1, N3–H3A \cdots O3, N3–H3A \cdots O4 and O1W–H1WA \cdots O4 all of which connect the cations and the water molecule to the anion, Table 1. These units are connected by the intermolecular hydrogen bonds, N2–H2A \cdots O2(-1+x,y,z), N4–H4A \cdots O3(-x+1/2,y-1/2,-z+1/2) and O1W–H1WB \cdots O2(-x+2,-y+1,-z+1), to form a three-dimensional network Table2 and Figure 1.

Experimental


0.83 g (5 mmol) of phthalic acid was added to in 10 ml water which was heated. A few drops of ethanol and 0.68 g (10 mmol) imidazole were added to the solution. The mixture was stirred until it reached ambient temperature, the liquid was filtered to give a clear solution. Colourless crystals suitable for X-ray structure analysis were obtained by the slow evaporation of the solution after several days at the ambient temperature.

Refinement


H atoms attached to C were placed in calculated positions (C–H = 0.93 Å for C_{sp2} atoms) while those attached to N and O were found in positions derived from a difference electron density map, with $U_{iso}(H) = 1.2 U_{iso}(C, N)$, $U_{iso}(H) = 1.5 U_{iso}(O)$. The -2 0 2 reflection was omitted since its measured value appear to be anomalous.

Computing details

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear* (Rigaku, 2005); data reduction: *CrystalClear* (Rigaku, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).


Figure 1

A diagram of the title compound, with displacement ellipsoids drawn at the 30% probability level. The atomic numbering scheme is shown.


Figure 2

A view of the packing diagram of the title compound, stacking along the *b* axis. Hydrogen bonds are shown as dashed lines.

Diimidazolium phthalate monohydrate

Crystal data



$M_r = 320.31$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 9.1250(18)\ \text{\AA}$

$b = 12.979(3)\ \text{\AA}$

$c = 13.549(3)\ \text{\AA}$

$\beta = 103.85(3)^\circ$

$V = 1558.0(6)\ \text{\AA}^3$

$Z = 4$

$F(000) = 672$

$D_x = 1.366\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3567 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 0.11\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colourless

$0.32 \times 0.28 \times 0.26\ \text{mm}$

Data collection

Rigaku SCXmini diffractometer	3567 independent reflections
Radiation source: fine-focus sealed tube	2758 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.044$
ω scans	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.967$, $T_{\text{max}} = 0.973$	$k = -16 \rightarrow 16$
15753 measured reflections	$l = -17 \rightarrow 17$
	3 standard reflections every 180 reflections
	intensity decay: none

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.058$	H-atom parameters constrained
$wR(F^2) = 0.141$	$w = 1/[\sigma^2(F_o^2) + (0.0667P)^2 + 0.3812P]$
$S = 1.10$	where $P = (F_o^2 + 2F_c^2)/3$
3567 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
208 parameters	$\Delta\rho_{\text{max}} = 0.32 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.50 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.67631 (14)	0.59458 (9)	0.53305 (10)	0.0375 (3)
O2	0.92261 (14)	0.61977 (10)	0.57079 (11)	0.0423 (3)
O3	0.45616 (13)	0.66464 (9)	0.28714 (10)	0.0386 (3)
O4	0.67947 (15)	0.59102 (10)	0.30286 (12)	0.0466 (4)
C1	0.66990 (17)	0.75577 (12)	0.37728 (12)	0.0242 (3)
C2	0.64193 (19)	0.85209 (13)	0.33196 (13)	0.0310 (4)
H2	0.5809	0.8575	0.2665	0.037*
C3	0.7038 (2)	0.93997 (13)	0.38309 (14)	0.0385 (4)
H3	0.6833	1.0041	0.3523	0.046*
C4	0.7959 (2)	0.93247 (13)	0.47976 (15)	0.0417 (5)
H4	0.8372	0.9915	0.5144	0.050*
C5	0.8267 (2)	0.83681 (14)	0.52503 (13)	0.0353 (4)
H5	0.8898	0.8319	0.5899	0.042*
C6	0.76452 (17)	0.74796 (12)	0.47483 (12)	0.0254 (3)
C7	0.79001 (18)	0.64612 (13)	0.52933 (12)	0.0266 (4)
C8	0.59838 (19)	0.66201 (12)	0.31870 (12)	0.0275 (4)

N1	0.41475 (17)	0.68481 (12)	0.53530 (12)	0.0390 (4)
H1A	0.5080	0.6559	0.5268	0.047*
N2	0.18863 (17)	0.69028 (12)	0.55394 (12)	0.0375 (4)
H2A	0.0940	0.6703	0.5552	0.045*
C9	0.2907 (2)	0.63160 (14)	0.52810 (15)	0.0385 (4)
H9	0.2770	0.5631	0.5079	0.046*
C10	0.2510 (2)	0.78435 (16)	0.58006 (18)	0.0499 (5)
H10	0.2043	0.8406	0.6021	0.060*
C11	0.3926 (2)	0.78088 (16)	0.5680 (2)	0.0548 (6)
H11	0.4625	0.8344	0.5799	0.066*
N3	0.37146 (18)	0.46491 (12)	0.23556 (11)	0.0375 (4)
H3A	0.4030	0.5339	0.2521	0.045*
N4	0.24322 (18)	0.32527 (12)	0.20991 (12)	0.0403 (4)
H4A	0.1710	0.2820	0.2070	0.048*
C12	0.2391 (2)	0.42411 (14)	0.23263 (15)	0.0398 (4)
H12	0.1560	0.4590	0.2446	0.048*
C13	0.3841 (2)	0.30217 (16)	0.19825 (16)	0.0446 (5)
H13	0.4179	0.2383	0.1820	0.053*
C14	0.4641 (2)	0.39014 (16)	0.21490 (16)	0.0452 (5)
H14	0.5645	0.3983	0.2127	0.054*
O1W	0.99551 (16)	0.57033 (11)	0.32839 (11)	0.0479 (4)
H1WA	0.9020	0.5841	0.3250	0.072*
H1WB	1.0210	0.5131	0.3660	0.072*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0274 (6)	0.0290 (6)	0.0568 (8)	-0.0006 (5)	0.0116 (6)	0.0092 (6)
O2	0.0250 (6)	0.0430 (8)	0.0569 (8)	0.0060 (6)	0.0058 (6)	0.0199 (6)
O3	0.0273 (7)	0.0295 (7)	0.0526 (8)	-0.0025 (5)	-0.0032 (6)	-0.0047 (6)
O4	0.0372 (8)	0.0359 (7)	0.0664 (9)	0.0010 (6)	0.0118 (7)	-0.0221 (6)
C1	0.0216 (7)	0.0219 (8)	0.0291 (8)	-0.0012 (6)	0.0060 (6)	-0.0019 (6)
C2	0.0316 (9)	0.0288 (9)	0.0295 (9)	0.0003 (7)	0.0015 (7)	0.0029 (7)
C3	0.0500 (12)	0.0205 (8)	0.0423 (10)	-0.0005 (8)	0.0057 (8)	0.0041 (7)
C4	0.0529 (12)	0.0236 (9)	0.0434 (11)	-0.0073 (8)	0.0016 (9)	-0.0066 (7)
C5	0.0380 (10)	0.0323 (9)	0.0307 (9)	-0.0044 (8)	-0.0013 (7)	-0.0039 (7)
C6	0.0219 (8)	0.0242 (8)	0.0299 (8)	0.0000 (6)	0.0056 (6)	0.0003 (6)
C7	0.0257 (8)	0.0252 (8)	0.0293 (8)	0.0029 (7)	0.0075 (6)	0.0009 (6)
C8	0.0265 (8)	0.0255 (8)	0.0288 (8)	-0.0020 (7)	0.0037 (6)	-0.0015 (6)
N1	0.0283 (8)	0.0405 (9)	0.0500 (9)	0.0023 (7)	0.0129 (7)	0.0026 (7)
N2	0.0290 (8)	0.0320 (8)	0.0522 (9)	-0.0025 (6)	0.0113 (7)	-0.0003 (7)
C9	0.0351 (10)	0.0295 (9)	0.0501 (11)	0.0001 (8)	0.0083 (8)	-0.0045 (8)
C10	0.0416 (12)	0.0293 (10)	0.0811 (16)	-0.0004 (9)	0.0194 (11)	-0.0093 (10)
C11	0.0416 (12)	0.0357 (11)	0.0872 (17)	-0.0119 (9)	0.0155 (11)	-0.0056 (11)
N3	0.0434 (9)	0.0313 (8)	0.0373 (8)	-0.0099 (7)	0.0088 (7)	-0.0018 (6)
N4	0.0394 (9)	0.0317 (8)	0.0490 (10)	-0.0117 (7)	0.0088 (7)	-0.0040 (7)
C12	0.0403 (11)	0.0346 (10)	0.0458 (11)	-0.0065 (8)	0.0126 (9)	-0.0049 (8)
C13	0.0446 (11)	0.0375 (10)	0.0508 (12)	0.0022 (9)	0.0100 (9)	-0.0068 (9)
C14	0.0340 (10)	0.0504 (12)	0.0527 (12)	-0.0067 (9)	0.0131 (9)	-0.0013 (10)
O1W	0.0393 (8)	0.0419 (8)	0.0624 (10)	0.0004 (6)	0.0116 (7)	0.0079 (7)

Geometric parameters (Å, °)

O1—C7	1.246 (2)	N2—C9	1.314 (2)
O2—C7	1.253 (2)	N2—C10	1.358 (2)
O3—C8	1.266 (2)	N2—H2A	0.9055
O4—C8	1.232 (2)	C9—H9	0.9300
C1—C2	1.389 (2)	C10—C11	1.342 (3)
C1—C6	1.399 (2)	C10—H10	0.9300
C1—C8	1.513 (2)	C11—H11	0.9300
C2—C3	1.383 (2)	N3—C12	1.310 (2)
C2—H2	0.9300	N3—C14	1.360 (3)
C3—C4	1.380 (3)	N3—H3A	0.9507
C3—H3	0.9300	N4—C12	1.322 (2)
C4—C5	1.384 (3)	N4—C13	1.365 (3)
C4—H4	0.9300	N4—H4A	0.8591
C5—C6	1.389 (2)	C12—H12	0.9300
C5—H5	0.9300	C13—C14	1.345 (3)
C6—C7	1.505 (2)	C13—H13	0.9300
N1—C9	1.309 (2)	C14—H14	0.9300
N1—C11	1.355 (3)	O1W—H1WA	0.8623
N1—H1A	0.9615	O1W—H1WB	0.8997
C2—C1—C6	119.30 (14)	C9—N2—H2A	125.4
C2—C1—C8	118.78 (14)	C10—N2—H2A	126.5
C6—C1—C8	121.91 (14)	N1—C9—N2	109.25 (16)
C3—C2—C1	120.73 (15)	N1—C9—H9	125.4
C3—C2—H2	119.6	N2—C9—H9	125.4
C1—C2—H2	119.6	C11—C10—N2	107.09 (18)
C4—C3—C2	120.01 (16)	C11—C10—H10	126.5
C4—C3—H3	120.0	N2—C10—H10	126.5
C2—C3—H3	120.0	C10—C11—N1	107.11 (18)
C3—C4—C5	119.80 (16)	C10—C11—H11	126.4
C3—C4—H4	120.1	N1—C11—H11	126.4
C5—C4—H4	120.1	C12—N3—C14	108.62 (16)
C4—C5—C6	120.81 (16)	C12—N3—H3A	127.8
C4—C5—H5	119.6	C14—N3—H3A	123.5
C6—C5—H5	119.6	C12—N4—C13	108.66 (16)
C5—C6—C1	119.34 (15)	C12—N4—H4A	125.5
C5—C6—C7	119.47 (14)	C13—N4—H4A	125.7
C1—C6—C7	121.02 (14)	N3—C12—N4	108.79 (18)
O1—C7—O2	124.02 (15)	N3—C12—H12	125.6
O1—C7—C6	117.34 (14)	N4—C12—H12	125.6
O2—C7—C6	118.61 (14)	C14—C13—N4	106.43 (18)
O4—C8—O3	124.83 (15)	C14—C13—H13	126.8
O4—C8—C1	119.42 (15)	N4—C13—H13	126.8
O3—C8—C1	115.72 (14)	C13—C14—N3	107.50 (18)
C9—N1—C11	108.42 (17)	C13—C14—H14	126.2
C9—N1—H1A	124.0	N3—C14—H14	126.2
C11—N1—H1A	127.1	H1WA—O1W—H1WB	108.6
C9—N2—C10	108.12 (16)		

C6—C1—C2—C3	-1.4 (3)	C2—C1—C8—O4	122.83 (18)
C8—C1—C2—C3	179.38 (16)	C6—C1—C8—O4	-56.3 (2)
C1—C2—C3—C4	0.8 (3)	C2—C1—C8—O3	-55.5 (2)
C2—C3—C4—C5	0.3 (3)	C6—C1—C8—O3	125.37 (17)
C3—C4—C5—C6	-0.7 (3)	C11—N1—C9—N2	0.7 (2)
C4—C5—C6—C1	0.1 (3)	C10—N2—C9—N1	-0.9 (2)
C4—C5—C6—C7	-175.09 (17)	C9—N2—C10—C11	0.7 (3)
C2—C1—C6—C5	1.0 (2)	N2—C10—C11—N1	-0.3 (3)
C8—C1—C6—C5	-179.85 (16)	C9—N1—C11—C10	-0.2 (3)
C2—C1—C6—C7	176.10 (15)	C14—N3—C12—N4	-0.5 (2)
C8—C1—C6—C7	-4.7 (2)	C13—N4—C12—N3	0.2 (2)
C5—C6—C7—O1	125.27 (18)	C12—N4—C13—C14	0.1 (2)
C1—C6—C7—O1	-49.8 (2)	N4—C13—C14—N3	-0.4 (2)
C5—C6—C7—O2	-52.5 (2)	C12—N3—C14—C13	0.6 (2)
C1—C6—C7—O2	132.45 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O1	0.96	1.71	2.665 (2)	170
N2—H2A \cdots O2 ⁱ	0.91	1.75	2.655 (2)	172
N3—H3A \cdots O3	0.95	1.80	2.747 (2)	177
N3—H3A \cdots O4	0.95	2.56	3.191 (2)	124
N4—H4A \cdots O3 ⁱⁱ	0.86	1.93	2.774 (2)	168
O1W—H1WA \cdots O4	0.86	1.98	2.834 (2)	170
O1W—H1WB \cdots O2 ⁱⁱⁱ	0.90	1.94	2.8326 (19)	172

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1/2, y-1/2, -z+1/2$; (iii) $-x+2, -y+1, -z+1$.