

RESEARCH

Open Access



Sexual transmission of *Anopheles gambiae* densovirus (AgDENV) leads to disseminated infection in mated females

Kristine L. Werling^{1,2}, Rebecca M. Johnson^{1,2,4}, Hillery C. Metz^{1,2} and Jason L. Rasgon^{1,2,3*}

Abstract

Background: *Anopheles gambiae* densovirus (AgDENV) is an insect-specific, single-stranded DNA virus that infects *An. gambiae* sensu stricto (s.s.), the major mosquito species responsible for transmitting malaria parasites throughout sub-Saharan Africa. AgDENV is a benign virus that is very specific to its mosquito host and therefore has the potential to serve as a vector control tool via paratransgenesis (genetic modification of mosquito symbionts) to limit transmission of human pathogens. Prior to being engineered into a control tool, the natural transmission dynamics of AgDENV between *An. gambiae* mosquitoes needs to be fully understood. Additionally, improved knowledge of AgDENV infection in male mosquitoes is needed. In the study presented here, we examined the tissue tropism of AgDENV in the male reproductive tract and investigated both venereal and vertical transmission dynamics of the virus.

Methods: *Anopheles gambiae* s.s. adult males were infected with AgDENV via microinjection, and reproductive tissues were collected and assayed for AgDENV using qPCR. Next, uninfected females were introduced to AgDENV-infected or control males and, after several nights of mating, both the spermatheca and female carcass were assessed for venereally transmitted AgDENV. Finally, F1 offspring of this cross were collected and assayed to quantify vertical transmission of the virus.

Results: AgDENV infected the reproductive tract of male mosquitoes, including the testes and male accessory glands, without affecting mating rates. AgDENV-infected males venereally transmitted the virus to females, and these venereally infected females developed disseminated infection throughout the body. However, AgDENV was not vertically transmitted to the F1 offspring of this cross.

Conclusions: Infected male releases could be an effective strategy to introduce AgDENV-based paratransgenic tools into naïve populations of *An. gambiae* s.s. females.

Keywords: Insect specific virus, Densovirus, *Anopheles gambiae*, Paratransgenesis, Male releases

Background

Anopheles gambiae densovirus (AgDENV) is an insect-specific virus (ISV) that efficiently infects *An. gambiae* sensu stricto (s.s.) mosquitoes [1, 2], one of the major

vectors of malaria-causing parasites. This non-enveloped, single-stranded DNA virus belongs to the *Parvoviridae* family and *Brevidensovirus* genus and has a compact genome of approximately 4.1 kb [1]. First identified in an *An. gambiae* cell line (Sua5B) [1], AgDENV is specific to *An. gambiae* s.s. mosquitoes [2] and can also likely infect the sibling species *Anopheles coluzzii* (due to the mixed origin of Sua5B cells). It replicates only minimally in closely related *Anopheles arabiensis* and cannot replicate in more distantly related *Anopheles* or other mosquito

*Correspondence: jlr54@psu.edu

¹ Department of Entomology, Pennsylvania State University, University Park, PA, USA

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

genera [2]. Importantly, AgDENV is also unable to infect vertebrates [2]. Due to this host specificity, ISVs like AgDENV offer a promising new avenue for targeted biological control of individual mosquito species and the pathogens they transmit [3–5]. In its natural *An. gambiae* s.s. host (hereinafter referred to as *An. gambiae*), AgDENV can infect both sexes and all developmental stages, replicating particularly well in adults post-emergence [6, 7]. Notably, AgDENV is a benign virus that does not cause mortality or even a marked transcriptional response in its mosquito host [8]. This is in contrast to several related mosquito-specific densoviruses that infect *Aedes* mosquitoes with high lethality [9–11]. While cytotoxic densoviruses can be harnessed to make novel biopesticides [5], as was done with the *Aedes aegypti* densovirus (AaeDENV) [12], the benign nature of AgDENV is highly advantageous for another vector control approach known as paratransgenesis, or the genetic modification of symbionts with the goal of decreasing pathogen transmission. Specifically, AgDENV could be genetically modified to encode a specific effector that, upon delivery and expression in its *An. gambiae* host, targets either the mosquito or the pathogens they transmit to ultimately decrease the incidence of mosquito-borne diseases like malaria. The potential use of AgDENV for paratransgenesis has previously been demonstrated [1, 7, 13], but for an AgDENV-based tool to be effective in field settings, we first need a complete understanding of AgDENV transmission dynamics between *An. gambiae* hosts. This information is critical for determining the best application and deployment strategy for an AgDENV-based vector control tool.

A previous study found that AgDENV can be venereally transmitted from infected males to the spermatheca of females post-mating [6], yet it is not known if venereally transmitted AgDENV remains contained to the spermatheca or if it can also lead to disseminated infection in females. This information is important for determining whether infected male releases could be a viable method for introducing AgDENV-based tools into field populations of female *An. gambiae*. As only female mosquitoes blood feed and transmit disease-causing pathogens, they are the main target of vector control programs, and as such, male mosquitoes can be safely released into a population without directly affecting pathogen transmission. Modified male releases is an approach that has been successful for certain *Aedes aegypti* control programs using either the endosymbiont bacterium *Wolbachia* [14] or transgenic male mosquitoes [15, 16]. In addition to venereal transmission, AgDENV can also be vertically transmitted from mother to offspring when the mother is infected during larval stages [1], but it is not yet known if mothers infected venereally can subsequently transmit AgDENV to the next generation. This knowledge would

reveal if AgDENV-based tools could disseminate transgenerationally into a population following infected male releases.

We also do not know the tissue tropism of AgDENV in male mosquitoes, particularly in their reproductive tissues where infection could impact mating success. Notably, *plugin* is one of the few genes that is differentially expressed in AgDENV-infected mosquitoes [8]. It encodes a fundamental component of the male mating plug, a coagulated mass of seminal secretions transferred to females during copulation that is essential for proper sperm storage in females [17]. Given that AgDENV influences the expression of this key mating factor, assessing whether AgDENV infection influences male mating rates is also an urgent question, as this could impinge on the success of AgDENV-based tools.

In the study reported here, we addressed these questions by examining AgDENV tropism and transmission dynamics in mosquitoes infected in the laboratory. We found that AgDENV spreads venereally from infected males to naïve females, such that females develop disseminated infection; however, AgDENV is not subsequently transmitted vertically to F1 offspring. Our findings demonstrate that AgDENV holds promise as a tool for paratransgenesis in *An. gambiae*, whereby infected male releases could be utilized for dissemination into field populations.

Methods

Production of AgDENV (GenBank: EU233812.1)

Moss55 cells, which naturally lack wild-type AgDENV [1], were transfected with a plasmid containing the complete wild-type genome of AgDENV, as previously described [1, 6, 13]. Briefly, 6-well plates were seeded with approximately 3×10^6 cells per well. The following day, transfections were performed using Lipofectamine LTX with Plus Reagent (Thermo Fisher Scientific, Waltham, MA, USA) and 2.5 μ g of AgDENV-containing plasmid (pAgDENV) per well. Three days after transfection, AgDENV virus was harvested as follows: cells were washed and re-suspended in phosphate-buffered saline (PBS), lysed via 3 freeze/thaw cycles (alternating -80°C and room temperature) and then spun down to remove cell debris. Supernatant was collected, aliquoted and stored at -80°C . This supernatant served as the AgDENV viral inoculum for all subsequent experiments.

Mosquito rearing

Anopheles gambiae s.s. (Keele strain) were reared at 27°C and approximately 80% humidity under a 12/12-h light/dark cycle. Larvae were fed ground Tetramin flakes (Tetra Werke, Melle, Germany) daily. At the pupal stage, males

and females were separated to maintain virgin adults. Adults were provided 10% sucrose solution throughout.

Microinjections

At 2–3 days post-emergence, virgin males were interthoracically injected with approximately 200 nl of either AgDENV viral stock or PBS (negative control) using a Nanoject II Auto-Nanoliter Injector (Drummond Scientific Co., Broomall, PA, USA) and glass capillary needles back-filled with mineral oil. Mosquitoes injected with AgDENV each received an estimated 10^6 viral genome equivalents (vge). Mosquitoes were cold-anesthetized using ice and a cold-block during injections, and immediately placed to recover at room temperature after being injected.

Mating

When AgDENV- and PBS-injected male mosquitoes reached 14 days post-injection (dpi), virgin females were introduced into their cages using a mouth aspirator (John W Hock Co., Gainesville, FL, USA). Females were added in a 1:2 ratio to males, and these mixed cages were left for 3 nights to allow for mating, following which the mixed sex cage was briefly chilled at 4 °C to anesthetize the mosquitoes. Males and females were then separated over ice using a paintbrush and forceps and transferred to new cages.

Blood-feeding and oviposition for F1 offspring

Female mosquitoes were provided with a blood meal at 1 and 7 days post-mating. Blood-feeding was achieved using a 37 °C water bath, glass membrane feeder, and anonymous human blood sourced from BioIVT (Westbury, NY, USA). Two days after each blood-feeding, females were provided with an oviposition site (petri dish with wet cotton and filter paper). Eggs were collected 3 days later and immediately hatched and reared as described above.

Tissue collections

Dissected tissues were either stored dry (carcasses) or collected in 50–100 µl of PBS (spermathecas, testes, male accessory glands [MAGs]) and stored at – 80 °C until DNA extraction.

Males Male reproductive tissues (testes and MAGs) were collected in pools of 12, while male carcasses (body without reproductive tract) were analyzed individually. Males were analyzed at 11, 13, 15 and 18 dpi.

Females Female mating status was determined by visual examination of the spermatheca for the presence or absence of sperm. Only mated females were further processed for AgDENV quantification. Spermathecas

were pooled into groups of 5–13 tissues, while female carcasses (body without spermatheca) were analyzed individually. Females were analyzed at 8–14 days post-mating (estimated, because exact time of mating was not known).

F1 offspring F1 offspring were collected as whole mosquitoes, and both males and females were analyzed in groups of four to five individuals. F1 adults were maintained in a mixed sex cage until collection.

DNA extraction from mosquito tissues

Tissues were first homogenized using either sterile pellet pestles (Fisherbrand™, Thermo Fisher Scientific) and an electric pestle grinder (Kimble®; Dwk Life Sciences LLC, Millville, NJ, USA) (spermathecas, testes, MAGs), or stainless steel beads (OPS Diagnostics, Lebanon, NJ, USA) and the TissueLyser II small bead mill (Qiagen, Hilden, Germany) (carcasses). After homogenization, TL buffer (Omega Bio-Tek Inc., Norcross, GA, USA) was added to reach a final volume of 200 µl. DNA was then extracted from the samples using the E.Z.N.A Tissue DNA kit (Omega Bio-Tek) according to the manufacturer's instructions. Proteinase K incubation was carried out for 60 min at 55 °C in a water bath with vortexing every 15–20 min, and final samples were eluted twice with 50 µl elution buffer warmed to 70 °C. Sample quantity and quality were checked using a Nanodrop spectrophotometer (Thermo Fisher Scientific) before proceeding to quantitative PCR (qPCR).

Quantification of AgDENV

Quantification of AgDENV viral stock produced from transfected cells One aliquot of virus stock was removed from storage at – 80 °C and thawed on ice. Using Turbo™ DNase (Invitrogen™, Thermo Fisher Scientific) the sample was treated to remove any remaining plasmid DNA that may have been left over from the transfection and would interfere with viral quantification. Next, viral DNA was extracted from the stock using the E.Z.N.A Tissue DNA kit (Omega Bio-Tek) according to the manufacturer's instructions for cultured cells, starting with Proteinase K treatment. qPCR was then run on a Roto-Gene Q real-time cyler (Qiagen) using the PerfeCTa SYBR Green Master Mix (Quantabio, Beverly, MA, USA) and primers targeting the viral protein (VP) gene on the AgDENV genome. Samples were run alongside a standard curve, consisting of a pAgDENV dilution series, to allow for quantification of vge per milliliter. VP primer sequences were the same as those used in a previous study [22]: 5'-GGC ATC AAT GTG GGA CCA

AG-3' (forward) and 5'-CCG TTA GCA AGC GTT GTC TG-3' (reverse). qPCR cycling conditions were an initial cycle of 95 °C for 5 min, followed by 40 cycles of 95 °C for 10 s, then by a data acquisition step of 60 °C for 30 s; a melt curve analysis was run at the end of cycling by going from 72 °C to 95 °C with 5 s per step.

Quantifying AgDENV from mosquito tissues Following DNA extraction, samples were analyzed by qPCR, as described above, except that a pAgDENV standard curve was not run. Instead, primers targeting the host ribosomal protein gene *S7* were run alongside primers targeting the viral VP gene. Relative AgDENV titers were then determined by normalizing VP quantification (viral genomes) to *S7* quantification (host genomes). This step normalizes for differences in the amount of total DNA obtained from different tissue types (e.g. testes vs. carcass). *S7* primer sequences were: 5'-AAG GGT TGC GTG CTA GTG AA-3' (forward) and 5'-TAA CGG CTT TTC TGC GTC CA-3' (reverse). AgDENV dissemination rates in mated female carcasses (see text associated with Fig. 2b) were estimated by using the highest detectable VP/*S7* value found in control female carcasses (0.0000845 VP/*S7*) as the cut-off for determining AgDENV-positive ($n=45$, 90%) or AgDENV-negative samples ($n=5$, 10%).

Statistical analyses

Data for relative AgDENV titers were first tested for normality (Kolmogorov–Smirnov, D’Agostino & Pearson) and then, where appropriate, log [Ln(y)] transformed to achieve normality. When analyzing two groups, unpaired Student t-tests were used for data with equal variances,

and Welch t-tests were used for data with unequal variances. If analyzing more than two groups, one-way analysis of variance (ANOVA) (equal variances) or Brown–Forsythe/Welch ANOVA (unequal variances) was used with either Sidak’s or Dunnett’s multiple testing correction, respectively. Mating and survival rates were analyzed using Fisher’s exact tests. Statistical tests are also indicated in the figure and table legends. A *P*-value cut-off of <0.05 was used to determine significance. Analyses were performed with either GraphPad Prism (GraphPad Software Inc., San Diego, CA, USA) or JMP software (SAS Institute, Cary, NC, USA).

Results

AgDENV infects male reproductive tissues without affecting mating rates

To determine the tissue tropism and infection dynamics of AgDENV in the reproductive tract of *An. gambiae* males, we infected young adult males with AgDENV via inter-thoracic microinjections and then assessed viral titers at four time points, relative to PBS-injected controls. We found that AgDENV infects both major male reproductive tissues, namely the testes (Fig. 1a) and the male accessory glands (MAGs) (Fig. 1b), as well as the remainder of the body (hereafter referred to as the carcass) (Fig. 1c). AgDENV was detectable in all tissues at 11, 13, 15, and 18 dpi, yet within each tissue type, relative viral titers (i.e. normalized to host genome) did not differ significantly between time points (Fig. 1a–c). Independent of time post-injection, we found significantly higher relative AgDENV titers in the MAGs compared to the testes (Fig. 1d), with MAG titers comparable to those of the carcass. We summarize these results as showing




Fig. 1 AgDENV infects the reproductive tract of *Anopheles gambiae* males. **a–c** Males were injected with either AgDENV (DNV on figure) or PBS (control). **a–c** At 11, 13, 15 and 18 dpi, AgDENV-treated males have infected: **a** testes [Ln(y) transformed, one-way ANOVA, Sidak’s correction], **b** MAGs [Ln(y) transformed, one-way ANOVA, Sidak’s correction], **c** carcass [Ln(y) transformed, Brown-Forsythe/Welch ANOVA, Dunnett’s correction]. However, viral titers do not change over time in any tissue. **d** Independent of time (including 11–18 dpi), AgDENV titers in the MAGs and carcass are significantly higher than those in the testes [Ln(y) transformed, Brown-Forsythe/Welch ANOVA, Dunnett’s correction]. NS indicates *P*-value > 0.05. Throughout, for testes and MAGs, each point represents a pool of 12 tissues; for carcasses, each point represents an individual carcass with the reproductive tract removed. Viral genomes are normalized to host genomes (*S7*). Abbreviations: AgDENV, *Anopheles gambiae* densovirus; ANOVA, analysis of variance; dpi, days post-injection; Ln(y), log transformed; MAGs, male accessory glands; PBS, phosphate-buffered saline; *S7*, host ribosomal protein gene; VP, viral protein




Fig. 2 Venereally transmitted AgDENV causes disseminated infection in mated females. **a, b** Females mated to AgDENV-injected males have **a** Detectable AgDENV in the spermatheca [Ln(y) transformed, unpaired Student’s t-test; each point = pool of 5–13 tissues], **b** disseminated AgDENV infection in the carcass [Ln(y) transformed, Welch’s t-test, each point = single carcass with spermatheca removed], when analyzed 8–14 days post-mating and relative to females mated to PBS-treated mates. Viral genomes are normalized to host genomes (S7)

that both components of the male reproductive tract are susceptible to infection, but that AgDENV likely replicates more efficiently in the MAGs than in the testes.

To assess whether AgDENV infection in male mosquitoes affects their ability to mate, we introduced uninfected, virgin females into cages with either AgDENV- or PBS-injected males. After 3 nights of mating, females were dissected and mating status was determined by the presence or absence of sperm in the spermatheca. We found no difference in the proportion of mated females between those exposed to AgDENV-injected males and those exposed to PBS-injected males (Table 1), indicating that AgDENV infection does not significantly hinder male mating ability. Additionally, we found no difference in the

Table 1 Mating rates between virgin females with either AgDENV- or PBS-injected male *Anopheles gambiae*

Mating rates ^a	PBS-injected males		AgDENV-injected male	
	Mated	Unmated	Mated	Unmated
P	29%	71%	36%	64%
N	46	111	63	114
Total number	157		177	

The mating rate between PBS-injected males and AgDENV-injected males did not differ (Fisher’s exact test, $P=0.2432$), as determined by the number of females that were inseminated after being left with males for 3 nights

AgDENV *Anopheles gambiae* densovirus, PBS phosphate-buffered saline

^a N = number of female mosquitoes, either with or without sperm present in the spermatheca (classified as mated or unmated, respectively). P = percentage of female mosquitoes that are mated or unmated

Table 2 Male *An. gambiae* survival rates

Survival rates ^a	PBS-injected males		AgDENV-injected males	
	Alive	Dead	Alive	Dead
P	71%	29%	75%	25%
N	371	151	428	141
Total number	522		569	

The survival of PBS-injected males and AgDENV-injected males (Fisher’s exact test, $P=0.1322$) did not differ when assessed at 18 dpays post-injection

^a N = number of male mosquitoes that are either alive or dead. P = percentage of male mosquitoes that are either alive or dead


survival of AgDENV-injected males compared to controls when assessed at 18 dpi (Table 2), supporting previous findings that AgDENV infection does not affect longevity of its *An. gambiae* host [8].

An. gambiae females develop disseminated infection after mating with AgDENV-infected males

Authors of previous studies reported that AgDENV is venereally transmitted from infected males to the spermatheca of mated females [6]. To test whether sexually-transmitted AgDENV can also lead to disseminated infection in mated females, we again allowed uninfected, virgin females to mate with either AgDENV- or PBS-injected males. After mating, the females were separated and maintained for 8–14 days before being analyzed. As expected, we detected AgDENV in the spermatheca of females mated to infected males (Fig. 2a). We also found significant levels of AgDENV in the female carcass (Fig. 2b). Although AgDENV titers were variable, an estimated 90% of mated females ($n=45$) developed disseminated infection after exposure to AgDENV-infected males (see Methods).

Females venereally infected with AgDENV fail to vertically transmit to F1 progeny

We next assessed whether females sexually infected with AgDENV can vertically transmit the virus to their offspring. Specifically, after females were mated to AgDENV- or PBS-injected males, they were provided a blood meal and an oviposition site. Oviposited eggs were reared to adulthood, and F1 male and female offspring were pooled and assessed for virus. Across all samples tested, AgDENV was not detectable in F1 progeny (whole bodies) from either a first or second blood-feeding (Fig. 3a, b). These findings reveal that even though venereally transmitted AgDENV leads to disseminated infection in mated females, they do not vertically transmit the infection to their offspring.



Discussion

Here we report that venereal transmission of AgDENV leads to disseminated infection in *An. gambiae* females. This finding reveals that infected male releases could be an effective way to introduce AgDENV-based paratransgenic tools into field populations of *An. gambiae* females. More specifically, released males harboring a modified AgDENV would mate with wild females and venereally transmit the virus to them. Then, as AgDENV titers build up in the female post-mating, a viral-encoded effector could interfere with the ability of the mosquito to transmit pathogens like malaria parasites. Male releases have already been shown to be a safe and effective way to deploy biocontrol agents into mosquito populations [14–16], largely because male mosquitoes do not blood feed and thus are not responsible for pathogen transmission. Treatment of larval breeding sites is another means by which ISV-based control tools could be deployed in the field, as was done for the AaeDENV-based larvicide [5, 12], but this approach poses significant challenges for *An. gambiae*. *Anopheles gambiae* females often lay their eggs in small temporary breeding sites produced by rain fall, making large-scale application of larvae-targeting tools generally impractical for this mosquito species in the field setting [18]. Therefore, infected male releases offer a more logistically feasible method to introduce AgDENV-based control tools into wild *An. gambiae* populations.

For infected male releases to be successful, AgDENV must not only be venereally transmitted, but AgDENV-infected males must also be fit and able to mate at rates competitive with wild males. In the present study, we

found that AgDENV infection in males does not affect mating rates, a good first indication of reproductive fitness. To fully assess mating competitiveness for field releases, mate-choice experiments with wild-type mosquitoes in a semi-field environment would still be needed. We also assessed the tissue tropism of AgDENV in male mosquitoes and found that both the testes and the MAGs become infected, indicating that venereally transmitted virus could be transferred via sperm (produced in the testes), the mating-plug (produced from the MAGs) or both. After mating, sperm is stored in the female spermatheca and the mating-plug is digested in the female atrium (uterus) [19]; since we observed AgDENV in both the spermatheca and the carcass of mated females, it is likely that both components contribute to sexual transmission of AgDENV. While we observed AgDENV in all spermatheca pools tested (Fig. 2a), due to the pooling of tissues, we are unable to determine the exact rate of venereal transmission. In the future, paired mating assays, such as mating captures, coupled with individual tissue analyses could provide this information as well as enable determination of how variation in male viral titers influences transmission and dissemination in mated females.

We also found no difference in AgDENV titers across time in any of the male tissues assessed, demonstrating that peak viral titers are achieved by 11 dpi, in agreement with the results of a previous study [6], which found that adult AgDENV titers increase up to ~12 dpi.

In this study, we used the microinjection method to infect male mosquitoes with AgDENV. This approach allowed for controlled and reproducible delivery of inoculating AgDENV doses for the characterization of infection and transmission dynamics. However, microinjections are laborious and would be an impractical strategy to infect male mosquitoes on a large scale, as would be needed for an AgDENV-based male release program. Therefore, although logistically challenging for field applications, treatment of larval breeding water with AgDENV could be an effective way of mass-producing infected *An. gambiae* in the laboratory. In fact, it was previously shown that adult mosquitoes can become infected with AgDENV following the exposure of larvae to AgDENV-treated water [1, 7]. This infection method would need to be further developed for operational purposes, but previous data suggest that when mosquitoes are infected via larval exposure, adult AgDENV titers are similar to those achieved by injection [6, 7] (Fig. 1c). Notably, viral titers may even reach their peak more quickly in adults that were infected as larvae [7] as compared to those infected by injection [6] (7 vs. 12 days post-emergence, respectively). This means that larval exposure to AgDENV could allow for the release of younger infected males, furthering the feasibility of this approach over microinjections

for operational purposes. Data from females suggest that AgDENV tissue tropism is consistent between infection routes [1, 13], yet male tissue tropism and venereal transmission dynamics following larval exposure still need to be tested. This information would help optimize the larval exposure method in the laboratory to maximize male infection and venereal transmission rates for any future AgDENV-based male release programs.

A previous study showed that AgDENV is transferred to the female spermatheca immediately post-mating [6], but the authors of that study did not detect significant dissemination in the female carcass. This discrepancy likely arises due to low statistical power in the previous study [6]: elevated DENV was detected in the carcass but it was not statistically significant. In addition, females in this study [6] were assessed immediately after mating, whereas in the present study we assayed for AgDENV in females 8–14 days post-mating, allowing time for the virus to replicate. Given that AgDENV titers are known to rise gradually in adult females [6, 7], it is not surprising that it took a few days for AgDENV titers to increase to significant levels in mated females. In fact, the gradual increase in AgDENV titers over time is one reason this ISV has been proposed as a candidate for a “late-life-acting” insecticide [7, 20]. In the present study, we did not assess any female fitness parameters following venereally acquired AgDENV infection because previous work [8] found that mosquitoes directly infected with high doses of AgDENV do not exhibit reduced fitness or even a marked transcriptional response. We do not expect venereally transmitted AgDENV to affect female fitness, but prior to implementation of AgDENV-based tools, this possibility would need to be tested.

Although venereally transmitted AgDENV produces disseminated infection in females, we found no evidence of vertical transmission from these females. Interestingly, previous studies demonstrated vertical transmission of AgDENV when both male and female mosquitoes were directly infected by exposure to virus during larval development [1]. There are several possible reasons why we did not observe vertical transmission in our experiments. First, although AgDENV was present in the female body, it may not have infected the ovaries/oocytes. Secondly, AgDENV present in the spermatheca may not be directly associated with sperm cells and therefore not likely to enter the egg during fertilization. Finally, it is possible that AgDENV titers simply were not high enough for vertical transmission to occur in these mosquitoes. Without vertical transmission, an AgDENV-based vector control tool would not spread trans-generationally following infected male releases. This means that a greater frequency of male releases would be required to have an impact on wild *An. gambiae* populations; on the other

hand, this feature may be advantageous from a public acceptance stand-point, because the virus and its encoded effectors would not be heritable across multiple generations. This feature would give more user control over the application of these paratransgenic tools.

Our studies here focused exclusively on the infection and transmission dynamics of wild-type AgDENV. This contrasts with several previous studies of AgDENV-based paratransgenesis that have largely focused on a two-virus helper/transducer system (consisting of one wild-type AgDENV and one defective, GFP-expressing AgDENV) [1, 6, 13]. While this helper/transducer approach offers a larger capacity for transgenic cargo, it is far more practical to utilize a single, self-replicating transgenic virus for application in field settings. Viral packaging efficiency limits the size of transgenes that can be encoded in a single AgDENV vector, but small anti-pathogenic effectors or short-hairpin RNAs targeting host processes could be expressed under an endogenous promoter to limit the mosquito’s ability to transmit pathogens. Further engineering and testing are still needed to identify and validate the best effectors for this system.

Finally, in addition to offering paratransgenic tools, ISVs can also have natural pathogen-blocking effects [21]. For example, AgDENV has an inhibitory effect on the emerging arbovirus Mayaro virus [22], which can be transmitted by *Anopheles* [23]. Wild-type AgDENV may also inhibit additional pathogens, such as the malaria parasite, although this remains to be explored. Together, our findings underscore the importance of basic research into the dynamics of wild-type ISVs such as AgDENV, which hold promise for both basic and applied purposes.

Conclusions

These findings reveal that if AgDENV is developed into a paratransgenic tool for *Anopheles gambiae* mosquitoes, it could be potentially deployed in the field via infected male releases. *Anopheles gambiae* spreads more fatal infections to humans than any other vector, yet to date we still lack genetic tools for targeting this species and AgDENV may fill this gap.

Abbreviations

AgDENV: *Anopheles gambiae* densovirus; dpi: Days post-injection; ISV: Insect-specific virus; MAGs: Male accessory glands; pAgDENV: Plasmid containing complete AgDENV genome; S7: Host ribosomal protein gene; vge: Viral genome equivalents; VP: Viral protein (gene).

Acknowledgements

The authors wish to thank Ms Amelia Romo and Ms Kaylee Montanari for their assistance rearing the *An. gambiae* colony mosquitoes. The graphical abstract was created using BioRender.com.

Author contributions

KLW and JLR conceived and designed the study and analyzed the data. KLW, JLR, and HCM wrote and edited the manuscript. KLW performed the

experiments. JLR provided materials and reagents. RMJ contributed protocols, reagents and experimental insights. K LW and HCM designed mosquito graphs. All authors read and approved the final manuscript.

Funding

This research was supported by NIH/NIAID grant R01AI128201 to JLR.

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Entomology, Pennsylvania State University, University Park, PA, USA. ²Center for Infectious Disease Dynamics, Pennsylvania State University, University Park, PA, USA. ³The Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA, USA. ⁴Present Address: Department of Environmental Sciences, The Connecticut Agricultural Experiment Station, New Haven, CT 06504, USA.

Received: 11 February 2022 Accepted: 20 May 2022

Published online: 20 June 2022

References

- Ren X, Hoiczyc E, Rasgon JL. Viral paratransgenesis in the malaria vector *Anopheles gambiae*. *PLoS Pathog*. 2008;4:e1000135.
- Suzuki Y, Barik TK, Johnson RM, Rasgon JL. In vitro and in vivo host range of *Anopheles gambiae* densovirus (AgDENV). *Sci Rep*. 2015;5:12701.
- Patterson EI, Villinger J, Muthoni JN, Dobel-Ober L, Hughes GL. Exploiting insect-specific viruses as a novel strategy to control vector-borne disease. *Curr Opin Insect Sci*. 2020;39:50–6.
- Bolling BG, Weaver SC, Tesh RB, Vasilakis N. Insect-specific virus discovery: significance for the arbovirus community. *Viruses*. 2015;7:4911–28.
- Johnson RM, Rasgon JL. Densovirus (densovirus) for mosquito and pathogen control. *Curr Opin Insect Sci*. 2018;28:90–7.
- Barik TK, Suzuki Y, Rasgon JL. Factors influencing infection and transmission of *Anopheles gambiae* densovirus (AgDENV) in mosquitoes. *PeerJ*. 2016;4:e2691.
- Ren X, Rasgon JL. Potential for the *Anopheles gambiae* densovirus to act as an “evolution-proof” biopesticide. *J Virol*. 2010;84:7726–9.
- Ren X, Hughes GL, Niu G, Suzuki Y, Rasgon JL. *Anopheles gambiae* densovirus (AgDENV) has negligible effects on adult survival and transcriptome of its mosquito host. *PeerJ*. 2014;2:e584.
- Hirunkanokpun S, Carlson JO, Kittayapong P. Evaluation of mosquito densovirus for controlling *Aedes aegypti* (Diptera: Culicidae): variation in efficiency due to virus strain and geographic origin of mosquitoes. *Am J Trop Med Hyg*. 2008;78:784–90.
- Ledermann JP, Suchman EL, Black WC, Carlson JO. Infection and pathogenicity of the mosquito densovirus AeDENV, HeDENV, and APeDENV in *Aedes aegypti* mosquitoes (Diptera: Culicidae). *J Econ Entomol*. 2004;97:1828–35.
- Kittayapong P, Baisley KJ, O'Neill SL. A mosquito densovirus infecting *Aedes aegypti* and *Aedes albopictus* from Thailand. *Am J Trop Med Hyg*. 1999;61:612–7.
- Buchatskiĭ LP, Kuznetsova MA, Lebedinets NN, Kononko AG. Development and basic properties of the viral preparation viroden. *Vopr Virusol*. 1987;32:729–33.
- Suzuki Y, Niu G, Hughes GL, Rasgon JL. A viral over-expression system for the major malaria mosquito *Anopheles gambiae*. *Sci Rep*. 2014;4:5127.
- Crawford JE, Clarke DW, Criswell V, Desnoyer M, Cornel D, Deegan B, et al. Efficient production of male Wolbachia-infected *Aedes aegypti* mosquitoes enables large-scale suppression of wild populations. *Nat Biotechnol*. 2020;38:482–92.
- Harris AF, McKemey AR, Nimmo D, Curtis Z, Black I, Morgan SA, et al. Successful suppression of a field mosquito population by sustained release of engineered male mosquitoes. *Nat Biotechnol*. 2012;30:828–30.
- Carvalho DO, McKemey AR, Garziera L, Lacroix R, Donnelly CA, Alphey L, et al. Suppression of a field population of *Aedes aegypti* in Brazil by sustained release of transgenic male mosquitoes. *PLoS Negl Trop Dis*. 2015;9:e0003864.
- Rogers DW, Baldini F, Battaglia F, Panico M, Dell A, Morris HR, et al. Transglutaminase-mediated semen coagulation controls sperm storage in the malaria mosquito. *PLoS Biol*. 2009;7:e1000272.
- Centers for Disease Control and Prevention GH, Division of Parasitic Diseases and Malaria. Larval control and other vector control interventions 2020. https://www.cdc.gov/malaria/malaria_worldwide/reduction/vector_control.html. Accessed 10 Feb 2022.
- Rogers DW, Whitten MM, Thailayil J, Soichot J, Levashina EA, Catteruccia F. Molecular and cellular components of the mating machinery in *Anopheles gambiae* females. *Proc Natl Acad Sci USA*. 2008;105:19390–5.
- Koella JC, Lynch PA, Thomas MB, Read AF. Towards evolution-proof malaria control with insecticides. *Evol Appl*. 2009;2:469–80.
- Carvalho VL, Long MT. Insect-specific viruses: an overview and their relationship to arboviruses of concern to humans and animals. *Virology*. 2021;557:34–43.
- Urakova N, Brustolin M, Joseph RE, Johnson RM, Pujhari S, Rasgon JL. *Anopheles gambiae* densovirus (AgDENV) negatively affects Mayaro virus infection in *Anopheles gambiae* cells and mosquitoes. *Parasit Vectors*. 2020;13:210.
- Brustolin M, Pujhari S, Henderson CA, Rasgon JL. *Anopheles* mosquitoes may drive invasion and transmission of Mayaro virus across geographically diverse regions. *PLoS Negl Trop Dis*. 2018;12:e0006895.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

