



ICTV Virus Taxonomy Profile: *Peribunyaviridae*

Holly R. Hughes¹, Scott Adkins², Sergey Alkhovskiy³, Martin Beer⁴, Carol Blair⁵, Charles H. Calisher⁵, Mike Drebot⁶, Amy J. Lambert^{1,*}, William Marciel de Souza⁷, Marco Marklewitz⁸, Márcio R. T. Nunes⁹, Xiāohóng Shí (石晓宏)¹⁰ and ICTV Report Consortium

Abstract

Peribunyaviruses are enveloped and possess three distinct, single-stranded, negative-sense RNA segments comprising 11.2–12.5 kb in total. The family includes globally distributed viruses in the genera *Orthobunyavirus*, *Herbevirus*, *Pacuvirus* and *Shangavirus*. Most viruses are maintained in geographically-restricted vertebrate–arthropod transmission cycles that can include transovarial transmission from arthropod dam to offspring. Others are arthropod-specific. Arthropods can be persistently infected. Human infection occurs through blood feeding by an infected vector arthropod. Infections can result in a diversity of human and veterinary clinical outcomes in a strain-specific manner. Segment reassortment is evident between some peribunyaviruses. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the taxonomy of the family *Peribunyaviridae*, which is available at ictv.global/report/peribunyaviridae.

Table 1. Characteristics of members of the family *Peribunyaviridae*

Typical member	<i>Bunyamwera virus</i> (S, D00353; M, M11852; L, X14383), species <i>Bunyamwera orthobunyavirus</i> , genus <i>Orthobunyavirus</i>
Virion	Enveloped, spherical or pleomorphic virions, 80–120 nm in diameter
Genome	Three single-stranded, negative-sense RNA molecules, S, M and L, each of about 1, 4 and 6.8 kb
Replication	Cytoplasmic; primary transcription is primed by ‘cap snatching’ of host RNAs
Translation	On ER-bound ribosomes for Gn and Gc and on free ribosomes in the cytoplasm for N and L
Host range	Vertebrates and invertebrates (including mammals, birds, mosquitoes, culicoids and psychodid sandflies)
Taxonomy	Phylum <i>Negarnaviricota</i> , subphylum <i>Polyploviricotina</i> , class <i>Ellioviricetes</i> , order <i>Bunyavirales</i> , several genera and >90 species

VIRION

Peribunyavirus virions are spherical or pleomorphic, 80–120 nm in diameter [1] with glycoprotein surface projections (5–18 nm) embedded in a lipid bilayer envelope (about 5 nm) (Table 1, Fig. 1). The genome comprises three single-stranded negative-sense RNAs (designated S, small, M, medium and L, large) (Fig. 2), each with complementary terminal nucleotide sequences that base-pair to form non-covalently closed, circular RNAs [2] that are individually encapsidated.

GENOME

The S segment encodes the nucleocapsid protein (N), which is abundant in infected cells; in some viruses an overlapping reading frame encodes the non-structural protein NSs [3–5]. The M segment encodes two structural glycoproteins (Gn and Gc). Some members also encode a non-structural protein (NSm) between the Gn and Gc coding regions [4]. The L segment encodes the L protein, which has RNA-directed RNA polymerase and endonuclease functions.

Received 18 November 2019; Accepted 18 November 2019; Published 17 December 2019

Author affiliations: ¹Centers for Disease Control and Prevention, Fort Collins, CO, USA; ²United States Department of Agriculture, Agricultural Research Service, Fort Pierce, FL, USA; ³D. I. Ivanovsky Institute of Virology, Moscow, Russia; ⁴Friedrich-Loeffler-Institut, Greifswald-Insel Riems, Germany; ⁵Colorado State University, Fort Collins, CO, USA; ⁶Public Health Agency of Canada, Winnipeg, MB, Canada; ⁷University of Campinas (UNICAMP), Campinas, São Paulo, Brazil; ⁸Charité-Universitätsmedizin Berlin, Humboldt-University Berlin, and Berlin Institute of Health, Berlin, Germany; ⁹Evandro Chagas Institute, Ministry of Health, Ananindeua, Pará, Brazil; ¹⁰MRC-University of Glasgow Centre for Virus Research, Glasgow, UK.

*Correspondence: Amy J. Lambert, ahk7@cdc.gov

Keywords: *Peribunyaviridae*; bunyavirus; ICTV Report; *Orthobunyavirus*; *Herbevirus*; *Pacuvirus*; *Shangavirus*; taxonomy.

Abbreviations: G, Glycoprotein; L, N and NS, large, nucleocapsid and non-structural proteins; S, M and L, Small, medium and large vRNA segments. 001365






Fig. 1. Peribunyavirus virion structure. (a) representation of a virion in cross-section. The surface spikes comprise the Gn and Gc glycoproteins. The helical nucleocapsids are circular and comprise one each of the unique ssRNA segments (L, large; M, medium; S, small) encapsidated by N protein and associated with the L protein. (b) negative-stained transmission electron microscopy photograph of California encephalitis virus virions (image: CDC/Drs Frederick Murphy and Erskine Palmer).

Bunyamwera virus




Fig. 2. Coding strategy of Bunyamwera virus. Translation of NSs protein is initiated at an alternative start codon. The Gn, NSm and Gc proteins are generated by co-translational cleavage of M polyprotein.

REPLICATION

Virions attach via surface glycoproteins, entering the cell through clathrin-mediated endocytosis. Fusion of the viral Gc protein fusion peptide with endosomal membranes facilitates the release of ribonucleocapsids into the cytoplasm. The complementary 5'- and 3'-terminal ends serve as promoters for both mRNA and antigenome synthesis. Viral mRNAs are not polyadenylated and are truncated relative to the vRNA; a 5'-methylated cap is derived from host mRNA via 'cap snatching' mediated by the endonuclease function of the L protein. Proteins are translated on free ribosomes (S and L segment mRNAs) or membrane-bound ribosomes (M segment mRNA). The Gn and Gc proteins are generated by co-translational cleavage and targeted to and retained in the Golgi complex. Ribonucleoproteins are targeted

near the Golgi complex. Genomes are packaged by signals from non-conserved sequences in the terminal untranslated regions. Virions bud into Golgi cisternae and are transported to the cell surface by the secretory pathway [6].

TAXONOMY

Genera are monophyletic based on analysis of the virus L protein; members of a genus have similar genomic organizations and transmission cycles. Peribunyaviruses form a group in the phylum *Negarnaviricota*, subphylum *Polyploviricotina*, class *Ellioviricetes*, order *Bunyavirales*, being most closely related to viruses of the families *Fimoviridae* and *Tospoviridae*. Peribunyaviruses share some of the following characteristics: (i) enveloped spherical or pleomorphic virions; (ii) three segments of single-stranded, negative-sense RNA, with all proteins encoded in the same sense; (iii) capped but not polyadenylated viral mRNA; (iv) establish a persistent infection in an arthropod vector.

RESOURCES

Current ICTV Report on the family *Peribunyaviridae*: ictv.global/report/peribunyaviridae.

Funding information

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA). W. M. S. is supported by Fundação de Amparo à Pesquisa do Estado de São Paulo, Brazil (17/13981-0).

Acknowledgements

Members of the ICTV Report Consortium are Stuart G. Siddell, Andrew J. Davison, Elliot J. Lefkowitz, Peter Simmonds, Sead Sabanadzovic, Donald B. Smith, Richard J. Orton and Jens H. Kuhn. The findings and conclusions in this manuscript are those of the authors and do not necessarily represent the views of the CDC or other organizations.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

- Martin ML, Lindsey-Regnery H, Sasso DR, McCormick JB, Palmer E. Distinction between *Bunyaviridae* genera by surface structure and comparison with Hantaan virus using negative stain electron microscopy. *Arch Virol* 1985;86:17–28.
- Raju R, Kolakofsky D. The ends of La Crosse virus genome and antigenome RNAs within nucleocapsids are base paired. *J Virol* 1989;63:122–128.
- de Melo AB, de Souza WM, Acrani GO, Carvalho VL, Romeiro MF et al. Genomic characterization and evolution of Tacaiuma orthobunyavirus (*Peribunyaviridae* family) isolated in Brazil. *Infect Genet Evol* 2018;60:71–76.
- Marklewitz M, Zirkel F, Rwegu IB, Heidemann H, Trippner P et al. Discovery of a unique novel clade of mosquito-associated bunyaviruses. *J Virol* 2013;87:12850–12865.
- Rodrigues DS, Medeiros DB, Rodrigues SG, Martins LC, de Lima CP et al. Pacui virus, Rio Preto da Eva virus, and Tapirape virus, three distinct viruses within the family *Bunyaviridae*. *Genome Announc* 2014;2:e00923–14.
- Ferron F, Weber F, de la Torre JC, Reguera J. Transcription and replication mechanisms of *Bunyaviridae* and *Arenaviridae* L proteins. *Virus Res* 2017;234:118–134.