

Circadian Tick-Talking Across the Neuroendocrine System and Suprachiasmatic Nuclei Circuits: The Enigmatic Communication Between the Molecular and Electrical Membrane Clocks

M. D. C. Belle

Faculty of Life Sciences, University of Manchester, Manchester, UK.

Journal of Neuroendocrinology

As with many processes in nature, appropriate timing in biological systems is of paramount importance. In the neuroendocrine system, the efficacy of hormonal influence on major bodily functions, such as reproduction, metabolism and growth, relies on timely communication within and across many of the brain's homeostatic systems. The activity of these circuits is tightly orchestrated with the animal's internal physiological demands and external solar cycle by a master circadian clock. In mammals, this master clock is located in the hypothalamic suprachiasmatic nucleus (SCN), where the ensemble activity of thousands of clock neurones generates and communicates circadian time cues to the rest of the brain and body. Many regions of the brain, including areas with neuroendocrine function, also contain local daily clocks that can provide feedback signals to the SCN. Although much is known about the molecular processes underpinning endogenous circadian rhythm generation in SCN neurones and, to a lesser extent, extra-SCN cells, the electrical membrane clock that acts in partnership with the molecular clockwork to communicate circadian timing across the brain is poorly understood. The present review focuses on some circadian aspects of reproductive neuroendocrinology and processes involved in circadian rhythm communication in the SCN, aiming to identify key gaps in our knowledge of cross-talk between our daily master clock and neuroendocrine function. The intention is to highlight our surprisingly limited understanding of their interaction in the hope that this will stimulate future work in these areas.

Correspondence to:

M. D. C. Belle, Faculty of Life Sciences, University of Manchester, AV Hill Building, Manchester M13 9PT, UK (e-mail: mino.belle@manchester.ac.uk).

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Key words: suprachiasmatic nuclei, circadian rhythm, clock genes, neuroendocrine system, reproduction, electrical activity, ion channels

doi: 10.1111/jne.12279

The developmental onset of some fundamental neuroendocrine processes of the body, such as those involved in reproductive maturity, ranges from months to years depending on the species in question, and relies on constellations of tightly regulated hormonal signalling processes within the brain and body. Once established, this timely coordinated balance in hormonal activity and reproduction is temporally regulated by our master circadian clock in the suprachiasmatic nuclei (SCN). Indeed, work spanning over 50 years is unravelling the important and intimate link between SCN activity and appropriate timing in neuroendocrine function.

Circadian rhythms are daily timing in behaviour and physiology that persist (free run) in the absence of external time cues, such as light. In mammals, the master circadian clock resides in the SCN, where the activity of thousands of cell-autonomous clock neurones is synchronised each day to output ensemble circadian time cues to

the rest of the brain and body. At its most basic level, the circadian clockwork is conceptualised as an intrinsic intracellular process involving a dynamic interplay between the protein products of core clock genes such as *Period1/2* (*Per1/2*), *Cryptochrome 1/2* (*Cry 1/2*), *Clock* and *Bmal1*. In the past three decades, significant progress has been made to dissect and understand the intricate inner workings of this molecular timing machine that operates as a near 24-h transcription-translation molecular feedback loop (TTFL) (1,2).

Our current understanding of circadian rhythm generation in the SCN is that the TTFL activity within single SCN neurones drives daily changes in membrane excitability (3,4) (although see also the later section on the membrane clock). Indeed, neurones of the central nervous system can alter their intrinsic excitability by modulating the activity of their ion channels that provide excitatory or inhibitory drive to the membrane (5). How the TTFL achieves this,

however, is largely unknown. Broadly speaking, activation of sodium and some calcium channels and/or reduction in potassium channel activity depolarise the resting membrane potential (RMP) of the neurones, causing action potential (AP) discharge. By contrast, activation of potassium and/or chloride channels provides hyperpolarising forces that suppress electrical activity. In SCN neurones, both the conductivity and transcript activity of several ion channels are under circadian control (3,4). This allows SCN neurones to generate daily rhythms in electrical outputs, a hallmark feature of SCN cells that is paramount to the functioning of the circadian timing system. Measurements of AP firing rate in unidentified SCN neurones both *in vivo* and *in vitro* SCN brain slice preparations show that, at the population level, SCN neurones are significantly more active during the day (an up state: discharging APs at 4–6 Hz) than at night (a down state: generating APs at 0.1–2 Hz) (Fig. 1). Even when isolated from the SCN tissue, single SCN neurones retain the intrinsic ability to generate daily rhythm in AP activity (6,7) that can be maintained for days. It is considered that this day–night alteration in electrical states synchronises the activity of SCN neurones and communicates circadian information to the brain and body (3). Evidence from imaging and electrophysiological studies, however, suggests that the manner through which SCN neurones encode and communicate circadian information is more complex

than first assumed and may not solely rely on day–night variation in AP firing rate.

Under natural conditions, SCN clock neurones are entrained/synchronised to the light and dark cycle by responding to the daily glutamatergic activity of the retino–hypothalamic tract (Fig. 2). This excitatory drive is conveyed to the SCN by specialised melanopsin-containing retinal ganglion cells that communicate ambient light levels directly to SCN neurones (8), causing *Per1* gene activation in these cells (9). This allows environmental light cues to reset SCN clock neurones and coordinate their activity to generate a stable internal representation of solar time.

Several internal physiological signals also feedback and fine tune timing precision in the SCN. Such intrinsic feedback cues [referred to as Zeitnehmer ('time-taker'), a term applied to input pathways that are rhythmically regulated by feedback from an oscillator] (10,11) emerge mainly from the body's homeostatic systems and are communicated to the SCN using a variety of signalling neuropeptides to mainly suppress *Per1/2* gene expression and electrical activity. These feedback signals include neuropeptide Y (NPY) neurones of the thalamic intergeniculate leaflet, which send monosynaptic projections to the SCN via the geniculo–hypothalamic tract (12), as well as arousal-promoting orexinergic neurones of the lateral hypothalamus (13), which also send axonal contacts to SCN




Fig. 1. A schematic view of the daily electrical profiles of suprachiasmatic nucleus (SCN) neurones. (A) Over the day–night cycle, SCN neurones show overt changes in their resting membrane potential (RMP), and traverse through several points of neutral rest state (indicated by where the solid line crosses the dashed line). During the day, the RMP of SCN neurones is depolarised. In some cells, reduced activity of L-type calcium and calcium activated potassium (K_{Ca}) channels partly underpins this up state (128). At night, increased conductivity of multiple potassium channels hyperpolarises the neurones, placing them into a down state (3,4). (B) In some SCN neurones, this daytime up state causes action potential (AP) discharge (b1). In others, however, the RMP becomes too positive (~ -30 mV) to sustain AP production (b2). Instead, these neurones display depolarised low-amplitude membrane oscillation (b2). At night, during the RMP down state, SCN neurones generate action potentials at a significantly reduced rate (b3). This shows the complexity, richness and diversity of electrical communication in SCN neurones. The light area indicates the day, whereas the shaded region shows the night. The increase or decrease underlying ion channel activity is indicated by upward- and downward-pointing blue arrows, respectively.




Fig. 2. Cross-talk between the suprachiasmatic nucleus (SCN) and neuroendocrine cells. (A) Cross-section of the mouse brain at the anatomical level of the SCN (in red). (B) Bidirectional communication between the molecular clock (transcription–translation molecular feedback loop; TTFL) and electrical/membrane clock. In this model, the TTFL clock drives day–night rhythms in the electrical activity of SCN neurones, and electrical activity feedback onto the TTFL clock through unknown mechanisms (light blue arrows). This may underlie how the TTFL outputs signals to neuroendocrine cells (C) and how neuroendocrine processes feedback to adjust circadian timing in the SCN (grey arrows). This drawing is superimposed on top of a modified image taken at the SCN mid-coronal section showing *Per1*-EGFP neurones (red arrows). Darker blue arrows represent inputs to the SCN, with their sizes denoting feedback magnitude. Solid arrows indicate an established link, whereas broken arrows show tentative interactions. (b1) Stylised waveform showing daily variation in SCN *Per1* and electrical activity. White and black bars underneath represent day and night, respectively. RHT, retino-hypothalamic tract; LHI, lateral hypothalamic input; GHT, geniculohypothalamic tract; OX, optic chiasm; 3V, third ventricle; NPY, neuropeptide Y; VIP, vasoactive intestinal polypeptide; AVP, arginine vasopressin.

neurones (14). In the SCN, these neuropeptidergic signals can converge onto single clock neurones to modulate electrical activity and circadian timing in this brain structure (14,15). As such, SCN neurones integrate both intrinsic and extrinsic signals and send collective circadian time cues by neural and paracrine pathways to the rest of the brain, including many neuroendocrine hypothalamic structures (Fig. 2). Thus, the physiology and behaviour of the organism are tuned to anticipate and adapt to the solar day–night cycle, a key physiological prerequisite for survival and reproductive success. The SCN therefore represents a good example of localised autonomous function in the brain, and provides a unique opportunity in neuroscience to link highly organised behaviours (e.g. sleep–wake cycle, daily feeding drive and neuroendocrine functions) to the activities of a known population of neurones in the brain.

Circadian regulation of neuroendocrine functions

The realisation of the importance of circadian timing in hormone function started with the work of Everett and Sawyer in the 1950s. In a pioneering study, it was discovered that, in nocturnal rodents, a stimulatory drive occurring at a narrow temporal window in the pro-oestrus afternoon is necessary for the induction of ovulation later at night (16). Subsequently, a number of studies both in primates (including humans) and rodents have reported circadian rhythms in gonadotrophin-releasing hormone (GnRH) gene expres-

sion (17) and in the levels of many of the body's endocrine hormones, such as luteinising hormone (LH), testosterone, prolactin (PRL), cortisol and gonadotrophins after the onset of puberty (18–22). A more in-depth description of hormonal secretion profiles is provided elsewhere (23).

Indeed, when placed in constant conditions, the oestrus cycle in rodents free-runs (24,25); a description of circadian terms is provided in Liu *et al.* (26). This supports its reliance on a master timing process governing daily cycles in mammalian endocrine physiology. Furthermore, ablation of the SCN or its output pathways abolishes the night-time LH surge and circadian rhythms in the release of a number of endocrine hormones, such as PRL and gonadotrophin (27–29).

Perhaps the most direct evidence of an SCN-dependent driven rhythm in hormonal secretion comes from elegant studies in female hamsters. Under normal conditions, both halves of the bilateral SCN in hamsters operate in synchrony. When these animals are housed in constant light, however, the SCN activity and behavioural rhythms 'split' in some hamsters, giving rise to two activity bouts in a 24-h period (30). Remarkably, these 'split' females display two daily LH surges, each in anti-phase and of half the concentration seen in 'nonsplit' controls (31). More recently, work in rodents has definitively linked circadian clock gene activity in the SCN with circadian timing in hormone synthesis and secretion (32).

Taken together, these studies demonstrate that important aspects of neuroendocrine physiology exhibit circadian variation, and that

the master circadian clock is central to the timing of endocrine processes. This provides an excellent model system in which to investigate circadian and endocrine interactions.

Circadian signalling to neuroendocrine targets

Neurons of the SCN are neurochemically and functionally heterogeneous and form distinct anatomical clusters within this brain structure. Ventral SCN neurones synthesise vasoactive intestinal polypeptide (VIP), whereas the dorsal neurones contain arginine vasopressin (AVP) (33), two neuropeptides that are rhythmically produced in the SCN and are critically involved in appropriate circadian function (6,34–36). Although the VPAC₂ receptor, the preferred receptor for VIP in the SCN, is expressed throughout this nucleus (37), VIP neurones generally project dorsally to the vicinity of AVP cells (33). Here, they form an ensemble bundle with AVP-axons that project away from the SCN.

In addition to forming conventional cell-to-cell synaptic contacts, SCN neurones can also signal circadian timing to the body in a paracrine fashion. Indeed, the timing of behavioural rhythms in rodents, such as locomotor activity, drinking and gnawing, are under the control of the SCN paracrine tone (38–41). An ingenious *in vitro* study has also recently shown that some intrinsic SCN clock function relies on paracrine communication amongst its neurones (mainly through diffusible neuropeptides such as VIP) (42), although further *in vivo* studies are needed to establish the role of paracrine signalling in SCN function. When sending circadian signals to the neuroendocrine system, however, the SCN does so via direct synaptic contacts onto endocrine cells or through populations of neurones that contact endocrine cells. This suggests that, unlike daily timing in behaviour (neuromodulation) (43), SCN communication to neuroendocrine cells requires the precision and speed of discrete synaptic transmission.

In the hypothalamic preoptic area (POA) of many species, VIP terminals originating from the SCN form direct contacts onto GnRH neurones, and indirectly connect to oestrodial-concentrating interneurons that communicate to GnRH cells (30,44–47). The VPAC₂ receptor is expressed by a subset of GnRH neurones (48,49) and VIP can directly modulate the activity of GnRH cells in a time-of-day related manner, therefore controlling the timing of LH release (50–53). Alteration of VIP signalling both *in vivo* and *in vitro* causes marked changes in GnRH, LH and gonadotrophin release (54–56), as well as in the magnitude and timing of the LH surge (55,57). Evidence also suggests that SCN VIP signalling to neuroendocrine dopaminergic neurones is involved in modulating pituitary PRL release (58). Indeed, paraventricular nucleus (PVN) dopaminergic neurones express the VPAC₂ receptor, hypothalamic VIP expression is in phase with PRL release, and work in rodents indicates that the activity of PVN dopaminergic cells suppresses pituitary PRL secretion (59–61).

There is also strong evidence in rodents suggesting that AVP signalling in the POA modulates the activity of GnRH neurones and is involved in PRL and LH secretion. AVP fibres contact GnRH neurones, and AVP secretion occurs in phase with GnRH release (45,62). Furthermore, work in female rats shows that mRNA for the AVP

receptor (*V1aR*) is expressed by a small population of POA GnRH neurones (63) and blockade of AVP receptors attenuates LH and PRL release (64). In SCN-lesioned animals, time-of-day-dependent AVP administration in the medial preoptic area rescues the LH surge (65–67). In the CLOCK- Δ 19 mutant mouse, where the activity of the circadian clock is severely disrupted, and the LH surge is dampened, the expression of AVP and *V1aR* in the SCN is also significantly reduced (68). Administration of AVP into the POA of these animals rescues the LH surge (68).

In the past few decades, it has also become increasingly clear that electrical and TFL activity of SCN neurones contributes to the regulation of seasonal rhythms (69). Day-length encoding by the SCN relies strongly on the plasticity of its neuronal network (70), which allows this brain structure to shorten and lengthen the phase relationship among its neurones, coding for the short winter and long summer days, respectively. This regulates the activity of key regions of the body that are important for seasonal changes in reproductive function, such as the seasonal release profile of the pineal gland neurohormone melatonin (69).

Taken together, these observations link the activity of the SCN, the plasticity of its neural circuits and the rhythmic release of some of its neuropeptides with the fundamental control of neurohormone secretion.

Extra-SCN endocrine circadian clocks

The discovery and understanding of the inner workings of the TFL come with the realisation that, although the SCN harbours the principal daily clock, many areas of the brain and body also have local circadian clockworks (71,72). These act in concert with the SCN to give rise to the extended circadian system. Indeed, core circadian clock genes are active in several neuroendocrine regions of the brain, pituitary and pineal glands (73–76), as well as in a number of peripheral endocrine tissues, such as the adrenal gland, testis and ovary (77,78). At the single cell level, clock gene expression is detected in GnRH and neuroendocrine dopaminergic neurones (79,80).

Furthermore, the daily expression profile of key clock genes, such as *Per1*, reveals rhythmic activity in several neuroendocrine brain regions and individual neurones, such as in dopaminergic cells that control PRL secretion (79,81–83). Studies using pituitary explants from transgenic animals bearing bioluminescent reporters of circadian clock activity have conclusively shown that pituitary cells can sustain intrinsic circadian rhythms in clock gene expression persisting over several days (84,85). Rhythmic expression of core clock genes is also reported in the pineal gland (86), in hypothalamic GnRH neurones (87–89) and in immortalised GnRH cell lines (80,87), where they serve to regulate GnRH cell activity and gate the sensitivity of these cells to daily hormonal signals (90,91). Disruption of clock gene activity in GnRH cell lines interrupts daily GnRH release in these cells (91) and, in preoptic-explants from *Bmal1* knockout mice, the stimulating drive in GnRH release is significantly compromised (88). Clock gene mutant rodent models are now confirming that disruption of appropriate circadian timing signals impairs hypothalamic and peripheral hormone secretion and interferes with reproductive success (92,93).

Taken together, these observations demonstrate that the functioning of local circadian clocks in neuroendocrine tissues and at the level of the SCN is important for normal neuroendocrine function. These neuroendocrine clock circuits may not only serve as gating mechanisms that determine whether and how neuroendocrine cells respond to periodic SCN time cues, but also provide tissue-specific local circadian time signals. This may influence output feedback signals from these tissues to the SCN (94). As such, the extended circadian system interacts to maintain optimal daily temporal alignment in physiology across the many tissues and organ systems of the body, although how SCN output molecules affect circadian gene expression (e.g. *Per1*) in neuroendocrine cells remains unknown.

Neuroendocrine feedback to the SCN: neuroendocrine neurones talking back

Although there is limited evidence of direct neuroendocrine cell projections to the SCN (95–98), several rhythmic hormonal output signals originating from neuroendocrine cell populations are known to act in the SCN. For example, in all animal species studied thus far, including humans, high-affinity receptors for melatonin, oestrogen, androgen and progesterone are present in the SCN (99–104). These hormones can act to modulate the electrical activity of SCN neurones and adjust the phase of the SCN clockworks (105–109). Furthermore, enzymatic conversion of these hormones, such as the aromatisation of testosterone to oestrogen, provides additional complexity to hormonal signals conveyed to the SCN (110). However, unlike the well-studied actions of feedback signals (e.g. NPY, serotonin and melatonin) on SCN clock functions, it remains unknown whether the phase of the molecular clockwork and the resulting electrical state of SCN neurones determines whether and how sex hormones act in the SCN. Moreover, how neuroendocrine feedback signals affect SCN circadian gene expression is also not known. Thus, more work is needed before the effects of these hormones on the SCN clockwork can be fully determined.

Although significant recent progress in circadian neuroendocrine studies now places the activity of the newly-identified kisspeptin and RFamide-related peptide at the core of the communication conduits interfacing SCN signals with neuroendocrine activity (111,112), many key questions still remain. For example, how does circadian information encoded by the hands of the clock (TTFL) translate into meaningful outputs that can be processed by the rest of the brain, including neuroendocrine cells, and how does the brain talk back to sculpt circadian timing in SCN clock neurones? Furthermore, retinal melanopsin neurones that project to the SCN via the RHT also form functional connectivity with a number of neuroendocrine hypothalamic regions (113,114). Therefore, can environmental light directly modulate the activity of hypothalamic neuroendocrine clock neurones and influence circadian and seasonal timing in these cells? To achieve such communication, signals travelling to and from the TTFL both in the SCN and neuroendocrine clock neurones must flow through the cell membrane (Fig. 3). However, the mechanistic nature of these signals that interweave the molecular clockwork and membrane activity to produce physiological circadian rhythms is largely unknown. This is a dilemma that




Fig. 3. A schematic view of communication between the transcription-translation molecular feedback loop (TTFL) in the suprachiasmatic nucleus (SCN) and neuroendocrine cells through the membrane clock. In this model, signals travelling to and from the TTFL both in the SCN and neuroendocrine clock neurones must flow through the cell membrane (bidirectional arrows) by unknown mechanisms. This process may underlie how the SCN TTFL outputs circadian signals to neuroendocrine cells (blue solid axon and arrow) and how neuroendocrine TTFL feedback to adjust circadian timing in SCN neurones (orange dotted axon and arrow). Through similar unknown processes, environmental and/or homeostatic cues can sculpt the activity of the TTFL in SCN and endocrine neurones (grey solid and dotted axon terminals, and arrows). Solid and broken axons/arrows show known and tentative links, respectively.

is now hampering major progress in circadian biology research, although some progress has been made in the SCN. Below, reference is made to the membrane clock as a model for addressing the mechanisms of information flow between the membrane and TTFLs: the all-important electrical-genetic interplay.

The membrane clock

Over the past 10 years, pioneering studies have revealed that the membrane properties, activity and excitability of SCN neurones are not only the proximal target of the TTFL, but also act to transmit information from the external world and body to the circadian molecular clockwork (Figs 2 and 3). Indeed, work on mammalian and *Drosophila* clock neurones strongly supports the concept that the electrical activity of clock neurones is integral to the functioning of the intracellular molecular clockwork (115–119). Indeed, the electrical state of clock neurones can impose time-of-day stamps onto its transcriptional programmes, thereby acting as an intrinsic zeitgeber (time-giver) (120). Studies performed in cultured neonatal SCN slices have also established a tight link between the TTFL activity, as measured by bioluminescence imaging, and membrane excitability in clock neurones. Here, abolishing AP discharge by tetrodotoxin (TTX) blockade of sodium channel activity, which also stops AP-dependent neurone-to-neurone communication, desynchronised the SCN clock and dampened TTFL rhythm amplitude within individual SCN neurones (121,122). Interestingly, however,

similar analysis using SCN brain slices from young adults showed that TTX administration had much less dramatic effects ((123); A. T. Hughes and H. D. Piggins, unpublished observations). In support, *in vivo* chronic TTX infusion in adult animals also failed to interfere with SCN function (124). This highlights the complex relationship between the TTFL and membrane clock and/or suggests developmental issues within the SCN neuronal network that require careful consideration (125). Therefore, from the paracrine influence to the electrical-genetic interplay in single SCN neurones, the SCN neural network plays a critical role in adult SCN function (70).

Nevertheless, growing evidence supports the idea that information flow between the membrane and TTFL may well underlie the processes by which neurotransmitters that are released from retinal cells and the body's homeostatic systems sculpt the timing precision in the SCN. TTFL-membrane communication might also underpin how the SCN network organises excitability of its neurones to accommodate extrinsic cues to communicate circadian and seasonal signals to extra-SCN TTFLs in the body (Fig. 3). Our limited mechanistic understanding of how this is achieved, however, may be inherently linked with the way we measure and report excitability in clock neurones, primarily in the SCN.

Further considerations: measuring electrical excitability in SCN neurones

Although the rate at which SCN neurones discharge APs is routinely used to report clock phase and functionality, measurement of AP firing frequency alone is not providing us with the complete picture. This is because SCN neurones also show overt day-night differences in their RMP and input resistance (Fig. 1) (126), which are other important modes of excitability in SCN neurones. Relying on AP firing rate measurements alone means that excitability in SCN neurones cannot be assessed when they are not producing APs; even in the up state during the day, single SCN neurones generate APs for 4–6 h only and not for the duration of the light phase. Further complexity arises when considering that not all SCN neurones contain a functional molecular clock (70).

Indeed, targeted recording of fluorescence tagged SCN neurones [from mice in which enhanced green fluorescent protein (EGFP) indicates *Per1* promoter activity (*Per1*-EGFP+ve cells)] (127) demonstrates that the electrical activity of SCN neurones is more complex and richer than was first assumed (128). It was revealed that the well-described day-night difference in AP generation in the SCN mostly comprises the activity of neurones in which the EGFP construct cannot be detected (EGFP-ve cells: presumed nonclock neurones). *Per1*-EGFP+ve SCN neurones stop discharging APs in the middle of the day as their RMP becomes too positive (~ -30 mV) to sustain AP generation. Instead, through reducing the activity of their calcium-activated potassium channels, these neurones enter a state of intrinsic depolarisation silencing (depolarisation blockade), producing L-type calcium channel dependent depolarised low-amplitude membrane oscillations (DLAMOs) instead of APs (128) (Fig. 1b). These modes of excitability may be important for the normal functioning of the circadian clock, and may be necessary to modulate intracellular calcium levels in SCN neurones, a key signal-

ling molecule in SCN circadian rhythm generation processes (3,4). Indeed, the relative inability of TTX to suppress TTFL rhythms in young adult SCN neurones (123) suggests that continuous AP production may not be the sole mechanism for intercellular communication and calcium signalling in this brain structure.

In recent years, several studies using genetically encoded calcium sensors (e.g. GCaMP3) or synthetic fluorescent probes (e.g. Fura-2) have documented the steady-state intracellular calcium $[Ca^{2+}]_i$ concentration in SCN neurones. Most (129–132), but not all (133), reported higher $[Ca^{2+}]_i$ levels during the day than at night. Furthermore, the concentration of $[Ca^{2+}]_i$ reported by these studies across the day and at night is not consistent. This ranges from 191 to 440 nM during the day and from 50 to 119 nM at night. Moreover, although, under some experimental conditions, TTX abolished the day-night difference in $[Ca^{2+}]_i$ levels (134), in others, TTX had no effect (131). Although, across experiments, animal age (neonate versus young adult tissues), species differences (rat versus mouse) and $[Ca^{2+}]_i$ detection methods (variety of genetically encoded calcium sensors used and Fura-2) may account for some of these discrepancies, there is no doubt that heterogeneity in SCN neurones and as yet unconfirmed excitability states of SCN cells (AP versus TTX insensitive depolarised DLAMOs) contribute significantly to these apparent inconsistencies. In support, from a single study by Hong *et al.* (135), TTX suppression of day-night $[Ca^{2+}]_i$ rhythms was seen in only half of SCN neurones, leaving daily cycles of this neuronal excitability measure unaffected in others.

Although the steady-state $[Ca^{2+}]_i$ concentration induced by DLAMOs is yet to be described, this electrical state may be crucial for optimal daytime calcium signalling and circadian rhythm generation by clock genes (115,119). Indeed, the predicted $[Ca^{2+}]_i$ concentration during DLAMOs (115) is consistent with the higher $[Ca^{2+}]_i$ concentration and calcium channel conductance values [440 nM (131) and > 40 pA (136), respectively] measured experimentally in SCN neurones. An extreme DLAMO-like depolarised state (~ -30 mV) is observed in cerebellar and habenular neurones (137–139) and can be induced in central amygdala neurones (140). Although we have yet to determine whether single neurones in these brain areas are cell-autonomous circadian oscillators, clock genes are expressed in these brain regions (71,81,141).

Taken together, these observations show that electrical signalling in the SCN is complex and that extreme depolarisation in central neurones extends beyond the borders of the SCN. This raises the possibility that these severe depolarised electrical states are more widespread than previously assumed and may be necessary for normal brain function. Studying these electrical states may reconcile some of the apparent inconsistencies seen in SCN neurophysiology and may extend our understanding of the all-important electrical-genetic interactions in this brain structure, which may also highlight functionality in other brain circuits containing circadian clock genes. Indeed, to date, we have no records of day-night electrical excitability and $[Ca^{2+}]_i$ levels in neuroendocrine neurones. We also have little understanding of whether and how neuroendocrine cells change their excitability to SCN and other hormonal cues over the circadian day, and whether changes in excitability of these cells can influence their genetic programmes (Fig. 3).

Conclusions and perspectives

Despite our growing understanding of the cell-autonomous processes that drive circadian rhythms in clock gene expression, our knowledge of how the molecular clockwork interacts with the membrane to drive daily changes in excitability of SCN neurones is limited. Furthermore, the nature of membrane feedback signals that convey external information to the SCN molecular clockwork is also unknown. Clock genes are found in many of the body's tissue and organ systems, and these extra-SCN clocks are considered to be vital in providing local circadian timing in a tissue-specific manner (142). Understanding the nature of the signals responsible for reciprocal connectivity between the molecular clockwork and membrane activity will unravel how circadian rhythms are generated and communicated in SCN and endocrine neurones. Furthermore, this will also reveal how circadian gene activity in neuroendocrine cells gates excitability in these cells to hormones, SCN signals and other physiological demands. This is important if we are to understand how circadian signals influence reproductive neuroendocrinology.

The observation that SCN neurones can convey circadian time cues to each other and to the body both by neural and paracrine signals demonstrates the diversity and richness of communication 'modes' in the SCN. Targeted recording of SCN neurones is also adding to our understanding that the electrical repertoire of SCN neurones is much broader than first appreciated. This highlights the challenges facing chronobiologists, physiologists, endocrinologists and neuroscientists in understanding circadian rhythm generation and communication in both the SCN and its targets. It is therefore prudent that, in addition to AP frequency measurements, multiple parameters of membrane excitability are assessed, preferably simultaneously, when interrogating the electrical state of SCN and extra-SCN neurones. This will greatly increase our understanding of the processes governing cross-talk between the master circadian clock and the neuroendocrine system, and how these signalling conduits are influenced by external cues.

Acknowledgements

I thank Professor Hugh Piggins for his critical reading of the manuscript and for his continuing support. I also would like to thank Drs N. Glossop, B. Bano-Otalora, A. Hughes and N. Nunn for their constructive comments and discussions. This work is for the Michael Harbuz Young Investigator Perspective Prize, awarded by the British Society for Neuroendocrinology, and is supported by a BBSRC grant to Hugh Piggins and David Bechtold with MDCB as co-investigator. The author has no conflicts of interest to declare.

Received 26 August 2014,
revised 25 March 2015,
accepted 26 March 2015

References

- Glossop NR. Circadian timekeeping in *Drosophila melanogaster* and *Mus musculus*. *Essays Biochem* 2011; **49**: 19–35.
- Mohawk JA, Takahashi JS. Cell autonomy and synchrony of suprachiasmatic nucleus circadian oscillators. *Trends Neurosci* 2011; **34**: 349–358.
- Colwell CS. Linking neural activity and molecular oscillations in the SCN. *Nat Rev Neurosci* 2011; **12**: 553–569.
- Brown TM, Piggins HD. Electrophysiology of the suprachiasmatic circadian clock. *Prog Neurobiol* 2007; **82**: 229–255.
- Hille B. *Ion Channels of Excitable Membranes*. Sunderland, MA: Sinauer Associates Inc., 2001.
- Aton SJ, Colwell CS, Harmar AJ, Waschek J, Herzog ED. Vasoactive intestinal polypeptide mediates circadian rhythmicity and synchrony in mammalian clock neurons. *Nat Neurosci* 2005; **8**: 476–483.
- Welsh DK, Logothetis DE, Meister M, Reppert SM. Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron* 1995; **14**: 697–706.
- Lucas RJ, Peirson SN, Berson DM, Brown TM, Cooper HM, Czeisler CA, Figueiro MG, Gamlin PD, Lockley SW, O'Hagan JB, Price LL, Provencio I, Skene DJ, Brainard GC. Measuring and using light in the melatonin age. *Trends Neurosci* 2014; **37**: 1–9.
- Lowrey PL, Takahashi JS. Genetics of the mammalian circadian system: photic entrainment, circadian pacemaker mechanisms, and posttranslational regulation. *Annu Rev Genet* 2000; **34**: 533–562.
- Hughes AT, Piggins HD. Feedback actions of locomotor activity to the circadian clock. *Prog Brain Res* 2012; **199**: 305–336.
- Roenneberg T, Kantermann T, Juda M, Vetter C, Allebrandt KV. Light and the human circadian clock. *Handb Exp Pharmacol* 2013; **217**: 311–331.
- Morin LP. Neuroanatomy of the extended circadian rhythm system. *Exp Neurol* 2013; **243**: 4–20.
- Mieda M, Sakurai T. Overview of orexin/hypocretin system. *Prog Brain Res* 2012; **198**: 5–14.
- Belle MD, Hughes AT, Bechtold DA, Cunningham P, Pierucci M, Burdakov D, Piggins HD. Acute suppressive and long-term phase modulation actions of orexin on the mammalian circadian clock. *J Neurosci* 2014; **34**: 3607–3621.
- Besing RC, Hablitz LM, Paul JR, Johnson RL, Prosser RA, Gamble KL. Neuropeptide Y-induced phase shifts of PER2:LUC rhythms are mediated by long-term suppression of neuronal excitability in a phase-specific manner. *Chronobiol Int* 2012; **29**: 91–102.
- Everett JW, Sawyer CH. A 24-hour periodicity in the 'LH-release apparatus' of female rats, disclosed by barbiturate sedation. *Endocrinology* 1950; **47**: 198–218.
- Gore AC. Diurnal rhythmicity of gonadotropin-releasing hormone gene expression in the rat. *Neuroendocrinology* 1998; **68**: 257–263.
- Plant TM. Time courses of concentrations of circulating gonadotropin, prolactin, testosterone, and cortisol in adult male rhesus monkeys (*Macaca mulatta*) throughout the 24 h light-dark cycle. *Biol Reprod* 1981; **25**: 244–252.
- Andrews WW, Ojeda SR. A detailed analysis of the serum luteinizing hormone secretory profile in conscious, free-moving female rats during the time of puberty. *Endocrinology* 1981; **109**: 2032–2039.
- Boyar RM, Wu RH, Roffwarg H, Kapen S, Weitzman ED, Hellman L, Finkelstein JW. Human puberty: 24-hour estradiol in pubertal girls. *J Clin Endocrinol Metab* 1976; **43**: 1418–1421.
- Jakacki RI, Kelch RP, Sauder SE, Lloyd JS, Hopwood NJ, Marshall JC. Pulsatile secretion of luteinizing hormone in children. *J Clin Endocrinol Metab* 1982; **55**: 453–458.
- Legan SJ, Karsch FJ. A daily signal for the LH surge in the rat. *Endocrinology* 1975; **96**: 57–62.
- Hastings MH. Neuroendocrine rhythms. *Pharmacol Ther* 1991; **50**: 35–71.
- Alleva JJ, Waleski MV, Alleva FR. A biological clock controlling the estrous cycle of the hamster. *Endocrinology* 1971; **88**: 1368–1379.
- Fitzgerald K, Zucker I. Circadian organization of the estrous cycle of the golden hamster. *Proc Natl Acad Sci USA* 1976; **73**: 2923–2927.
- Liu AC, Lewis WG, Kay SA. Mammalian circadian signaling networks and therapeutic targets. *Nat Chem Biol* 2007; **3**: 630–639.

- 27 Antunes-Rodrigues J, McCann SM. Effect of suprachiasmatic lesions on the regulation of luteinizing hormone secretion in the female rat. *Endocrinology* 1967; **81**: 666–670.
- 28 Pan JT, Gala RR. Central nervous system regions involved in the estrogen-induced afternoon prolactin surge. I. Lesion studies. *Endocrinology* 1985; **117**: 382–387.
- 29 Wiegand SJ, Terasawa E. Discrete lesions reveal functional heterogeneity of suprachiasmatic structures in regulation of gonadotropin secretion in the female rat. *Neuroendocrinology* 1982; **34**: 395–404.
- 30 de la Iglesia HO, Meyer J, Schwartz WJ. Lateralization of circadian pacemaker output: activation of left- and right-sided luteinizing hormone-releasing hormone neurons involves a neural rather than a humoral pathway. *J Neurosci* 2003; **23**: 7412–7414.
- 31 Swann JM, Turek FW. Multiple circadian oscillators regulate the timing of behavioral and endocrine rhythms in female golden hamsters. *Science* 1985; **228**: 898–900.
- 32 Poletini MO, McKee DT, Kennett JE, Doster J, Freeman ME. Knockdown of clock genes in the suprachiasmatic nucleus blocks prolactin surges and alters FRA expression in the locus coeruleus of female rats. *Am J Physiol Endocrinol Metab* 2007; **293**: E1325–E1334.
- 33 Morin LP, Allen CN. The circadian visual system, 2005. *Brain Res Rev* 2006; **51**: 1–60.
- 34 Hughes AT, Guilding C, Lennox L, Samuels RE, McMahon DG, Piggins HD. Live imaging of altered period1 expression in the suprachiasmatic nuclei of *Vipr2*^{-/-} mice. *J Neurochem* 2008; **106**: 1646–1657.
- 35 Maywood ES, Reddy AB, Wong GK, O'Neill JS, O'Brien JA, McMahon DG, Harmar AJ, Okamura H, Hastings MH. Synchronization and maintenance of timekeeping in suprachiasmatic circadian clock cells by neuropeptidergic signaling. *Curr Biol* 2006; **16**: 599–605.
- 36 Kalsbeek A, Fliers E, Hofman MA, Swaab DF, Buijs RM. Vasopressin and the output of the hypothalamic biological clock. *J Neuroendocrinol* 2010; **22**: 362–372.
- 37 Usdin TB, Bonner TI, Mezey E. Two receptors for vasoactive intestinal polypeptide with similar specificity and complementary distributions. *Endocrinology* 1994; **135**: 2662–2680.
- 38 Lehman MN, Silver R, Gladstone WR, Kahn RM, Gibson M, Bittman EL. Circadian rhythmicity restored by neural transplant. Immunocytochemical characterization of the graft and its integration with the host brain. *J Neurosci* 1987; **7**: 1626–1638.
- 39 Ralph MR, Foster RG, Davis FC, Menaker M. Transplanted suprachiasmatic nucleus determines circadian period. *Science* 1990; **247**: 975–978.
- 40 Silver R, LeSauter J, Tresco PA, Lehman MN. A diffusible coupling signal from the transplanted suprachiasmatic nucleus controlling circadian locomotor rhythms. *Nature* 1996; **382**: 810–813.
- 41 Silver R, Lehman MN, Gibson M, Gladstone WR, Bittman EL. Dispersed cell suspensions of fetal SCN restore circadian rhythmicity in SCN-lesioned adult hamsters. *Brain Res* 1990; **525**: 45–58.
- 42 Hastings MH, Brancaccio M, Maywood ES. Circadian pacemaking in cells and circuits of the suprachiasmatic nucleus. *J Neuroendocrinol* 2014; **26**: 2–10.
- 43 Bucher D, Marder E. SnapShot: neuromodulation. *Cell* 2013; **155**: 482.
- 44 Van Der Beek EM, Horvath TL, Wiegant VM, Van den Hurk R, Buijs RM. Evidence for a direct neuronal pathway from the suprachiasmatic nucleus to the gonadotropin-releasing hormone system: combined tracing and light and electron microscopic immunocytochemical studies. *J Comp Neurol* 1997; **384**: 569–579.
- 45 Mahoney MM, Smale L. Arginine vasopressin and vasoactive intestinal polypeptide fibers make appositions with gonadotropin-releasing hormone and estrogen receptor cells in the diurnal rodent *Arvicantis niloticus*. *Brain Res* 2005; **1049**: 156–164.
- 46 Van Der Beek EM, Wiegant VM, van der Donk HA, Van den Hurk R, Buijs RM. Lesions of the suprachiasmatic nucleus indicate the presence of a direct vasoactive intestinal polypeptide-containing projection to gonadotropin-releasing hormone neurons in the female rat. *J Neuroendocrinol* 1993; **5**: 137–144.
- 47 Van Der Beek EM, van Oudheusden HJ, Buijs RM, van der Donk HA, Van den Hurk R, Wiegant VM. Preferential induction of c-fos immunoreactivity in vasoactive intestinal polypeptide-innervated gonadotropin-releasing hormone neurons during a steroid-induced luteinizing hormone surge in the female rat. *Endocrinology* 1994; **134**: 2636–2644.
- 48 Smith MJ, Jiennes L, Wise PM. Localization of the VIP2 receptor protein on GnRH neurons in the female rat. *Endocrinology* 2000; **141**: 4317–4320.
- 49 Olcese J, McArdle CA, Middendorff R, Greenland K. Pituitary adenylate cyclase-activating peptide and vasoactive intestinal peptide receptor expression in immortalized LHRH neurons. *J Neuroendocrinol* 1997; **9**: 937–943.
- 50 Christian CA, Moenter SM. Vasoactive intestinal polypeptide can excite gonadotropin-releasing hormone neurons in a manner dependent on estradiol and gated by time of day. *Endocrinology* 2008; **149**: 3130–3136.
- 51 Alexander MJ, Clifton DK, Steiner RA. Vasoactive intestinal polypeptide effects a central inhibition of pulsatile luteinizing hormone secretion in ovariectomized rats. *Endocrinology* 1985; **117**: 2134–2139.
- 52 Stobie KM, Weick RF. Vasoactive intestinal peptide inhibits luteinizing hormone secretion: the inhibition is not mediated by dopamine. *Neuroendocrinology* 1989; **49**: 597–603.
- 53 Weick RF, Stobie KM. Vasoactive intestinal peptide inhibits the steroid-induced LH surge in the ovariectomized rat. *J Endocrinol* 1992; **133**: 433–437.
- 54 Ohtsuka S, Miyake A, Nishizaki T, Tasaka K, Tanizawa O. Vasoactive intestinal peptide stimulates gonadotropin-releasing hormone release from rat hypothalamus *in vitro*. *Acta Endocrinol (Copenh)* 1988; **117**: 399–402.
- 55 Van Der Beek EM, Swarts HJ, Wiegant VM. Central administration of antiserum to vasoactive intestinal peptide delays and reduces luteinizing hormone and prolactin surges in ovariectomized, estrogen-treated rats. *Neuroendocrinology* 1999; **69**: 227–237.
- 56 Kimura F, Mitsugi N, Arita J, Akema T, Yoshida K. Effects of preoptic injections of gastrin, cholecystokinin, secretin, vasoactive intestinal peptide and PHI on the secretion of luteinizing hormone and prolactin in ovariectomized estrogen-primed rats. *Brain Res* 1987; **410**: 315–322.
- 57 Harney JP, Scarbrough K, Rosewell KL, Wise PM. *In vivo* antisense antagonism of vasoactive intestinal peptide in the suprachiasmatic nuclei causes aging-like changes in the estradiol-induced luteinizing hormone and prolactin surges. *Endocrinology* 1996; **137**: 3696–3701.
- 58 Gerhold LM, Horvath TL, Freeman ME. Vasoactive intestinal peptide fibers innervate neuroendocrine dopaminergic neurons. *Brain Res* 2001; **919**: 48–56.
- 59 Mai LM, Shieh KR, Pan JT. Circadian changes of serum prolactin levels and tuberoinfundibular dopaminergic neuron activities in ovariectomized rats treated with or without estrogen: the role of the suprachiasmatic nuclei. *Neuroendocrinology* 1994; **60**: 520–526.
- 60 Poletini MO, Kennett JE, McKee DT, Freeman ME. Central clock regulates the cervically stimulated prolactin surges by modulation of dopamine and vasoactive intestinal polypeptide release in ovariectomized rats. *Neuroendocrinology* 2010; **91**: 179–188.
- 61 Egli M, Bertram R, Sellix MT, Freeman ME. Rhythmic secretion of prolactin in rats: action of oxytocin coordinated by vasoactive intestinal polypeptide of suprachiasmatic nucleus origin. *Endocrinology* 2004; **145**: 3386–3394.
- 62 Funabashi T, Shinohara K, Mitsushima D, Kimura F. Gonadotropin-releasing hormone exhibits circadian rhythm in phase with

- arginine-vasopressin in co-cultures of the female rat preoptic area and suprachiasmatic nucleus. *J Neuroendocrinol* 2000; **12**: 521–528.
- 63 Kalamatianos T, Kallo I, Goubillon ML, Coen CW. Cellular expression of V1a vasopressin receptor mRNA in the female rat preoptic area: effects of oestrogen. *J Neuroendocrinol* 2004; **16**: 525–533.
- 64 Funabashi T, Aiba S, Sano A, Shinohara K, Kimura F. Intracerebroventricular injection of arginine-vasopressin V1 receptor antagonist attenuates the surge of luteinizing hormone and prolactin secretion in proestrous rats. *Neurosci Lett* 1999; **260**: 37–40.
- 65 Palm IF, Van Der Beek EM, Wiegant VM, Buijs RM, Kalsbeek A. Vasopressin induces a luteinizing hormone surge in ovariectomized, estradiol-treated rats with lesions of the suprachiasmatic nucleus. *Neuroscience* 1999; **93**: 659–666.
- 66 Kalsbeek A, Van Heerikhuizen JJ, Wortel J, Buijs RM. A diurnal rhythm of stimulatory input to the hypothalamo-pituitary-adrenal system as revealed by timed intrahypothalamic administration of the vasopressin V1 antagonist. *J Neurosci* 1996; **16**: 5555–5565.
- 67 Palm IF, Van Der Beek EM, Wiegant VM, Buijs RM, Kalsbeek A. The stimulatory effect of vasopressin on the luteinizing hormone surge in ovariectomized, estradiol-treated rats is time-dependent. *Brain Res* 2001; **901**: 109–116.
- 68 Miller BH, Olson SL, Levine JE, Turek FW, Horton TH, Takahashi JS. Vasopressin regulation of the proestrous luteinizing hormone surge in wild-type and Clock mutant mice. *Biol Reprod* 2006; **75**: 778–784.
- 69 Coomans CP, Ramkisoensing A, Meijer JH. The suprachiasmatic nuclei as a seasonal clock. *Front Neuroendocrinol* 2014; doi:10.1016/j.yfrne.2014.11.002.
- 70 Welsh DK, Takahashi JS, Kay SA. Suprachiasmatic nucleus: cell autonomy and network properties. *Annu Rev Physiol* 2010; **72**: 551–577.
- 71 Guilding C, Piggins HD. Challenging the omnipotence of the suprachiasmatic timekeeper: are circadian oscillators present throughout the mammalian brain? *Eur J Neurosci* 2007; **25**: 3195–3216.
- 72 Mohawk JA, Green CB, Takahashi JS. Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci* 2012; **35**: 445–462.
- 73 Shieh KR. Distribution of the rhythm-related genes rPERIOD1, rPERIOD2, and rCLOCK, in the rat brain. *Neuroscience* 2003; **118**: 831–843.
- 74 von GC, Garabette ML, Kell CA, Frenzel S, Dehghani F, Schumm-Draeger PM, Weaver DR, Korf HW, Hastings MH, Stehle JH. Rhythmic gene expression in pituitary depends on heterologous sensitization by the neurohormone melatonin. *Nat Neurosci* 2002; **5**: 234–238.
- 75 Sitzmann BD, Lemos DR, Ottinger MA, Urbanski HF. Effects of age on clock gene expression in the rhesus macaque pituitary gland. *Neurobiol Aging* 2010; **31**: 696–705.
- 76 Fukuhara C, Yamazaki S, Liang J. Pineal circadian clocks gate arylalkylamine N-acetyltransferase gene expression in the mouse pineal gland. *J Neurochem* 2005; **93**: 156–162.
- 77 Bittman EL, Doherty L, Huang L, Paroskie A. Period gene expression in mouse endocrine tissues. *Am J Physiol Regul Integr Comp Physiol* 2003; **285**: R561–R569.
- 78 Zylka MJ, Shearman LP, Weaver DR, Reppert SM. Three period homologs in mammals: differential light responses in the suprachiasmatic circadian clock and oscillating transcripts outside of brain. *Neuron* 1998; **20**: 1103–1110.
- 79 Kriegsfeld LJ, Korets R, Silver R. Expression of the circadian clock gene Period 1 in neuroendocrine cells: an investigation using mice with a Per1:GFP transgene. *Eur J Neurosci* 2003; **17**: 212–220.
- 80 Olcese J, Domagalski R, Bednorz A, Weaver DR, Urbanski HF, Reuss S, Middendorff R. Expression and regulation of mPer1 in immortalized GnRH neurons. *NeuroReport* 2003; **14**: 613–618.
- 81 Abe M, Herzog ED, Yamazaki S, Straume M, Tei H, Sakaki Y, Menaker M, Block GD. Circadian rhythms in isolated brain regions. *J Neurosci* 2002; **22**: 350–356.
- 82 Guilding C, Hughes AT, Brown TM, Namvar S, Piggins HD. A riot of rhythms: neuronal and glial circadian oscillators in the mediobasal hypothalamus. *Mol Brain* 2009; **2**: 28.
- 83 Sellix MT, Egli M, Poletini MO, McKee DT, Bosworth MD, Fitch CA, Freeman ME. Anatomical and functional characterization of clock gene expression in neuroendocrine dopaminergic neurons. *Am J Physiol Regul Integr Comp Physiol* 2006; **290**: R1309–R1323.
- 84 Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Siepka SM, Hong HK, Oh WJ, Yoo OJ, Menaker M, Takahashi JS. PERIOD2: LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci USA* 2004; **101**: 5339–5346.
- 85 Hughes AT, Guilding C, Piggins HD. Neuropeptide signaling differentially affects phase maintenance and rhythm generation in SCN and extra-SCN circadian oscillators. *PLoS ONE* 2011; **6**: e18926.
- 86 Andrade-Silva J, Cipolla-Neto J, Pelicari-Garcia RA. The *in vitro* maintenance of clock genes expression within the rat pineal gland under standard and norepinephrine-synchronized stimulation. *Neurosci Res* 2014; **81–82**: 1–10.
- 87 Gillespie JM, Chan BP, Roy D, Cai F, Belsham DD. Expression of circadian rhythm genes in gonadotropin-releasing hormone-secreting GT1-7 neurons. *Endocrinology* 2003; **144**: 5285–5292.
- 88 Choe HK, Kim HD, Park SH, Lee HW, Park JY, Seong JY, Lightman SL, Son GH, Kim K. Synchronous activation of gonadotropin-releasing hormone gene transcription and secretion by pulsatile kisspeptin stimulation. *Proc Natl Acad Sci USA* 2013; **110**: 5677–5682.
- 89 Hickok JR, Tischkau SA. *In vivo* circadian rhythms in gonadotropin-releasing hormone neurons. *Neuroendocrinology* 2010; **91**: 110–120.
- 90 Zhao S, Kriegsfeld LJ. Daily changes in GT1-7 cell sensitivity to GnRH secretagogues that trigger ovulation. *Neuroendocrinology* 2009; **89**: 448–457.
- 91 Chappell PE, White RS, Mellon PL. Circadian gene expression regulates pulsatile gonadotropin-releasing hormone (GnRH) secretory patterns in the hypothalamic GnRH-secreting GT1-7 cell line. *J Neurosci* 2003; **23**: 11202–11213.
- 92 Miller BH, Olson SL, Turek FW, Levine JE, Horton TH, Takahashi JS. Circadian clock mutation disrupts estrous cyclicity and maintenance of pregnancy. *Curr Biol* 2004; **14**: 1367–1373.
- 93 Miller BH, Takahashi JS. Central circadian control of female reproductive function. *Front Endocrinol (Lausanne)* 2013; **4**: 195.
- 94 Duguay D, Cermakian N. The crosstalk between physiology and circadian clock proteins. *Chronobiol Int* 2009; **26**: 1479–1513.
- 95 de la Iglesia HO, Blaustein JD, Bittman EL. Oestrogen receptor- α -immunoreactive neurones project to the suprachiasmatic nucleus of the female Syrian hamster. *J Neuroendocrinol* 1999; **11**: 481–490.
- 96 Van Der Beek EM, Wiegant VM, van Oudheusden HJ, van der Donk HA, Van den Hurk R, Buijs RM. Synaptic contacts between gonadotropin-releasing hormone-containing fibers and neurons in the suprachiasmatic nucleus and perichiasmatic area: an anatomical substrate for feedback regulation? *Brain Res* 1997; **755**: 101–111.
- 97 Merchenthaler I, Gorcs T, Setalo G, Petrusz P, Flerko B. Gonadotropin-releasing hormone (GnRH) neurons and pathways in the rat brain. *Cell Tissue Res* 1984; **237**: 15–29.
- 98 Jennes L, Stumpf WE. LHRH-systems in the brain of the golden hamster. *Cell Tissue Res* 1980; **209**: 239–256.
- 99 Gundlach C, Kohama SG, Mirkes SJ, Garyfallou VT, Urbanski HF, Bethea CL. Distribution of estrogen receptor beta (ERbeta) mRNA in hypothalamus, midbrain and temporal lobe of spayed macaque: continued expression with hormone replacement. *Brain Res Mol Brain Res* 2000; **76**: 191–204.
- 100 Kruijver FP, Swaab DF. Sex hormone receptors are present in the human suprachiasmatic nucleus. *Neuroendocrinology* 2002; **75**: 296–305.

- 101 Su JD, Qiu J, Zhong YP, Chen YZ. Expression of estrogen receptor -alpha and -beta immunoreactivity in the cultured neonatal suprachiasmatic nucleus: with special attention to GABAergic neurons. *NeuroReport* 2001; **12**: 1955–1959.
- 102 Clancy AN, Whitman C, Michael RP, Albers HE. Distribution of androgen receptor-like immunoreactivity in the brains of intact and castrated male hamsters. *Brain Res Bull* 1994; **33**: 325–332.
- 103 Fernandez-Guasti A, Kruijver FP, Fodor M, Swaab DF. Sex differences in the distribution of androgen receptors in the human hypothalamus. *J Comp Neurol* 2000; **425**: 422–435.
- 104 Dubocovich ML, Benloucif S, Masana MI. Melatonin receptors in the mammalian suprachiasmatic nucleus. *Behav Brain Res* 1996; **73**: 141–147.
- 105 Scott FF, Belle MD, Delagrange P, Piggins HD. Electrophysiological effects of melatonin on mouse Per1 and non-Per1 suprachiasmatic nuclei neurons *in vitro*. *J Neuroendocrinol* 2010; **22**: 1148–1156.
- 106 Butler MP, Karatsoreos IN, LeSauter J, Silver R. Dose-dependent effects of androgens on the circadian timing system and its response to light. *Endocrinology* 2012; **153**: 2344–2352.
- 107 Kow LM, Pfaff DW. Suprachiasmatic neurons in tissue slices from ovariectomized rats: electrophysiological and neuropharmacological characterization and the effects of estrogen treatment. *Brain Res* 1984; **297**: 275–286.
- 108 Pevet P, Challet E. Melatonin: both master clock output and internal time-giver in the circadian clocks network. *J Physiol Paris* 2011; **105**: 170–182.
- 109 Morin LP, Cummings LA. Effect of surgical or photoperiodic castration, testosterone replacement or pinealectomy on male hamster running rhythmicity. *Physiol Behav* 1981; **26**: 825–838.
- 110 Roy EJ, Wade GN. Role of estrogens in androgen-induced spontaneous activity in male rats. *J Comp Physiol Psychol* 1975; **89**: 573–579.
- 111 Smarr BL, Morris E, de la Iglesia HO. The dorsomedial suprachiasmatic nucleus times circadian expression of Kiss1 and the luteinizing hormone surge. *Endocrinology* 2012; **153**: 2839–2850.
- 112 Khan AR, Kauffman AS. The role of kisspeptin and RFamide-related peptide-3 neurons in the circadian-timed preovulatory luteinizing hormone surge. *J Neuroendocrinol* 2012; **24**: 131–143.
- 113 Hattar S, Kumar M, Park A, Tong P, Tung J, Yau KW, Berson DM. Central projections of melanopsin-expressing retinal ganglion cells in the mouse. *J Comp Neurol* 2006; **497**: 326–349.
- 114 Brown TM, Wynne J, Piggins HD, Lucas RJ. Multiple hypothalamic cell populations encoding distinct visual information. *J Physiol* 2011; **589**: 1173–1194.
- 115 Diekmann CO, Belle MD, Irwin RP, Allen CN, Piggins HD, Forger DB. Causes and consequences of hyperexcitation in central clock neurons. *PLoS Comput Biol* 2013; **9**: e1003196.
- 116 Nitabach MN, Wu Y, Sheeba V, Lemon WC, Strumbos J, Zelensky PK, White BH, Holmes TC. Electrical hyperexcitation of lateral ventral pacemaker neurons desynchronizes downstream circadian oscillators in the fly circadian circuit and induces multiple behavioral periods. *J Neurosci* 2006; **26**: 479–489.
- 117 Nitabach MN, Blau J, Holmes TC. Electrical silencing of *Drosophila* pacemaker neurons stops the free-running circadian clock. *Cell* 2002; **109**: 485–495.
- 118 Wu Y, Cao G, Pavlicek B, Luo X, Nitabach MN. Phase coupling of a circadian neuropeptide with rest/activity rhythms detected using a membrane-tethered spider toxin. *PLoS Biol* 2008; **6**: e273.
- 119 Lundkvist GB, Kwak Y, Davis EK, Tei H, Block GD. A calcium flux is required for circadian rhythm generation in mammalian pacemaker neurons. *J Neurosci* 2005; **25**: 7682–7686.
- 120 Mizrak D, Ruben M, Myers GN, Rhrissorakrai K, Gunsalus KC, Blau J. Electrical activity can impose time of day on the circadian transcriptome of pacemaker neurons. *Curr Biol* 2012; **22**: 1871–1880.
- 121 Yamaguchi S, Isejima H, Matsuo T, Okura R, Yagita K, Kobayashi M, Okamura H. Synchronization of cellular clocks in the suprachiasmatic nucleus. *Science* 2003; **302**: 1408–1412.
- 122 Maywood ES, O'Neill JS, Chesham JE, Hastings MH. Minireview: The circadian clockwork of the suprachiasmatic nuclei—analysis of a cellular oscillator that drives endocrine rhythms. *Endocrinology* 2007; **148**: 5624–5634.
- 123 Myung J, Hong S, Hatanaka F, Nakajima Y, De SE, Takumi T. Period coding of Bmal1 oscillators in the suprachiasmatic nucleus. *J Neurosci* 2012; **32**: 8900–8918.
- 124 Schwartz WJ, Gross RA, Morton MT. The suprachiasmatic nuclei contain a tetrodotoxin-resistant circadian pacemaker. *Proc Natl Acad Sci USA* 1987; **84**: 1694–1698.
- 125 Ono D, Honma S, Honma K. Cryptochromes are critical for the development of coherent circadian rhythms in the mouse suprachiasmatic nucleus. *Nat Commun* 2013; **4**: 1666.
- 126 Kuhlman SJ, McMahon DG. Rhythmic regulation of membrane potential and potassium current persists in SCN neurons in the absence of environmental input. *Eur J Neurosci* 2004; **20**: 1113–1117.
- 127 Kuhlman SJ, Quintero JE, McMahon DG. GFP fluorescence reports Period 1 circadian gene regulation in the mammalian biological clock. *NeuroReport* 2000; **11**: 1479–1482.
- 128 Belle MD, Diekmann CO, Forger DB, Piggins HD. Daily electrical silencing in the mammalian circadian clock. *Science* 2009; **326**: 281–284.
- 129 Brancaccio M, Maywood ES, Chesham JE, Loudon AS, Hastings MH. A Gq-Ca²⁺ axis controls circuit-level encoding of circadian time in the suprachiasmatic nucleus. *Neuron* 2013; **78**: 714–728.
- 130 Enoki R, Ono D, Hasan MT, Honma S, Honma K. Single-cell resolution fluorescence imaging of circadian rhythms detected with a Nipkow spinning disk confocal system. *J Neurosci Methods* 2012; **207**: 72–79.
- 131 Ikeda M, Sugiyama T, Wallace CS, Gompf HS, Yoshioka T, Miyawaki A, Allen CN. Circadian dynamics of cytosolic and nuclear Ca²⁺ in single suprachiasmatic nucleus neurons. *Neuron* 2003; **38**: 253–263.
- 132 Irwin RP, Allen CN. Neuropeptide-mediated calcium signaling in the suprachiasmatic nucleus network. *Eur J Neurosci* 2010; **32**: 1497–1506.
- 133 Ikeda M, Yoshioka T, Allen CN. Developmental and circadian changes in Ca²⁺ mobilization mediated by GABA_A and NMDA receptors in the suprachiasmatic nucleus. *Eur J Neurosci* 2003; **17**: 58–70.
- 134 Colwell CS. Circadian modulation of calcium levels in cells in the suprachiasmatic nucleus. *Eur J Neurosci* 2000; **12**: 571–576.
- 135 Hong JH, Jeong B, Min CH, Lee KJ. Circadian waves of cytosolic calcium concentration and long-range network connections in rat suprachiasmatic nucleus. *Eur J Neurosci* 2012; **35**: 1417–1425.
- 136 Pennartz CM, de Jeu MT, Bos NP, Schaap J, Geurtsen AM. Diurnal modulation of pacemaker potentials and calcium current in the mammalian circadian clock. *Nature* 2002; **416**: 286–290.
- 137 Sakhi K, Belle MD, Gossan N, Delagrange P, Piggins HD. Daily variation in the electrophysiological activity of mouse medial habenula neurons. *J Physiol* 2014; **592**: 587–603.
- 138 Pugh JR, Raman IM. Nothing can be coincidence: synaptic inhibition and plasticity in the cerebellar nuclei. *Trends Neurosci* 2009; **32**: 170–177.
- 139 Raman IM, Gustafson AE, Padgett D. Ionic currents and spontaneous firing in neurons isolated from the cerebellar nuclei. *J Neurosci* 2000; **20**: 9004–9016.
- 140 Pape HC, Driesang RB. Ionic mechanisms of intrinsic oscillations in neurons of the basolateral amygdaloid complex. *J Neurophysiol* 1998; **79**: 217–226.
- 141 Guilding C, Hughes AT, Piggins HD. Circadian oscillators in the epithalamus. *Neuroscience* 2010; **169**: 1630–1639.
- 142 Green CB, Takahashi JS, Bass J. The meter of metabolism. *Cell* 2008; **134**: 728–742.