

2,2,2-Trifluoro-*N*-(2-iodophenyl)-acetamide

Yang Ruchun,* Zhang Hui and Cao BanPeng

Jiangxi Key Laboratory of Organic Chemistry, Jiangxi Science & Technology Normal University, Nanchang 330013, People's Republic of China
Correspondence e-mail: ouyangruchun@aliyun.com


Received 12 October 2013; accepted 31 October 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; disorder in main residue; R factor = 0.044; wR factor = 0.105; data-to-parameter ratio = 11.5.

The three F atoms in the title compound, $\text{C}_8\text{H}_5\text{F}_3\text{INO}$, are disordered over two sets of sites [relative occupancies = 0.615 (14):0.385 (14)]. In the crystal, molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming chains running along the c -axis direction. The dihedral angle between the ring and the amide group is 62.1 (3)°.

Related literature

For effects of fluorine on the properties of compounds, see: Jeschke (2004); Mueller *et al.* (2007); Purser *et al.* (2008). For the synthesis, see: Konfink *et al.* (2007).



Experimental

Crystal data

$\text{C}_8\text{H}_5\text{F}_3\text{INO}$
 $M_r = 315.02$

Tetragonal, $I4_1/a$
 $a = 15.8871$ (1) Å

$c = 15.9300$ (2) Å
 $V = 4020.7$ (6) Å³
 $Z = 16$
Mo $K\alpha$ radiation

$\mu = 3.20$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Agilent Xcalibur (Eos, Gemini) diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2011)
 $T_{\min} = 0.447$, $T_{\max} = 0.567$

6063 measured reflections
1775 independent reflections
1153 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.105$
 $S = 1.06$
1775 reflections
155 parameters

450 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.42$ e Å⁻³
 $\Delta\rho_{\min} = -0.54$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.86	2.08	2.917 (6)	163

Symmetry code: (i) $y + \frac{1}{4}, -x + \frac{3}{4}, z + \frac{1}{4}$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Science Fund of the Education Office of Jiangxi (GJJ12583) and the Bureau of Science and Technology of Nanchang City.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FF2122).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Jeschke, P. (2004). *Chembiochem*, **5**, 570–589.
- Konfink, C. C., Blank, B., Pagano, B., Gotz, N. & Knochel, P. (2007). *Chem. Commun.* **19**, 1954–1956.
- Mueller, K., Faeh, C. & Diederich, F. (2007). *Science*, **317**, 1881–1886.
- Purser, S., Moore, P. R., Swallow, S. & Gouverneur, V. (2008). *Chem. Soc. Rev.* **37**, 320–330.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2013). E69, o1749 [doi:10.1107/S1600536813029851]

2,2,2-Trifluoro-N-(2-iodophenyl)acetamide

Yang Ruchun, Zhang Hui and Cao BanPeng

1. Comment

Aromatic compounds play an important role in the pharmaceutical and agrochemical industries as important intermediates. The fluorine atom has a strong electron withdrawing character and small atomic radius. Introduction of one or more F atoms can change the physiological activity, pharmacological activity and metabolic stability properties of the compound (Jeschke, 2004; Mueller, *et al.*, 2007; Purser, *et al.*, 2008). In our study, we report an aromatic compounds containing fluorine, 2,2,2-trifluoro-N-(2-iodophenyl)acetamide, which could be used in the synthesis of many bioactive compounds.

There is a single molecule in the asymmetric unit of the title compound 2,2,2-trifluoro-N-(2-iodophenyl)acetamide, $C_8H_5F_3INO$. In the crystal, The three F atoms are disordered over two sites [relative occupancies 0.615 (14):0.385 (14)]. The crystal packing is stabilized by $N1-H1\cdots O1$ hydrogen bond that connect molecules into chains running along the *c* direction.

2. Experimental

The title compound was synthesized according to the previously reported procedure (Konfink, *et al.*, 2007).

3. Refinement

H atoms bond to N were located in a difference map and refined with $N-H = 0.86 \text{ \AA}$ and $U_{iso}(H) = 1.2U_{eq}(N)$ other H atoms attached to C were fixed geometrically and treated as riding with $C-H = 0.96 \text{ \AA}$ (methyl) or $0.93 \text{ \AA}$ (aromatic) and with $U_{iso}(H) = 1.2U_{eq}(\text{aromatic})$ or $U_{iso}(H) = 1.5U_{eq}(\text{methyl})$.

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO* (Agilent, 2011); data reduction: *CrysAlis PRO* (Agilent, 2011); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).




Figure 1

Thermal ellipsoid plot of $C_8H_5F_3INO$. Ellipsoids are drawn at the 30% probability level and H atoms are represented as small spheres of arbitrary radius.

2,2,2-Trifluoro-*N*-(2-iodophenyl)acetamide

Crystal data

$C_8H_5F_3INO$

$M_r = 315.02$

Tetragonal, $I4_1/a$

Hall symbol: $-I\ 4ad$

$a = 15.8871(1)\text{ \AA}$

$c = 15.9300(2)\text{ \AA}$

$V = 4020.7(6)\text{ \AA}^3$

$Z = 16$

$F(000) = 2368$

$D_x = 2.075\text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\text{ \AA}$

Cell parameters from 1114 reflections

$\theta = 3.1\text{--}21.8^\circ$

$\mu = 3.20\text{ mm}^{-1}$

$T = 293\text{ K}$

Block, colourless

$0.30 \times 0.20 \times 0.20\text{ mm}$

Data collection

Agilent Xcalibur (Eos, Gemini)
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2011)

$T_{\min} = 0.447$, $T_{\max} = 0.567$

6063 measured reflections

1775 independent reflections

1153 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.055$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 3.1^\circ$

$h = -17 \rightarrow 15$

$k = -18 \rightarrow 12$

$l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.105$

$S = 1.06$

1775 reflections

155 parameters

450 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0324P)^2 + 3.0148P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.54 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
I1	0.61704 (3)	0.35482 (3)	0.43754 (3)	0.0704 (2)	
N1	0.7883 (3)	0.4465 (3)	0.5004 (3)	0.0465 (12)	
H1	0.7771	0.4175	0.5446	0.056*	
O1	0.8853 (3)	0.4661 (3)	0.3977 (2)	0.0627 (13)	
F1	0.9082 (8)	0.3600 (9)	0.5849 (6)	0.095 (4)	0.615 (14)
F2	0.9342 (9)	0.3088 (7)	0.4454 (6)	0.098 (4)	0.615 (14)
F3	0.9903 (8)	0.4012 (8)	0.5219 (9)	0.109 (4)	0.615 (14)
F1'	0.8791 (11)	0.3219 (12)	0.5583 (13)	0.081 (5)	0.385 (14)
F2'	0.8837 (13)	0.2878 (9)	0.4878 (15)	0.099 (5)	0.385 (14)
F3'	0.9922 (12)	0.3650 (14)	0.4827 (12)	0.091 (5)	0.385 (14)
C1	0.6481 (4)	0.4831 (4)	0.4479 (3)	0.0468 (15)	
C2	0.5885 (4)	0.5432 (4)	0.4246 (4)	0.0594 (18)	
H2	0.5342	0.5272	0.4099	0.071*	
C3	0.6121 (5)	0.6273 (4)	0.4237 (4)	0.0649 (19)	
H3	0.5728	0.6682	0.4094	0.078*	
C4	0.6921 (5)	0.6507 (5)	0.4434 (4)	0.0634 (18)	
H4	0.7076	0.7071	0.4400	0.076*	
C5	0.7503 (4)	0.5913 (4)	0.4685 (4)	0.0543 (16)	
H5	0.8045	0.6079	0.4830	0.065*	
C6	0.7280 (4)	0.5074 (4)	0.4720 (3)	0.0440 (14)	
C7	0.8600 (4)	0.4334 (4)	0.4613 (3)	0.0431 (14)	
C8	0.9161 (5)	0.3650 (5)	0.5022 (5)	0.0577 (18)	

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.0683 (4)	0.0521 (4)	0.0909 (4)	−0.0005 (2)	−0.0240 (3)	0.0036 (3)
N1	0.052 (3)	0.047 (3)	0.041 (2)	0.010 (2)	−0.001 (2)	0.008 (2)
O1	0.055 (3)	0.086 (4)	0.047 (2)	0.011 (3)	0.007 (2)	0.018 (2)
F1	0.112 (8)	0.113 (8)	0.061 (5)	0.062 (6)	−0.008 (5)	0.010 (5)
F2	0.129 (8)	0.082 (6)	0.083 (5)	0.057 (6)	−0.001 (6)	−0.014 (5)
F3	0.077 (6)	0.118 (8)	0.133 (8)	0.006 (6)	−0.065 (6)	0.013 (6)

F1'	0.084 (9)	0.087 (10)	0.073 (9)	0.015 (7)	0.017 (7)	0.041 (8)
F2'	0.112 (10)	0.066 (8)	0.118 (10)	0.016 (7)	−0.039 (9)	0.003 (8)
F3'	0.063 (8)	0.116 (11)	0.092 (9)	0.036 (8)	0.018 (8)	0.027 (8)
C1	0.055 (4)	0.040 (3)	0.045 (3)	0.008 (3)	0.002 (3)	−0.001 (3)
C2	0.055 (4)	0.058 (4)	0.065 (4)	0.022 (3)	−0.007 (3)	−0.001 (3)
C3	0.080 (5)	0.046 (4)	0.069 (4)	0.026 (4)	−0.007 (4)	0.005 (3)
C4	0.078 (5)	0.045 (4)	0.067 (4)	0.013 (4)	0.004 (4)	−0.001 (3)
C5	0.066 (4)	0.048 (4)	0.049 (3)	0.003 (3)	0.004 (3)	−0.001 (3)
C6	0.053 (4)	0.042 (3)	0.037 (3)	0.014 (3)	0.006 (3)	0.004 (3)
C7	0.047 (4)	0.045 (4)	0.037 (3)	0.000 (3)	−0.005 (3)	−0.005 (3)
C8	0.052 (4)	0.062 (5)	0.059 (4)	0.015 (4)	−0.002 (4)	−0.004 (4)

Geometric parameters (Å, °)

I1—C1	2.103 (6)	C1—C6	1.382 (8)
N1—C7	1.316 (7)	C1—C2	1.394 (8)
N1—C6	1.433 (7)	C2—C3	1.389 (9)
N1—H1	0.8600	C2—H2	0.9300
O1—C7	1.208 (6)	C3—C4	1.361 (9)
F1—C8	1.326 (12)	C3—H3	0.9300
F2—C8	1.303 (11)	C4—C5	1.380 (9)
F3—C8	1.349 (13)	C4—H4	0.9300
F1'—C8	1.271 (16)	C5—C6	1.381 (8)
F2'—C8	1.351 (16)	C5—H5	0.9300
F3'—C8	1.248 (18)	C7—C8	1.548 (9)
C7—N1—C6	122.5 (5)	C5—C6—C1	119.6 (6)
C7—N1—H1	118.8	C5—C6—N1	119.6 (6)
C6—N1—H1	118.8	C1—C6—N1	120.8 (6)
C6—C1—C2	120.4 (6)	O1—C7—N1	128.1 (6)
C6—C1—I1	120.5 (4)	O1—C7—C8	117.6 (6)
C2—C1—I1	118.9 (5)	N1—C7—C8	114.3 (6)
C3—C2—C1	118.6 (7)	F3'—C8—F1'	128.5 (12)
C3—C2—H2	120.7	F2—C8—F1	132.0 (10)
C1—C2—H2	120.7	F2—C8—F3	105.1 (10)
C4—C3—C2	120.8 (7)	F1—C8—F3	82.9 (11)
C4—C3—H3	119.6	F3'—C8—F2'	109.1 (13)
C2—C3—H3	119.6	F1'—C8—F2'	56.9 (16)
C3—C4—C5	120.4 (7)	F3'—C8—C7	116.9 (10)
C3—C4—H4	119.8	F1'—C8—C7	114.1 (9)
C5—C4—H4	119.8	F2—C8—C7	108.4 (6)
C4—C5—C6	120.0 (7)	F1—C8—C7	113.9 (7)
C4—C5—H5	120.0	F3—C8—C7	107.6 (8)
C6—C5—H5	120.0	F2'—C8—C7	110.3 (8)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
-------------------------------	-------------	---------------------	----------------------------	-------------------------------

N1—H1 $\cdots$ O1 ⁱ	0.86	2.08	2.917 (6)	163
--------------------------------	------	------	-----------	-----

Symmetry code: (i) $y+1/4, -x+5/4, z+1/4$.