

POSTER PRESENTATION

Open Access

Tumor-specific donor lymphocyte infusion therapy with allogeneic T cells utilizing novel retrovirus vector silencing endogenous TCR expression

Hiroaki Ikeda^{1*}, Hiroaki Ueno¹, Ayumi Kawamura², Naoko Imai³, Sachiko Okamoto², Junichi Mineno², Kazutoh Takesako², Hiroshi Shiku¹

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Background

Donor Lymphocyte infusion (DLI) is a therapy for the patients with relapsed hematological malignancy after allogeneic hematopoietic stem cell transplantation. However, the development of Graft-Versus-Host Disease (GVHD) is a serious adverse event and the efficacy is limited when one needs to control the GVHD. To inhibit the development of GVHD with increased tumor-specificity of transferred allogeneic lymphocytes, here we demonstrate the development of DLI with lymphocytes engineered to express tumor-specific T cell receptor (TCR) in combination with decreased GVHD-inducing potential utilizing the novel retrovirus vector (siTCR vector) that specifically silences endogenous TCR in gene-engineered T cells.

Methods

Human PBMC were transduced with a high affinity TCR specific to a cancer/testis antigen, NY-ESO-1, by the retrovirus vector with siRNA specific to the endogenous TCR. Resulting TCR gene-transduced T cells were examined for their reactivity to allogeneic LCL by ³H uptake proliferation assay. Immunodeficient NOG mice were inoculated with a NY-ESO-1-expressing human melanoma cell line NW-MEL-38, received TCR gene-transduced T cells, and monitored for tumor growth and the development of GVHD.

Results

Human lymphocytes that were genetically engineered to express a high affinity NY-ESO-1-specific TCR with siTCR vector showed reduced expression of endogenous

TCR associated with the dramatically diminished reactivity to allogeneic lymphocytes. When administrated into NOG mice, these TCR gene-transduced T cells induced tumor regression without the development of GVHD.

Conclusions

The results here suggest that the allogeneic T cells transduced with a tumor-specific TCR by siTCR vector showed diminished GVHD potential. These T cells will be applicable to the donor lymphocytes infusion therapy after allogeneic stem cell transplantation for the treatment of hematological malignancy, providing diminished GVHD and increased tumor eradication.

Authors' details

¹Mie University Graduate School of medicine, Department of Immuno-Gene Therapy, Tsu, Japan. ²Takara Bio Inc., Otsu, Japan. ³Department of Medicine, Division of Hematology/Oncology, The Tisch Cancer Institute at Hess Center for Science and Medicine, Icahn School of Medicine at Mt. Sinai, Japan.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P20

Cite this article as: Ikeda et al.: Tumor-specific donor lymphocyte infusion therapy with allogeneic T cells utilizing novel retrovirus vector silencing endogenous TCR expression. *Journal for ImmunoTherapy of Cancer* 2014 2(Suppl 3):P20.

¹Mie University Graduate School of medicine, Department of Immuno-Gene Therapy, Tsu, Japan

Full list of author information is available at the end of the article