

The Implementation of the International Health Regulations on Vector-Borne Diseases: A Scoping Review of the Qualitative Evidence Performed Worldwide

Ana Margarida Alho^a Carlos Quelhas^b Inês Subtil^c Tiago Adrego^d
José Durão^e

^aPublic Health Medical Resident at USP Francisco George, ACES Lisboa Norte, Lisboa, Portugal; ^bPublic Health Medical Resident at USP Arnaldo Sampaio, ACES Arco-Ribeirinho, Barreiro, Portugal; ^cPublic Health Medical Resident at USP ACES Loures-Odivelas, Santo António dos Cavaleiros, Portugal; ^dPublic Health Medical Resident at USP ACES Pinhal Interior Norte, Lousã, Portugal; ^ePublic Health Medical Resident at USP Higeia, ACES Almada Seixal, Almada, Portugal

Keywords

International health regulations · Vector-borne diseases · Implementation measures · Scoping review

Abstract

Introduction: The International Health Regulations were created by the World Health Organization with the purpose of preventing the international spread of diseases. The 196 signatory countries are bound by this international agreement. In this article, we conducted a scoping review of the literature concerning the implementation and operationalization of measures at both the continental and national levels to prevent the spread of vector-borne diseases. The purpose was to understand the main strategies and policies adopted, as well as how they have been operating. **Methods:** Out of an initial search on PubMed™, SCOPUS™, and Web of Science™ using combinations of “International Health Regulations” and “vector-borne diseases”, 75 references were obtained, of which 27 were included after careful qualitative analysis. **Results:** Publication dates of included articles ranged from 1996 to

2022. Four major categories of measures were identified: (a) surveillance and epidemic intelligence; (b) Declaration of Public Health Emergency of International Concern; (c) measures in Points of Entry; and (d) vaccination status. Implemented measures were found in all continents: Africa, Oceania (Australia), Asia (China, India, Taiwan), Europe (Ireland, the Netherlands, and Mediterranean countries), North America (USA), and South America (Brazil). Yellow fever, Zika, Dengue, and Chikungunya were the most cited vector-borne diseases but Crimean-Congo hemorrhagic fever, Japanese encephalitis, Lyme disease, Malaria, Leishmaniasis, Tick-borne encephalitis, and West Nile fever were also mentioned. **Conclusion:** There are severe asymmetries across countries on the implementation of international regulations with regards to vector-borne diseases, particularly on the issue of surveillance systems. State Parties should consider the lessons learned from the pandemic and perfect their core capacities to prevent future outbreaks of infectious diseases.

© 2023 The Author(s). Published by S. Karger AG, Basel on behalf of NOVA National School of Public Health

A implementação do Regulamento Sanitário Internacional relativamente a doenças transmitidas por vetores – uma scoping review da evidência qualitativa conduzida no mundo

Palavras-Chave

Regulamento Sanitário Internacional · Doenças transmitidas por vetores · Medidas implementadas · Scoping review

Resumo

Introdução: O Regulamento Sanitário Internacional foi criado pela Organização Mundial da Saúde com o objetivo de prevenir a propagação de doenças a nível internacional. Os 196 países signatários estão vinculados a este acordo internacional. Neste artigo realizámos uma revisão de âmbito sobre a implementação e a operacionalização de medidas a nível continental e nacional para prevenir a propagação de doenças transmitidas por vetores. O objetivo foi compreender quais as principais estratégias e políticas adotadas e como têm sido implementadas. **Métodos:** A partir de uma pesquisa inicial nas bases PubMed™, SCOPUS™ e Web of Science™ utilizando as combinações dos termos “Regulamento Sanitário Internacional” e “Doenças transmitidas por vetores”, foram obtidas 75 referências, das quais 27 foram incluídas no trabalho após análise qualitativa. **Resultados:** Foram incluídos artigos de 1996 até 2022. Identificaram-se quatro grandes categorias de medidas: a) Vigilância e inteligência epidemiológica; b) Declaração de Emergência de Saúde Pública de Importância Internacional; c) Medidas nos Pontos de Entrada; e d) Situação vacinal. As medidas implementadas foram encontradas em todos os continentes: África, Oceania (Austrália), Ásia (China, Índia, Taiwan), Europa (Irlanda, Países Baixos e Países da região do Mediterrâneo) e América do Norte (Estados Unidos da América) e do Sul (Brasil). Febre amarela, Zika, Dengue e Chikungunya foram as doenças transmitidas por vetores mais referidas, tendo sido também citadas a febre hemorrágica Crimeia-Congo, a encefalite japonesa, a doença de Lyme, a malária, a leishmaniose, a encefalite transmitida por carraça e a infecção pelo vírus do Nilo Ocidental. **Conclusão:** Foram observadas grandes assimetrias entre países na implementação do Regulamento Internacional nas doenças transmitidas por vetores, particularmente, nos sistemas de vigilância. Os Estados signatários devem ter em conta as lições

aprendidas com a pandemia e aperfeiçoar as suas capacidades básicas para a prevenção de futuros surtos de doenças infeciosas.

© 2023 The Author(s). Published by S. Karger AG, Basel
on behalf of NOVA National School of Public Health

Introduction

The International Health Regulations (IHR) were created by the World Health Organization (WHO) in 1969 with the purpose of controlling the international spread of six diseases (cholera, plague, relapsing fever, smallpox, typhoid, and yellow fever). The Regulations outline the responsibilities of states regarding the notification of communicable diseases and are designed to maintain the highest level of security against the global spread of diseases with the least amount of disruption to global trade [1]. The most recent version of the IHR is from 2005. The changes in the IHR(2005) made it so that the spectrum of diseases it aims to prevent was broadened over the years. To this day, it remains a part of international law that binds 196 countries. The primary goal now is to prevent the spread of any disease that could potentially become a Public Health Emergency of International Concern (PHEIC). Each country is responsible for the national implementation of the IHR (2005). The role of the WHO is to assist countries with the implementation at the national level and to coordinate the response at the international level. Countries should be able to detect, assess, and respond to public health events, as well as report them to the WHO [2].

Vector-borne diseases (VBD) are diseases spread by “vectors”, which are living organisms that can transmit infectious pathogens between humans, or from animals to humans, and cause illnesses [3]. These diseases include several mosquito-borne viral infections with a significant public health impact, such as West Nile, Dengue, Chikungunya, Japanese encephalitis, Yellow fever, Rift Valley fever, and Zika virus, as well as parasitic infections such as Malaria and Lymphatic filariasis. Other well-established and important VBD are those transmitted by ticks, such as Lyme Borreliosis, Crimean-Congo hemorrhagic fever, Tick-borne encephalitis, and Typhus (the last, also transmitted by mites, lice, and fleas). Other relevant vector-borne diseases include those transmitted by triatomine bugs (Chagas disease), by flies (Onchocerciasis and Human African Trypanosomiasis), by

snails (Schistosomiasis), or by phlebotomine sandflies, such as cutaneous, mucocutaneous, and visceral Leishmaniasis. Vector avoidance strategies are therefore critical to the prevention of these infections. According to the WHO's Global Vector Control Response 2017–2030, 80% of the world's population is at risk of one VBD, and more than half are at risk of two or more VBD. Additionally, VBD account for 17% of the global burden of communicable diseases, with over 700,000 deaths annually. VBD has recently expanded into new regions, with illnesses that were previously only seen in tropical and subtropical regions becoming prevalent in temperate regions. Climate and environmental changes, increased worldwide travel and trade, changes in land use patterns such as deforestation, unplanned urbanization, inadequate solid waste management, and lack of piped water are all major factors contributing to the emergence or re-emergence of VBD [4].

The interplay between demographic, social, and environmental factors has resulted in the geographic dispersion and extended duration of transmission seasons for VBD, emphasizing the immediate need for a comprehensive approach to vector control. In this article, we aim to perform a scoping review of the literature on the implementation and operationalization of measures to control the spread of VBD within the scope of the IHR. The focus of this article was specifically on surveillance at Points of Entry (PoE), epidemic intelligence, entomological surveillance, and control of the vaccination status of travelers at airports, ports, and ground crossings [5]. We searched for articles within the scope of the implementation of such measures at both the continental and national levels. The purpose was to understand how these measures are being implemented and shed light on the successes and struggles of these implementations in different countries around the globe.

Materials and Methods

Review Question

The specific review question aimed to characterize the evidence of the IHR measures implemented to control the occurrence of VBD worldwide.

Objectives

1. To identify the effects of existing IHR strategies and interventions that have been developed for monitoring, restricting, and controlling the occurrence of VBD.

2. To classify strategies and interventions that have been implemented, according to their suitability and efficacy in controlling and preventing the development of VBD in human populations.

Type of Review and Justification

We conducted a scoping review as the body of literature concerning the implementation of the IHR on VBD has not yet been comprehensively reviewed. A second reason was the heterogeneous nature of content and measures found, which was not amenable to a more precise systematic review. Therefore, the authors opted for a scoping review as it provides a preliminary assessment of the potential scope and size of the available literature, allowing for the identification of the nature and extent of research evidence.

Review Process

In order to seek and map the evidence in such a broader approach, the authors standardized the conduct and reporting of this review taking into account the guidance developed by the members of the Joanna Briggs Institute and Collaborating Centres and the Search, Appraisal, Synthesis and Analysis (SALSA) Framework [6, 7]. Prior to the review itself, a scoping review protocol was developed where the objectives and methods were predefined. The search strategy was comprehensive to identify both published and unpublished (grey literature) references. We searched relevant databases, namely SCOPUS™, Web of Science™, and PubMed™, to search for articles published since 1990.

Combinations of the terms: "International Health Regulations," and "vector-borne diseases," contained in the title, abstract, and index terms were searched. Our final queries for the search were: "vector-borne" AND ("international health regulations" OR "IHR") on SCOPUS™, "vector-borne" AND ("international health regulations" OR "IHR") on Web of Science™ and ("International Health Regulations" OR "IHR") AND "Vector-borne" (Mesh) on PubMed™.

Articles and reports published in English or Portuguese were included. Studies written in other languages were identified and excluded during eligibility screening (see Table 1).

The primary screening in PubMed™ yielded 60 articles, with 16 detected in SCOPUS™ and 11 in Web of Science™. All results were exported into a Microsoft Excel file. After removing duplicates, 75 results were considered. The primary screening was performed independently by two researchers and conducted on the title and abstract of each article. The answers were compared, and disagreements were discussed to solve any discrepancies through consensus or a third researcher. Articles that did not meet one or more of the inclusion criteria were excluded ($n = 42$). Articles that met any combination or were unclear to one of the listed criteria proceeded to secondary screening. Based on defined criteria, 33 articles were included for full-text assessment. The secondary screening was performed on the full-text articles independently by two researchers. The consensus was obtained using the same methods as on the primary screening. Full-text careful qualitative analysis was performed [8–40], with 6 studies being excluded [9, 16, 17, 20, 29, 37]. Overall, 27 articles met all the described criteria and were included in this scoping review (shown in Fig. 1).

Table 1. Eligibility criteria for population, interventions, comparisons, outcomes, timing, setting, study design, and publication language

	Inclusion criteria	Exclusion criteria
Population	Geographical areas at risk for VBD	None
Intervention	A. Surveillance (syndromic surveillance, vector surveillance, epidemic intelligence) B. PHEIC C. Measures in Points of Entry D. Vaccination status	Complementary and alternative interventions, interventions outside of the scope of the IHR
Comparison	Standard measures of IHR	None
Outcomes	Prevalence of vectors detected during entomological collections Prevalence of human cases of VBD Number of countries affected Vaccine coverage	Studies outside the scope of the outcomes established
Timing	Studies published since 1990	None
Settings	International Health	None
Study design	Original data Any sample size Observational studies Case series Cross-sectional studies Qualitative studies Narrative reviews Systematic reviews or meta-analyses	Non-original data (e.g., book chapters, letters, or erratum) Surveys In vitro studies
Language	Studies published in English and Portuguese	Other foreign language studies

IHR, International Health Regulations; PHEIC, Public Health Emergency of International Concern; VBD, Vector-borne diseases.

For each included study, the following data were extracted:

1. Strategies/interventions on VBD applied in accordance with the IHR.
2. Effects of interventions directed toward the prevalence of human cases regarding VBD or toward the abundance of vectors.
3. Data related to other characteristics, study period, country of origin, study population, type of disease for which the IHR was applied.

Congo hemorrhagic fever, malaria, Japanese encephalitis, West Nile fever, and Leishmania) as others. A quotation is provided to exemplify each category.

General Characteristics of the Studies Included

Of the included articles, sixteen were research articles [8, 12, 14, 15, 17, 23, 25, 27, 30–33, 35, 36, 38, 40], five were review articles [10, 11, 14, 22, 39], and six were other types of publications (policy analysis, commentaries, and editorial) [19, 21, 24, 26, 28, 34]. The year of included articles ranged from 1996 [22] to 2022 [8, 31, 34] with 74% ($n = 20$) being published in 2016 or after [8, 10–12, 15, 18, 19, 21, 23, 24, 27, 30–34, 36, 38, 40]. Seven studies were conducted in India [19, 28, 30–33, 36], three in China [8, 12, 23], three in the Mediterranean area [27, 34, 40], two in Brazil [21, 25], two in Africa [22, 38], two in the Netherlands [8, 24], and two in Asia-Pacific, Australia, and French Polynesia [15, 21]. There was also one study from Ireland [11], one from the USA [26], one from Indonesia [24], one from Taiwan [18], and another one from several developing countries [35]. Three studies/articles had no specific geographic focus [10, 14, 39].

Results

An overview of the selected articles is provided in Table 2. For each article, author, year of publication, category(ies) of IHR measures type of publication, diseases, and geographic area are indicated. A more detailed description according to each of the four IHR measures categories is highlighted in Table 3. For simplification, Table 3 lists results for the four most prevalent diseases (Yellow fever, Zika, Dengue, and Chikungunya) and groups the rest (Lyme disease, Tick-borne encephalitis, Crimean-




Fig. 1. Flow diagram describing the steps and number of studies selected for inclusion during the scoping review process.

Some of the articles were related to specific diseases, mainly Yellow fever [8, 10, 12, 22, 26, 30–32, 38], Zika [8, 11, 18, 21, 25, 27, 30–32, 38], Dengue [8, 19, 24, 28, 30–33, 35, 38], and Chikungunya [8, 19, 24, 28, 30–33, 35, 38]. All the selected articles included IHR implementation measures to strengthen prevention and response to public health events. The main categories of measures were related to surveillance (whether syndromic or vector surveillance or epidemic intelligence), Declaration of PHEIC, measures in Points of Entry (PoE, such as ports and airports), and vaccination status.

Surveillance and Epidemic Intelligence

It is emphasized in the IHR that epidemiological and syndromic surveillance of VBD is crucial to expedite national and international reporting. There are risk characterization guidelines in place for several vectors which help countries identify specific triggers for scaling up or down surveillance and public health activities, such as monitoring geographic spread and distribution and detecting changing transmission patterns or atypical clinical presentations [21]. However, there have been reported operational difficulties due to barriers within countries. Among these barriers, one should consider

Table 2. Overview of the selected articles included in this scoping review

Ref.	Author (year of publication)	IHR measure category	Type of publication	Diseases included	Geographic area
[8]	Onstwedder et al. (2022)	Surveillance, PoE	Research article	Zika, Chikungunya, Dengue, Yellow fever, and other VBD	Beijing and the Netherlands
[10]	Simons and Patel (2016)	PHEIC, vaccination	Review article	Yellow fever	Global
[11]	Glynn et al. (2016)	PoE	Review article	Ebola, Zika	Ireland
[12]	Lock-Wah-Hoon et al. (2020)	Surveillance	Research article	Yellow fever	Beijing
[14]	Patel and Simons (2013)	Vaccination	Review article	Yellow fever	Global
[15]	Bennett and Carney (2017)	PHEIC	Research article	VBD and others	Asia-Pacific and Australia
[18]	Ho et al. (2017)	Surveillance, PoE	Research article	Zika	Taiwan
[19]	Pilot et al. (2020)	Surveillance	Qualitative policy analysis	Dengue	India
[21]	Singer et al. (2017)	PHEIC, Surveillance	Policy analysis	Zika	Brazil and French Polynesia
[22]	Robertson et al. (1996)	Vaccination	Review article	Yellow fever	Senegal and Ivory Coast
[23]	Vlieg et al. (2017)	Surveillance	Research article	Dengue, Chikungunya, Zika, others	China and the Netherlands
[24]	Satoto et al. (2018)	PoE	Research article	Dengue	Indonesia
[25]	Bueno FTC (2017)	PHEIC	Research article	Zika	Brazil, South America
[26]	Wilder-Smith et al. (2008)	Vaccination	Editorial	Yellow fever, others	USA
[27]	Minh et al. (2016)	Surveillance, PoE	Research article	Zika	Mediterranean
[28]	Sharma et al. (2005)	PoE	Research article	Dengue	India
[30]	Regu et al. (2021)	PoE	Research article	Dengue, Chikungunya, Zika, and Yellow fever	India
[31]	Rajendran et al. (2022)	PoE	Research article	Dengue, Chikungunya, Zika, and Yellow fever	India
[32]	Sharma et al. (2020)	PoE	Research article	Dengue, Zika, and Yellow fever	India
[33]	Sharma et al. (2021)	PoE	Research article	Dengue	India
[34]	Al Ariqi et al. (2022)	Surveillance	Commentary	Emerging and Epidemic Prone Infectious Diseases	Eastern Mediterranean
[35]	May et al. (2009)	Surveillance	Research article	Malaria, Dengue, other VBD, Foodborne illness, respiratory illness, sexually transmitted infections	Developing countries

Table 2 (continued)

Ref.	Author (year of publication)	IHR measure category	Type of publication	Diseases included	Geographic area
[36]	Aggrawal et al. (2020)	Surveillance	Research article	VBD and others	India
[38]	Williams et al. (2021)	Surveillance	Research article	Dengue, Chikungunya, Zika, West Nile fever, Yellow fever, Malaria, Leishmania	Africa
[39]	Mouchtouri et al. (2010)	PoE	Review article	VBD and others	Global
[40]	Drakou et al. (2020)	PoE	Research article	None in particular	Cyprus

PoE, Points of Entry; PHEIC, Public Health Emergency of International Concern; VBDs, Vector-borne diseases.

factors such as poor data quality and poor information flow, political barriers, and weak surveillance infrastructure [12]. International cooperation and sharing of best practices can be of the utmost importance for the implementation of the IHR. Comparing surveillance systems allows each country to learn valuable lessons and effectively address international challenges [8]. These recommendations are in accordance with Articles 5–10 of the IHR, 2005.

Declaration of PHEIC

The Emergency Committee of the WHO may declare a PHEIC under the IHR. In May 2014 this was the case for the wild poliovirus following the acknowledgment that travelers contributed to the spread of the virus. Recommendations were made to the affected countries, and measures were taken to reduce the potential spread [10]. A PHEIC declaration coordinates timelines and mechanisms for international response, including protocols for an outbreak and case limitation [21]. These recommendations are in accordance with Articles 12–14 of the IHR, 2005.

Measures at PoE

The outbreak of the Ebola Virus Disease in 2014–2015 and the emergence of Zika in 2016, both considered PHEIC, provided an impetus for countries to review their PoE preparedness [11]. Given the rapid global growth of human connectivity, the entry and establishment of VBD in seaports and airports pose a potential threat of disease outbreaks and epidemics. To address this, the IHR recommends sanitary standards at international borders and PoE, including knowledge awareness for PoE authorities, early de-

tection of potentially infected persons, dissemination of health information for travelers and protocols for assessment and case management [11]. Vector surveillance at these sites includes specific parameters. For instance, the IHR specifically states that if specific vectors are found within a 400-m range of PoE sites, public health authorities must be notified so that control measures may be applied [8, 31, 32]. These recommendations are in accordance with Articles 19–29 of the IHR, 2005.

Vaccination Status

Some VBD are vaccine-preventable. Yellow fever is one of those examples. This vaccine can be recommended for personal protection or it may be a mandatory requirement for entry into certain country. The WHO keeps a record of the at-risk countries and the national requirements for entry. As proof of vaccination under the IHR, an International Certificate of Vaccination or Prophylaxis (ICVP) is issued at the time of vaccination [10]. Policymakers at a national level are sometimes faced with the lag that happens between new scientific evidence and the lack of concurrent amendments to the IHR, such as with the shift from the 10-yearly revaccination to the declaration of life-long protection from a single dose of the Yellow fever vaccine. Nevertheless, risk assessment and risk management should always be considered by health practitioners [12]. The implementation of the IHR and the ICVP is crucial to prevent the international spread of infectious diseases and for the coordinated global response to public health events [26]. These recommendations are in accordance with Articles 18, 23, and 31 of the IHR, 2005.

Table 3. Characterization of the studies included in this scoping review

Measure category	Description	Studies included, n	Publication type [n]	Publication year [n]	Geographic area	Included diseases [n]	Example	References
Surveillance and epidemic intelligence	Countries' core capacity requirements with regard to disease and vector surveillance, including reporting, notification, verification, response, and collaboration	11	Research article [8], Policy analysis [2], Commentary [1]	2009 [1], 2016 [1], 2017 [3], 2020 [3], 2021 [1], 2022 [2]	Africa (several countries, developing countries), Asia (China, Taiwan, India), Europe (the Netherlands), Oceania (French Polynesia), South America (Brazil), Mediterranean region, and Eastern Mediterranean region	Chikungunya [3], Dengue [5], Yellow fever [3], Zika [6], Others [6]	"A strong entomological surveillance system is crucial for obtaining data on the distribution of vector(s), the extent and types of breeding habitats, and the intensity and seasonal fluctuations of breeding of mosquitoes. Entomological sampling methods to assess Aedes population density and evaluate the control interventions have been largely implemented, especially in South-East Asian countries. There is a need to introduce and adapt those methods in countries through standardized protocols, including traditional Stegomyia indices to be used in outbreak prevention." [27]	[8, 12, 18, 19, 21, 23, 27, 34-36, 38]
PHEIC declaration	Public health measures under the scope of a PHEIC declaration for a VBD.	4	Review article [1], Research article [2], Policy analysis [1]	2016 [1], 2017 [3]	Oceania (Australia, French Polynesia), South America (Brazil), Asia-Pacific region, Global	Chikungunya [1], Dengue [1], Yellow fever [2], Zika [3], Others [2]	"With the PHEIC declaration of neurologic complications associated with Zika virus infection, the WHO will follow its surveillance algorithm in accordance with International Health Regulations protocol and procedures." [21]	[10, 15, 21, 25]

Table 3 (continued)

Measure category	Description	Studies included, n	Publication type [n]	Publication year [n]	Geographic area	Included diseases [n]	Example	References
Points of Entry	Countries' core capacity requirements with specific regards to VBD control at airports, ports, and ground crossings	12	Review article [2], Research article [10]	2005 [1] 2010 [1], 2016 [2], 2017 [1], 2018 [1], 2020 [2], 2021 [2], 2022 [2]	Asia (China, India, Indonesia, Taiwan), Europe (Cyprus, Ireland, the Netherlands), Mediterranean region, Global	Chikungunya [4], Dengue [8], Yellow fever [5], Zika [8], Others [3]	'The IHR envisages that all international seaports and airports should be kept free from all types of mosquito vectors for a minimum distance of 400 m around the perimeter of the ports in order to achieve the ultimate aim of public health security. This necessitates periodic and regular vector surveillances and adoption of appropriate vector control measures in and around the ports.' [31]	[18, 11, 18, 24, 27, 28, 30–33, 39, 40]
Vaccination status	Measures pertaining to vaccination and prophylaxis recommendations for VBD, as well as related certificates	4	Review article [3], Editorial [1]	1996 [1], 2008 [1], 2013 [1], 2016 [1]	Africa (Senegal, Ivory Coast), North America (USA), Global	Yellow fever [4], others [2]	'An important distinction is that an individual country's requirements are not designed to protect individual travelers from a risk of yellow fever, but rather to protect the country from the introduction of yellow fever by the traveler.' [26]	[10, 14, 22, 26]

Discussion

Literature reviews play a significant role in healthcare practice. Evidence-based practice is gaining recognition as the benchmark for care [6]. Consequently, reviews of primary research are becoming more frequent as a useful tool for researchers, clinicians, and decision-makers. Scoping reviews are a relatively new methodology, generally performed to synthesize research evidence, and clarify definitions and conceptual boundaries within a determined field or topic, particularly when the body of certain literature has not yet been comprehensively reviewed. They allow one to perform a summary of research findings (quantitative and/or qualitative), identify research gaps, and make recommendations for future research topics.

The perceived strengths of a scoping review are the characteristics they share with the systematic reviews, namely their systematic, replicable, and transparent methodology. On the other hand, the perceived weaknesses and limitations of scoping reviews are the fact that they cannot be regarded as a final output due to potential limitations in their rigor and duration. Typically, they do not include a process of quality assessment, and consequently, there is the risk of including studies by their existence and not exactly by their intrinsic quality. Subsequently, their findings cannot be used to recommend policy/practice, as a formal assessment of the methodological quality of the included studies is generally not performed [7].

Overall, this scoping review stresses the relevance of IHR in a world increasingly susceptible to VBD. It is expected that VBD outbreaks will become more common and that new PHEIC will be declared as a result of climate change, globalization, tourism, and increased movement of goods across international borders. This will require a high level of preparedness from all countries, which should ensure that their core capacities are updated and robust.

Vaccination is the mainstay for the prevention of events caused by vaccine-preventable VBD (e.g., Yellow fever), and the main purpose of requiring travelers to be vaccinated is for public health benefit, i.e., to prevent the introduction of the pathogen in the destination country [10]. There is potential for improvement in keeping the IHR in accordance with the best available evidence and allowing the law to reflect the best clinical practice [14].

The formal declaration of a PHEIC constitutes an important tool for international cooperation and coordinated response to events and allows the countries to follow procedures already in place for incident response. There are, however, other interests influencing the im-

plementation of the IHR and a general need to strengthen and build capacity in the institutions responsible for responding to public health threats at a national, regional, and global level [15]. State Parties do not always follow WHO recommendations with regard to a PHEIC, such as when countries like Russia and China ceased pork meat imports from countries affected by Influenza H1N1 in 2009, despite no evidence linking pork to the pathogen. Likewise, several countries imposed considerable travel barriers contrary to WHO guidelines during the Ebola outbreak in 2014, with the USA, for example, applying 21-day quarantines on healthcare workers returning from afflicted countries [15]. Modifications to the formal processes described in the IHR may also be relevant to respond to specific and emerging threats in a scalable, adaptable, and flexible way [21].

Surveillance and epidemic intelligence are mandated by the IHR as important tools for the early detection of outbreaks. Syndromic surveillance extends the detection capability beyond confirmed cases of specific diseases characteristic of typical epidemic surveillance and should be built on existing public health surveillance infrastructure [35]. Across countries, epidemic surveillance varies in capabilities according to the resources available to each country and is not uniformly applied. For example, clinical surveillance in mass gathering events enables early disease detection and response [36] as well as provides early detection, even if the causative agent is yet unknown [35]. Comparing surveillance systems allows each country to learn valuable lessons. Nevertheless, differences in preparedness and response are rooted in specific national VBD epidemiology and surveillance structure. China, for instance, relies on syndromic surveillance first, while the Netherlands focus on laboratory-confirmed diagnosis [8].

Vector surveillance at PoE encompasses the most common implementation measures found in literature, with clear standards required by the IHR. These provide early warnings for the introduction of vectors or diseases at critical points (e.g., airports and seaports) [30–32].

An important gap found in the literature that runs across different studies is the systematic underinvestment in the necessary infrastructure to conduct the activity required to uphold the IHR. From lack of dedicated transport for field teams to insufficient information technology IT equipment, underfunded laboratories, and not enough human resources, or financial support to conduct surveillance activities, there is a clearly identified need to strengthen the institutions responsible for the implementation of the IHR at all levels of governance. This systemic lack of investment inevitably leads to marked differences in the implementation of the IHR across countries, with North

America and Europe investing more heavily in syndromic surveillance. Member states of the WHO African and Southeast Asia regions, on the other hand, face logistical, human, and financial challenges that countries from richer regions do not face, which can undermine efforts to adequately respond to such threats [15, 38].

Thus, this scoping review highlights the persistent and stark asymmetries between countries with regard to the full implementation of IHR. The fact that most of the vector-borne pathogens are endemic to the global south may be a limitation to the scope of this review. Moreover, the restriction of sources to the English and Portuguese languages may explain the relative lack of coverage of certain geographic regions, such as Southeast Asia. This could also be a reason why India is overrepresented in the literature selected in this review.

To conclude, this scoping review offers a summary of research findings regarding the implementation of the IHR to control the occurrence and transmission of VBD worldwide. To the authors' knowledge, this review represents the first assessment of available literature on this topic, as the implementation of the IHR on VBD has not yet been comprehensively reviewed. Indeed, the relative lack of robust, systematic literature found on this topic is cause for alert. The IHR measures are inconsistently referenced across pathogens, diseases, countries, and continents, often with different outcomes and reporting standards. Whether concerning surveillance, vaccination status, PoE, or PHEIC declaration, the IHR effectively lays the groundwork, albeit in need of a careful revision, and those tools and know-how should be approached in a consistent, responsible way, regardless of location and politics. Considering global phenomena, like climate change, and the consequent increasing spread of vectors and VBD to new geographical areas (especially higher latitudes and altitudes), adopting IHR is essential to ensure the highest level of protection against the spread of diseases with the least amount of disruption to the global movement of goods and people.

Further studies should focus on the preparedness level of countries in the northern hemisphere, as they are likely to face an increased risk of VBD outbreaks. Specifically,

surveillance systems should not only reflect the recommendations stated by the IHR but must allow for a flexible and continuous flow of information across borders, so that an international response can quickly and effectively be implemented between governments.

The continued relevance of the IHR was widely acknowledged during the COVID-19 pandemic, an event that strained State Parties' core capacities to their limits and allowed for successes and shortcomings of the IHR to be better identified. The continued sense of urgency regarding recent outbreaks of Poliomyelitis, Dengue, Malaria, and Ebola, as well as the unavoidable recognition of the interdependency of nations regarding their populations' health, should be the driving force behind an effective implementation of IHR measures.

Statement of Ethics

Ethics approval was not required for this study.

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

Funding Sources

No funding was received for this study.

Author Contributions

All authors designed the experiment, analyzed the data, and wrote the manuscript. Ana Margarida Alho and José Durão reviewed and prepared the manuscript. All authors approved the final version of the manuscript.

Data Availability Statement

All data generated or analyzed during this study are included in this article. Further inquiries can be directed to the corresponding author.

References

- 1 World Health Organization. *International health regulations*. Geneva: WHO; 1969.
- 2 World Health Organization. *International health regulations*. 2nd ed. Geneva: WHO; 2005.
- 3 World Health Organization. *Vector-borne diseases*. Geneva: WHO; 2020.
- 4 World Health Organization. *Global vector control response 2017–2030*. Geneva: WHO; 2017.
- 5 Huntington M, Allison J, Nair D. Emerging vector-borne diseases. *Am Fam Physician*. 2016;94(7):551–7.
- 6 Peters MDJ, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc*. 2015;13(3):141–6.
- 7 Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J*. 2009;26(2):91–108.
- 8 Onstwedder C, Lock-Wah-hoon J, Van Dorp S, Braks M, Van Asten L, Zheng Y, et al. Comparing vector-borne disease surveillance and response in Beijing and The Netherlands. *Ann Glob Heal*. 2022;88(1):59.

- 9 Queyriaux B, Pradines B, Hasseine L, Coste S, Rodriguez P, Coffinet T, et al. Airport malaria. *Presse Med*. 2009;38(7–8):1106–9.
- 10 Simons H, Patel D. International Health Regulations in practice: focus on yellow fever and poliomyelitis. *Hum Vaccin Immunother*. 2016;12(10):2690–3.
- 11 Glynn RW, Boland M; HSE Port Health Groups, Ireland. Ebola, Zika and the international health regulations: implications for port health preparedness. *Global Health*. 2016;12(1):74.
- 12 Lock-Wah-hoon J, Zheng Y, Braks M, van Asten L, Liu Q, Sushama P, et al. Exploring vector-borne disease surveillance and response systems in Beijing, China: a qualitative study from the health system perspective. *Int J Environ Res Public Heal*. 2020;17(22):e8512.
- 13 Steffen R, Baños A, DeBernardis C. Vaccination priorities. *Int J Antimicrob Agents*. 2003;21(2):175–80.
- 14 Patel D, Simons H. Yellow fever vaccination: is one dose always enough? *Travel Med Infect Dis*. 2013;11(5):266–73.
- 15 Bennett B, Carney T. Public health emergencies of international concern: global, regional, and local responses to risk. *Med Law Rev*. 2017;25(2):223–39.
- 16 Gostin LO. Our shared vulnerability to dangerous pathogens. *Med Law Rev*. 2017; 25(2):185–99.
- 17 Katz R, Allen H. Domestic understanding of the revised international health regulations. *Public Health Rep*. 2009;124(6):806–12.
- 18 Ho LL, Tsai YH, Lee WP, Liao ST, Wu LG, Wu YC. Taiwan's travel and border health measures in response to Zika. *Heal Secur*. 2017;15(2):185–91.
- 19 Pilot E, Murthy GVS, Nittas V. Understanding India's urban dengue surveillance: a qualitative policy analysis of Hyderabad district. *Glob Public Health*. 2020;15(11):1702–17.
- 20 Kindhauser MK, Allen T, Frank V, Santhana RS, Dye C. Zika: the origin and spread of a mosquito-borne virus. *Bull World Health Organ*. 2016;94(9):675–86C.
- 21 Singer L, Vest KG, Beadling CW. Zika virus: a review of management considerations and controversies at six months. *Disaster Med Public Health Prep*. 2017;11(3):279–84.
- 22 Robertson SE, Hull BP, Tomori O, Bele O, LeDuc JW, Esteves K. Yellow fever: a decade of reemergence. *JAMA*. 1996;276(14):1157–62.
- 23 Vlieg WL, Fanoy EB, Van Asten L, Liu X, Yang J, Pilot E, et al. Comparing national infectious disease surveillance systems: China and The Netherlands. *BMC Public Health*. 2017;17(1):415.
- 24 Satoto TBT, Listyantanto A, Agustjahani SD, Josef HK, Widartono BS. Vertical transmission of dengue virus in the Yogyakarta airport area. *Environ Health Prev Med*. 2018;23(1):22.
- 25 Bueno FTC. Health surveillance and response on a regional scale: a preliminary study of the Zika virus fever case. *Cien Saude Colet*. 2017; 22(7):2305–14.
- 26 Wilder-Smith A, Hill DR, Freedman DO. The revised International Health Regulations (2005): impact on yellow fever vaccination in clinical practice. *Am J Trop Med Hyg*. 2008; 78(3):359–60.
- 27 Minh NNT, Huda Q, Asghar H, Samhouri D, Abubakar A, Barwa C, et al. Zika virus: no cases in the Eastern Mediterranean Region but concerns remain. *East Mediterr Health J*. 2016;22(5):350–5.
- 28 Sharma SN, Kumar S, Das BP, Thomas TG, Kumar K, Katyal R, et al. Entomological indices of Aedes aegypti at some international airports and seaports of southern India: a report. *J Commun Dis*. 2005;37(3):173–81.
- 29 Hoch M, Wieser A, Löscher T, Margos G, Pürner F, Zühl J, et al. Louse-borne relapsing fever (*Borrelia recurrentis*) diagnosed in 15 refugees from northeast Africa: epidemiology and preventive control measures, Bavaria, Germany, July to October 2015. *Euro Surveill*. 2015;20(42):30046.
- 30 Regu K, Rajendran R, Rajendran A, Anusree SB, Tamizharasu W, Poduval R. Entomological surveillance of vectors of yellow fever, dengue, Chikungunya and Zika virus at Kannur International Airport, Kerala, India. *Int J Curr Microbiol App Sci*. 2021;10(8):728–35.
- 31 Rajendran R, Regu K, Anusree SB, Rajendran A, Tamizharasu W, Sharma SN. Need for strict adherence to international health regulations: entomological surveillance at mangalore international airport. *J Commun Dis*. 2022;54(2):61–6.
- 32 Sharma SN, Singh R, Kumawat R, Singh SK. Guidelines for vector surveillance and its control at international airports and ports in India. *J Commun Dis*. 2020;52(1):38–60.
- 33 Sharma SN, Sn S, Singh SK, Singh SK, Sk S. Assessment of the receptivity of Dengue vector breeding through pre- and post-monsoon vector surveillance at some international airports in India. *J Commun Dis*. 2021;53(3):16–22.
- 34 Al Ariqi I, Buliva E, Chughtai AA, Barakat A, Kodama C, Khan W, et al. How far are we? National preparedness and response capacities for emerging infectious disease outbreaks in the WHO Eastern Mediterranean Region. *BMJ Glob Heal*. 2022;7(Suppl 4): e009826.
- 35 May L, Chretien JP, Pavlin JA. Beyond traditional surveillance: applying syndromic surveillance to developing settings—opportunities and challenges. *BMC Public Health*. 2009;9(1):242.
- 36 Aggrawal V, Dikid T, Jain SK, Pandey A, Khasnabish P, Choudhary S, et al. Disease surveillance during a large religious mass gathering in India: the Prayagraj Kumbh 2019 experience. *Int J Infect Dis*. 2020;101: 167–73.
- 37 M'ikanatha NM. Infectious disease surveillance and the international health regulations. 2nd ed. In: M'ikanatha NM, Lynfield R, van Beneden CA, de Valk H, editors. *Infectious disease surveillance*. London: Wiley-Blackwell; 2013. p. 62–80.
- 38 Williams GS, Impouma B, Mboussou F, Lee TMH, Ogundiran O, Okot C, et al. Implementing epidemic intelligence in the WHO African region for early detection and response to acute public health events. *Epidemiol Infect*. 2021;149:e261.
- 39 Mouchtouri VA, Nichols G, Rachiotis G, Kremastinou J, Arvanitoyannis IS, Riemer T, et al. State of the art: public health and passenger ships. *Int Marit Health*. 2010;61(2):49–98.
- 40 Drakou K, Nikolaou T, Vasquez M, Petric D, Michaelakis A, Kapranas A, et al. The effect of weather variables on mosquito activity: a snapshot of the main point of entry of Cyprus. *Int J Environ Res Public Health*. 2020; 17(4):1403.