

# Voltage-dependent processes in the electroneutral amino acid exchanger ASCT2

Catherine B. Zander, Thomas Albers, and Christof Grewer

Department of Chemistry, Binghamton University, Binghamton, NY 13902

Neutral amino acid exchange by the alanine serine cysteine transporter (ASCT)2 was reported to be electroneutral and coupled to the cotransport of one Na<sup>+</sup> ion. The cotransported sodium ion carries positive charge. Therefore, it is possible that amino acid exchange is voltage dependent. However, little information is available on the electrical properties of the ASCT2 amino acid transport process. Here, we have used a combination of experimental and computational approaches to determine the details of the amino acid exchange mechanism of ASCT2. The [Na<sup>+</sup>] dependence of ASCT2-associated currents indicates that the Na<sup>+</sup>/amino acid stoichiometry is at least 2:1, with at least one sodium ion binding to the amino acid-free *apo* form of the transporter. When the substrate and two Na<sup>+</sup> ions are bound, the valence of the transport domain is +0.81. Consistently, voltage steps applied to ASCT2 in the fully loaded configuration elicit transient currents that decay on a millisecond time scale. Alanine concentration jumps at the extracellular side of the membrane are followed by inwardly directed transient currents, indicative of translocation of net positive charge during exchange. Molecular dynamics simulations are consistent with these results and point to a sequential binding process in which one or two modulatory Na<sup>+</sup> ions bind with high affinity to the empty transporter, followed by binding of the amino acid substrate and the subsequent binding of a final Na<sup>+</sup> ion. Overall, our results are consistent with voltage-dependent amino acid exchange occurring on a millisecond time scale, the kinetics of which we predict with simulations. Despite some differences, transport mechanism and interaction with Na<sup>+</sup> appear to be highly conserved between ASCT2 and the other members of the solute carrier 1 family, which transport acidic amino acids.

## INTRODUCTION

The amino acid transporter alanine serine cysteine transporter (ASCT)2 transports small, neutral amino acids across the membrane in exchange with intracellular amino acids (Utsunomiya-Tate et al., 1996; Bröer et al., 1999, 2000; Bode, 2001). ASCT2 belongs to the solute carrier 1 (SLC1) family of transporters. The structure of a bacterial glutamate transporter homologue of ASCT2 (glutamate transporter homologue from *Pyrococcus horikoshii* [GltPh]) is known in complex with the transported amino acid substrate (aspartate) and two sodium ions (Yernool et al., 2004; Boudker et al., 2007). Considering the high similarity of the amino acid sequence in the transmembrane domains, it is reasonable to assume that the ASCT2 structure is based on the same fold as that of the glutamate transporters (Albers et al., 2012).

Despite this homology with the glutamate transporters, which have been much more thoroughly studied, the functional properties of the ASCTs are not very well established. For example, the coupling stoichiometries of Na<sup>+</sup> and amino acid in system ASC (transport system that prefers alanine, serine, and cysteine) and ASCT2

are unknown, although models with a 1:1 ratio have been proposed (Bussolati et al., 1992; Bröer et al., 2000; Grewer and Grabsch, 2004). However, mammalian glutamate transporters cotransport three Na<sup>+</sup> ions with one glutamate molecule (Zerangue and Kavanaugh, 1996b), raising the possibility that the stoichiometry of ASCT2 amino acid exchange is also more complex than 1 amino acid: 1 Na<sup>+</sup> ion. Furthermore, amino acid exchange by system ASC and ASCT2 was proposed to be electroneutral (Valdeolmillos et al., 1986; Utsunomiya-Tate et al., 1996), whereas transport by the second ASCT subtype, ASCT1, was proposed to be electrogenic (Zerangue and Kavanaugh, 1996a). Because the amino acid is most likely transported in its zwitterionic, net neutral form, the cotransport of the positively charged sodium ions may lead to at least partial electrogenicity of ASCT2 transport (Bussolati et al., 1992). However, electrogenic reaction steps in ASCTs have not yet been identified.

In this paper, we characterized the mechanism of sodium interaction with ASCT2 and identified the voltage dependence of amino acid exchange. Using a computational approach, based on continuum electrostatic Poisson–Boltzmann (PB) theory, we estimated the net

Correspondence to Christof Grewer: cgrewer@binghamton.edu

Abbreviations used in this paper: APBS, adaptive Poisson–Boltzmann solver; ASCT, alanine serine cysteine transporter; EAAT, excitatory amino acid transporter; GltPh, glutamate transporter homologue from *Pyrococcus horikoshii*; MD, molecular dynamics; MNI, methoxynitroindoline; PB, Poisson–Boltzmann; RL2, reentrant loop 2; SLC1, solute carrier 1.

© 2013 Zander et al. This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

charge translocation of ASCT2 with various numbers of cations bound, predicting that the main translocation reaction is associated with the movement of positive charge across the membrane. Consistent with this prediction, charge movement was observed experimentally by applying step changes in the membrane potential, or alanine concentration jumps to ASCT2. In addition, we obtained experimental evidence that at least two Na<sup>+</sup> ions are cotransported with amino acid. Results from molecular dynamics (MD) simulations also suggest the existence of at least two, possibly three, Na<sup>+</sup>-binding sites. Our results are explained with a sequential transport mechanism in which one Na<sup>+</sup> binds to the empty transporter, followed by binding of the amino acid and the subsequent binding of another Na<sup>+</sup> ion, followed by electrogenic amino acid translocation across the cell membrane.

## MATERIALS AND METHODS

### Cell culture and transfection

cDNA coding for rat ASCT2 was provided by S. Bröer (The Australian National University, Acton, Australia; Bröer et al., 1999, 2000). The coding region of the cDNA was subcloned into the EcoRI site of the pBK-CMV vector (Agilent Technologies). The cDNA construct was used to transiently transfect human embryonic kidney (HEK) cells (HEK293; American Type Culture Collection no. CGL 1573) with reagent (Fugene HD; Roche), according to the instructions of the supplier. Cells were transfected in the exponential growth phase. Electrophysiological experiments were done 24 h after transfection.

### Electrophysiological methods

ASCT2-mediated currents were recorded with an amplifier (Adams and List EPC7; HEKA) under voltage-clamp conditions. The whole-cell current-recording configuration was used. Electrode resistances were 2–3 M $\Omega$ , and the series resistance ( $R_s$ ) was 5–8 M $\Omega$ .  $R_s$  was not compensated in the steady-state recordings, as compensation had no effect on the magnitude of the observed currents. However, up to 60%  $R_s$  compensation was used in the voltage-jump experiments to accelerate the time needed for charging of the membrane. We have previously estimated the instrument time constant under these conditions as 200  $\mu$ s (Mim et al., 2007). The extracellular bath solution contained (mM): 140 NaMES or NaSCN, 2 Mg gluconate<sub>2</sub>, 2 Ca gluconate<sub>2</sub>, and 10 HEPES (MES, pH 7.4/NaOH). The pipette solution contained (mM): 130 NaMES or NaSCN, 2 MgCl<sub>2</sub>, 10 EGTA, 10 HEPES, and 10 liter-alanine/cysteine, pH 7.3/NaOH. Using this intracellular solution, the transporters operate in the exchange mode, in which external amino acid is exchanged with internal amino acid in the absence of net transport. The exchange mode is associated with the activation of an uncoupled anion current, which was used to determine the [Na<sup>+</sup>] dependence. For experiments in the presence of intracellular K<sup>+</sup>, Na<sup>+</sup> in the above solutions was replaced with K<sup>+</sup>. For [Na<sup>+</sup>]-dependent experiments, Na<sup>+</sup> was replaced with *N*-methylglucamine<sup>+</sup> (NMG<sup>+</sup>). The currents were low-pass filtered at 3 kHz (EPC7 built-in filter) and digitized with a digitizer board (Digidata 1200; Axon Instruments) at a sampling rate of 10–50 kHz (controlled by Axon PClamp7 software). All experiments were performed at room temperature.

Rapid solution exchange was essentially performed as described previously (Watzke et al., 2001). In brief, substrates and inhibitors were applied to the ASCT2-expressing HEK293 cells

with a quartz tube (350- $\mu$ m tube diameter) positioned at a distance of  $\approx$ 0.5 mm to the cell. The linear flow rate of the solutions emerging from the opening of the tube was  $\sim$ 5–10 cm/s, resulting in typical rise times of the whole-cell current of 30–50 ms (10–90%).

### Laser-pulse photolysis

Laser-pulse photolysis experiments were performed as described previously (Zhang et al., 2007). In brief, the cells were equilibrated with neutral amino acids or methoxynitroindoline (MNI)-caged alanine using a fast solution exchange device. Photolysis was initiated with a light flash generated by a frequency-tripled Nd-Yag laser (Minilite II; Continuum). The laser light was delivered to the cell with an optical fiber (350- $\mu$ m diameter), and its energy was adjusted with neutral density filters (Andover Corporation). Typical laser energies, measured at the emitting end of the optical fiber using an energy meter (Gentec), were in the range of 50 to 400 mJ/cm<sup>2</sup>. A standard alanine concentration of 1 mM was applied to the cell by rapid solution exchange before and after laser-pulse photolysis to estimate the concentration of photolytically released alanine and to test the cell for damage by the laser pulse.

### Data analysis

Nonlinear regression fits of experimental data were performed with Origin (OriginLab) or Clampfit (pClamp8 software; Axon Instruments). Dose–response relationships of currents were fitted with a Michaelis–Menten-like equation, yielding  $K_m$  and  $I_{max}$ . Each experiment was performed in triplicate with at least two different cells. The error bars represent mean  $\pm$  SD, unless stated otherwise.

### PB calculations using adaptive PB solver (APBS)mem

Calculations of electrostatic energies were essentially performed as described previously (Grewer et al., 2012). The parameters for the APBSmem routine (Callenberg et al., 2010) were as follows. The temperature was set to 298 K. The symmetrical ion concentrations were set to 0.1 M. The membrane thickness was 30  $\text{Å}$ , with an upper and lower exclusion radius of 31 and 23  $\text{Å}$ , respectively. The dielectric constants were 80 for the intracellular and extracellular water and within the extracellular bowl created within the exclusion radius, 2.0 for the membrane, and 2–10 for the protein. The lipid headgroup region of the bilayer was not modeled. For the PB calculations, the finest focusing step used 161 grid points, in a 130- $\text{Å}$  grid, in x, y, and z directions. Three focusing steps were used, and with these parameters the energy values converged. The C and N termini were charged. Neutralization of the termini only has a minor effect on the results. To assess the effect of the z-position within the membrane, the transporter was moved up and down by  $\pm$ 2  $\text{Å}$  relative to the membrane. This caused a reduction of the calculated valence by 21–25% but in no case changed the sign of the valence.

### MD simulation protocol

The simulated system consists of the trimeric ASCT2 transporter, modeled on the structure of the GltPh transporter (Protein Data Bank accession nos. 2NWX [Boudker et al., 2007] and 3KBC [Reyes et al., 2009]; sequence alignment shown in Fig. S1), set up in a patch of POPE membrane in its box of water using visual MD (Humphrey et al., 1996). The overall system (neutralized by adding counterions) is composed of  $\sim$ 13,3000 atoms (illustrated in Fig. S2). The Modeller algorithm was used to build the homology model (Eswar et al., 2003). The C- and N-terminal domains of ASCT2 are not present in the GltPh crystal structures. Therefore, the ASCT2 sequence was truncated before V56 (N terminus) and after T487 (C terminus). The simulation uses the CHARMM27 force field with the CMAP corrections and the TIP3P model for explicit water. MD calculations were performed with periodic boundary conditions with a time step of 2 fs using NAMD 2.8

(Phillips et al., 2005). First, lipid tails were melted while all other atoms were held fixed for 400 ps at constant volume and constant temperature of 300 K. Then, the system was equilibrated at constant temperature and constant pressure of 1 bar for 500 ps while the atoms of the protein were restrained by harmonic potentials ( $k = 1 \text{ kcal/mol/\AA}^2$ ) to allow for packing of the lipid molecules around the protein. Next, the simulation was continued for 1 ns without constraints.

Production runs of up to 12 ns were performed after 8 ns of equilibration at constant area of the lipid bilayer and at constant pressure of 1 bar. Constant temperature was maintained by using Langevin dynamics with a damping coefficient of 5 ps. The Langevin piston method was used to maintain a constant pressure of 1.0 atm with a piston period of 200 fs. Short-range nonbonded interactions were calculated using a cutoff distance of 12 Å, and long-range electrostatic interactions were calculated using the particle mesh Ewald method (Darden et al., 1993).

#### Online supplemental material

The online supplemental material contains tables showing data from PB calculations (Tables S1 and S2), the sequence alignment between ASCT2 and GltPh for homology modeling (Fig. S1), an illustration of the MD setup (Fig. S2), and additional results from MD simulations (Fig. S3). The online supplemental material is available at <http://www.jgp.org/cgi/content/full/jgp.201210948/DC1>.

## RESULTS


### One or two sodium ions bind to the apo form of the transporter

Previous results showed that  $\text{Na}^+$  can activate a leak anion conductance in the amino acid-free apo form of ASCT2 (Greuer and Grabsch, 2004). In agreement with these results, anion current induced by  $\text{Na}^+$  in the absence of amino acid was activated at low  $[\text{Na}^+]$  (Fig. 1), with an apparent  $K_m$  of  $0.25 \pm 0.06 \text{ mM}$ . High  $[\text{Na}^+]$ , however, led to a small but significant reduction of the anion current ( $K_m$  estimated as 51 mM, although an exact value cannot be determined because of lack of saturation at 140 mM  $\text{Na}^+$ ; Fig. 1). This biphasic behavior can be explained by the binding of at least one but possibly two  $\text{Na}^+$  ions to the apo form of the transporter (see fit to two-site model in Fig. 1). In this interpretation,  $[\text{Na}^+]$  below 5 mM results in occupation of only the first high affinity binding site with an apparent  $K_m$  of 0.25 mM and activation of the leak anion conductance. The intrinsic dissociation constant of this binding step will be termed " $K_{\text{Na1}}$ " throughout this paper. At  $[\text{Na}^+]$  of 140 mM and higher, the low affinity binding site with an apparent  $K_m$  in the 50-mM range is also occupied, resulting in a state with two  $\text{Na}^+$  ions bound. This state has slightly lower anion conductance than the singly occupied state, causing the reduction of the leak anion current at high  $[\text{Na}^+]$  in Fig. 1.

### ASCT2 harbors at least two $\text{Na}^+$ -binding sites

It has been hypothesized that amino acid transport by ASCT2 is coupled to the exchange of one  $\text{Na}^+$  ion (Bussolati

et al., 1992; Bröer et al., 2000; Greuer and Grabsch, 2004). However, the highly homologous bacterial aspartate transporter GltPh has two identified  $\text{Na}^+$ -binding sites (cotransporting three  $\text{Na}^+$  ions; Boudker et al., 2007; Groeneveld and Slotboom, 2010), and mammalian glutamate transporters may have three  $\text{Na}^+$  sites (Zerangue and Kavanaugh, 1996b). To test for potential multiple  $\text{Na}^+$ -binding sites, we determined the  $\text{Na}^+$  dependence of ASCT2 function. This was done using the well-established anion conductance of ASCT2 as a readout (Greuer and Grabsch, 2004), as this conductance can be activated in the exchange mode at steady state. Fig. 2 A shows anion currents at various extracellular  $[\text{Na}^+]$ . The current was inwardly directed at high  $[\text{Na}^+]$ , as expected from an alanine-induced activation of the anion conductance ( $\text{SCN}^-$  outflow). At  $[\text{Na}^+]$  below 2 mM, the leak anion current was reduced after the application of alanine (Fig. 2 A). This apparent outward current was induced by alanine inhibition of the tonic leak anion conductance, resulting in inhibition of  $\text{SCN}^-$  outflow (see Greuer and Grabsch, 2004). This current- $[\text{Na}^+]$  relationship is quantified in Fig. 2 B. The outward current at low  $[\text{Na}^+]$  indicates that alanine is an inhibitor of the leak anion conductance under these conditions, as previously found for competitive ASCT2 inhibitors (Albers et al., 2012). This biphasic  $[\text{Na}^+]$  dependence of the anion current is incompatible with a single- $\text{Na}^+$  site model but clearly requires the existence of at least two  $\text{Na}^+$ -binding sites. Furthermore, our data show that low sodium concentration cannot be overcome by saturating [alanine], directly demonstrating that at least one  $\text{Na}^+$  ion must bind to the alanine-bound form of the transporter.




**Figure 1.** At least one but possibly two  $\text{Na}^+$  ions bind to the apo form of ASCT2.  $[\text{Na}^+]$  dependence of the leak anion current (normalized to the response at 140 mM  $\text{Na}^+$ ). The solid line represents a fit to the equation  $I = A [\text{Na}^+]/(K_1 + [\text{Na}^+]) + B [\text{Na}^+]/(K_2 + [\text{Na}^+])$ . Here,  $A$  and  $B$  are scaling parameters, and the values for  $K_1$  and  $K_2$  were  $0.25 \pm 0.06 \text{ mM}$  and  $51 \pm 10 \text{ mM}$ , respectively. The intracellular solution contained 120 mM  $\text{NaSCN}$  and 10 mM alanine. The extracellular solution contained 140 mM  $\text{NaMes}$  ( $V = 0 \text{ mV}$ ).






Grabsch, 2004; unpublished data). Furthermore, no amino acid-sensitive currents were detected in non-transfected control cells.



**Figure 3.** Amino acid exchange by ASCT2 is associated with transmembrane charge movement. (A) Voltage-jump protocol used in the experiment. (B) Original current traces in response to the voltage jump in the absence (red) and presence (black) of 1 mM cysteine (inset shows original currents not cut off at 0.5 nA). (C) Current trace obtained by subtraction of currents in the presence from that in the absence of cysteine. (D) Voltage dependence of the charge movement of the on-response obtained by integrating the current over time. The red line illustrates a fit of the Boltzmann equation ( $Q(V_m) = Q_{\min} + Q_{\max} \cdot [I + \exp(Z_Q(V_Q - V_m) \cdot F/(RT))]^{-1}$ ;  $Q$  represents the charge moved,  $Z_Q$  is the valence of this charge,  $V_Q$  is the midpoint potential of the charge movement,  $R$  is the gas constant,  $T$  is the temperature, and  $F$  is the Faraday constant) to the data with a  $z_q$  of 0.74 and a midpoint potential of  $-19$  mV. (E) Voltage dependence of the relaxation rate constant for decay of the transient current. The line represents a fit to the equation  $1/\tau = k_1(0) \exp(z_q FV/2RT) + k_2(0) \exp(-z_q FV/2RT)$ , with  $z_q = 0.41$ .

#### Continuum electrostatic models predict electrogenic nature of ASCT2 conformational transitions

To rationalize the results from the voltage-jump experiments, we used a computational approach to predict valences associated with the ASCT2 exchange reaction. The approach is based on solving the linearized PB equation for different structural states of the transporter in the presence of an internal biasing potential. The APBS was used for the calculations (Baker et al., 2001). If two states of a membrane protein are connected by an electrogenic transition, one state should be stabilized by the transmembrane potential relative to the other in terms of its total electrostatic energy (Callenberg et al., 2010). This approach has been successfully applied in the past to voltage-gated ion channels (Grabe et al., 2004; Silva et al., 2009; Choudhary et al., 2010). The setup of the system, using an implicit membrane with dielectric constant  $\epsilon = 2$ , is shown in Fig. 4 A. The transport proteins, which are homology models of ASCT2 based on the GltPh structures (Protein Data Bank



**Figure 4.** Valence associated with transitions of several forms of the ASCT2 transport domain. (A) Setup used for the APBS calculations. Blue color indicates inside positive or negative membrane potential (+ or  $-100$  mV), and red color indicates 0-mV membrane potential. (B) Illustration of the conformational change of an individual transport domain from outward- to inward-facing. (C) Relative change in the total electrostatic energy for the outward-facing to inward-facing transition of an individual subunit for the *apo* transporter (red), and the forms with 2  $\text{Na}^+$  bound (black) and 2  $\text{Na}^+$  and alanine bound ( $\Delta E$  was set to 0  $k_B T$  at 0 mV). (D) Valences calculated from the slopes of C.

accession nos. 2NWX [Boudker et al., 2007] and 3KBC [Reyes et al., 2009]), were inserted into the membrane using the APBSmem routine (Callenberg et al., 2010).

We first determined the valence of what is thought to be the major conformational transition from outward- to inward-facing configuration (Fig. 4, A and B; Crisman et al., 2009; Reyes et al., 2009). As illustrated in Fig. 4 (C and D), inside negative potential slightly stabilized the outward-facing configuration in the empty transporter (valence,  $z = -0.35$ ), suggesting that in the absence of bound cations, the transporter carries a net negative charge (assuming  $pK_a$  values for the buried charges to be similar to their  $pK_a$  values in water). The addition of cations to their presumed binding sites (modeled in analogy to the Na1 and Na2 sites of GltPh; see below) resulted in a positive valence of this transition (Fig. 4, B and C). As expected, the overall neutral zwitterionic amino acid substrate (alanine) contributed only minimally to the valence (Fig. 4, C and D). Glutamate transporters have a third  $Na^+$ -binding site, which was proposed to be formed by conserved amino acid side chains deeply buried within the transmembrane domain (for example, D367 and T101; Tao et al., 2006, 2010; Huang and Tajkhorshid, 2010). These amino acid residues are conserved in ASCT2. Thus, it is possible that ASCT2 harbors a third  $Na^+$ -binding site in this position. After adding a third  $Na^+$  ion to this site, the valence of the loaded transporter was calculated as  $z = +1.18$  (Fig. 4 D). The results of the PB calculations are summarized in [Table S1](#). Fig. 4 D also shows results from control PB calculations in which all charges but seven conserved ones in the transmembrane domain were removed to test for contribution of the movement of charges in the soluble domain of the transporter to the valence. Although the valence was slightly reduced after removal of these charges, the sign of the valence was unchanged (Fig. 4 D). Collectively, our continuum electrostatic analysis suggests that the transition from the outward- to the inward-facing state is associated with positive charge movement in the amino acid/ $Na^+$ -loaded transporter and is, thus, electrogenic.

Ionizable amino acid residues deeply buried in the transmembrane domain may have perturbed  $pK_a$  values, as recently suggested for the aspartate side chain coordinating sodium in the Na1 site of glutamate transporters (Mwaura et al., 2012). To test this hypothesis, we estimated  $pK_a$  values of relevant side chains using the empirical PROPKA approach (Rostkowski et al., 2011). The estimated values suggest that at least D473, which is expected to contribute to  $Na^+$  binding, has a strongly elevated  $pK_a$  (10.3 outward-facing, 10.9 inward facing) in the absence of ions. Assuming that the D473 side chain is protonated, the valence of the conformational transition of the *apo* form of ASCT2 is +0.05 and that of the fully loaded form (3  $Na^+$  and Ala) is +1.57 (Fig. 4 D).

### Rapid application of amino acid induces transient currents caused by relaxation of the electrogenic exchange equilibrium

The exchange equilibrium can be perturbed by voltage jumps as well as by rapid application of amino acid. In this single-turnover experiment, the transporter-binding sites are initially distributed according to the equilibrium constant associated with the inward-outward-facing transition in the absence of amino acid. When amino acid is rapidly applied to the extracellular side, binding to ASCT2 initiates transition to the inward-facing configuration. A typical current recording is shown in Fig. 5 B in the presence of intracellular  $K^+$ , a cation that is not thought to interact with ASCT2. Rapid alanine application through photolysis of caged alanine (Zhang et al., 2007) within less than a microsecond (Fig. 5 A; released [alanine],  $\sim 600 \mu M$ ) resulted in a small transient inward current that rapidly relaxed back to a steady state within  $\sim 10$  ms (Fig. 5 B). The inward direction of the current proves that positive charge is moving into the cell during the exchange equilibration process. Control experiments with nontransfected cells demonstrate the absence of current signals upon application of the laser flash. Further control experiments are shown in Zhang et al. (2007).

The observed time constant associated with the current decay,  $\tau_{obs} = 1/k_{obs}$ , can be analyzed using the simplified kinetic scheme (Scheme 1) shown in the Appendix. According to this scheme, the equation for  $k_{obs}$  is:

$$k_{obs} = k_1 + k_{-1}. \quad (1)$$

In the presence of intracellular  $K^+$ ,  $\tau_{obs}$  was  $3.1 \pm 0.5$  ms. As expected, no steady-state current was observed because ASCT2 is incapable of catalyzing steady-state forward transport. According to Eq. 1,  $k_{obs} = k_1$  under these conditions because the reverse reaction ( $T_{in}$  to  $T_{out}$ ) cannot be initiated in the absence of internal  $Na^+$  and amino acid ( $k_{-1} = 0$ ).


To further analyze the exchange reaction, we repeated the experiment under exchange conditions (140 mM  $Na^+$  and 10 mM alanine inside, rather than  $K^+$ ). Here, binding sites initially face outward (state  $T_{out}$  in Scheme 1; see Appendix), caused by the application of high amino acid and  $Na^+$  concentrations to the intracellular side in the absence of extracellular amino acid (Fig. 5 C, inset). External alanine application initiates  $Na^+$ /alanine exchange (establishing a new equilibrium between  $T_{out}$  and the inward-facing configuration,  $T_{in}$ ; Scheme 1; see appendix). A significantly larger transient current was observed than in the presence of internal  $K^+$ , which decayed with a slightly faster time constant of  $1.7 \pm 0.4$  ms (Fig. 5 C). The larger current amplitude in the exchange mode suggests that internal  $Na^+$ /alanine is more efficient in driving the binding sites to an outward-facing configuration than intracellular  $K^+$ . The faster time constant



indicates that the relaxation rate is now determined by the sum of the rate constants for the bidirectional forward and backward translocation reactions ( $k_{\text{obs}} = k_1 + k_{-1}$ ), whereas in the presence of intracellular  $\text{K}^+$ , only the unidirectional rate constant for forward translocation is observed ( $k_{\text{obs}} = k_{-1}$ ). With  $k_1 = 330 \text{ s}^{-1}$ , we can estimate  $k_{-1} = 600 \text{ s}^{-1} - 330 \text{ s}^{-1} = 270 \text{ s}^{-1}$ . Therefore, these data suggest that translocation occurs with rates of similar order of magnitude in both directions but

with a slight favoring of the inward-facing configuration at 0 mV.

We also determined the voltage dependence of the transient currents. As shown in Fig. 5 (D and E), the charge movement saturates at very negative transmembrane potentials, but it decreases at positive potentials. This indicates that at  $V < -60 \text{ mV}$ , the translocation equilibrium is fully driven toward the inward-facing configuration. The half-maximal potential of about +40 mV confirms the prediction that at 0 mV, the exchange equilibrium is shifted toward the inward-facing configuration.



**Figure 5.** Amino acid exchange is associated with inward charge movement. (A) Mechanism of photolysis of MNI-caged alanine. (B and C) Current in response to liberation of  $\sim 200 \mu\text{M}$  of free alanine from 3 mM of MNI-caged alanine at  $t = 0$  in the presence of internal  $\text{K}^+$  (B) or  $\text{Na}^+$ /alanine (C). The anion was  $\text{Mes}^-$  in all cases ( $V = 0 \text{ mV}$ ). (D) Current response to alanine application at  $t = 0$  at membrane potentials of  $-60 \text{ mV}$  (red),  $0 \text{ mV}$  (black), and  $+60 \text{ mV}$  (blue). (E) Charge movement as a function of the transmembrane potential. The charge was obtained by integrating the transient current response over time. The red line shows a fit to the Boltzmann equation (see legend to Fig. 3;  $z_Q = 0.58$  and  $V_Q = 41 \text{ mV}$ ).


Amino acid interaction with ASCT2 is electroneutral, but  $\text{Na}^+$  interaction is electrogenic

For several  $\text{Na}^+$ -driven transporters, it was suggested that  $\text{Na}^+$  binding and/or conformational changes associated with it contribute to transporter electrogenicity (Loo et al., 1993; Wadiche et al., 1995; Lu and Hilgemann, 1999; Watzke et al., 2001; Forster et al., 2002). We tested this possibility by measuring current responses to voltage jumps in the absence of extracellular amino acid, restricting ASCT2 to reaction steps associated with  $\text{Na}^+$  binding. As shown in Fig. 6 A (left), transient, inhibitor-sensitive currents were observed upon stepping the membrane potential from 0 to  $-60$  or  $+60 \text{ mV}$  when the extracellular  $[\text{Na}^+]$  was  $0.5 \text{ mM}$ . A concentration of  $0.5 \text{ mM}$  was chosen because it is close to the  $K_m$  of the empty transporter for  $\text{Na}^+$  ( $K_{\text{Na1}}$ ), which was in the  $0.05$ - and  $0.5$ -mM range, dependent on the various methods used to determine this value. At  $[\text{Na}^+] = K_m$ , the maximum charge movement should be observed. In contrast, at saturating  $[\text{Na}^+]$ , no or little charge movement should be observed, because the jump to more negative voltage cannot drive additional  $\text{Na}^+$  into the already occupied binding site. Consistent with this expectation, no currents were observed at  $140 \text{ mM}$  of external  $\text{Na}^+$ , when the  $\text{Na}^+$ -binding site(s) is fully saturated (Fig. 6 A, right). The charge movement was capacitive in nature, with the charge of the off-response being of equal magnitude but of opposite sign of that of the on-response (Fig. 6 A). The voltage dependence of the charge movement could be represented with the Boltzmann equation, yielding an apparent valence,  $z_Q$ , of  $0.41 \pm 0.1$  (Fig. 6 B). Collectively, these results indicate that  $\text{Na}^+$  binding, or a conformational change induced by this binding, is electrogenic. The results also confirm the existence of a high affinity  $\text{Na}^+$ -binding site with an apparent affinity in the submillimolar range.

We next tested whether the PB computations predict voltage dependence of  $\text{Na}^+$  binding. The models were based on the movement of a  $\text{Na}^+$  ion from the water phase ( $\epsilon = 80$ ) to one of the three predicted, occluded  $\text{Na}^+$ -binding sites, respectively. The calculated valences are shown in Fig. 6 D. These valences are positive (Table S2), as expected for movement of a positively charged ion into the low dielectric environment of the protein. These

computational results support our experimental results on electrogenic  $\text{Na}^+$  association. The actual  $\text{Na}^+$  association process may be separated into two steps. First,  $\text{Na}^+$  may move into its binding site through an aqueous access channel in a voltage-independent fashion. Second, occlusion of the bound  $\text{Na}^+$  results in charge displacement. This possibility is supported by MD simulations shown in Fig. S3, demonstrating that the Na1 site becomes hydrated from the extracellular side in the absence of any bound  $\text{Na}^+$ . Therefore, it is likely that the main electrogenic step is the occlusion of the  $\text{Na}^+$  in its binding site after initial electroneutral binding through an aqueous pathway.

The amino acid substrate presumably binds to ASCT2 in its zwitterionic form. Although the overall charge of the substrate is zero, the positive and negative charges on the amino and carboxy group may lead to transmembrane charge movement. However, the  $K_m$  of ASCT2 for alanine was independent of the transmembrane potential (Fig. 6 C). Furthermore, PB calculations indicate that alanine binding is associated with little valence ( $z = 0.04$ ;



**Figure 6.**  $\text{Na}^+$  binding to the empty transporter is electrogenic, but alanine binding is electroneutral. (A) Currents in response to voltage jumps (top panel indicates voltage protocol) at 0.5 mM of extracellular  $\text{Na}^+$  (left) and 140 mM  $\text{Na}^+$  (right) (counterion was  $\text{Mes}^-$ ). The intracellular solution contained 120 mM  $\text{NaMes}$ /10 mM alanine. (B) Voltage dependence of the charge movement of the on-response (solid line represent fit to a Boltzmann equation). (C) Voltage dependence of the  $K_m$  of ASCT2 for alanine. The intracellular solution contained 120 mM  $\text{NaSCN}$ /10 mM alanine, and the extracellular solution contained 140 mM  $\text{NaMes}$ . (D) Predicted valences computed by PB analysis for alanine binding as well as for the binding of  $\text{Na}^+$  to three hypothesized binding sites. The inset shows an illustration of the alanine-binding reaction used for calculation of the valence.

Fig. 6 D; see illustration of setup in the inset). Therefore, it is likely that the amino acid association reaction is electroneutral.

#### Evidence for conserved $\text{Na}^+$ sites from MD simulations

A high resolution crystal structure is available for the aspartate transporter GltPh (Boudker et al., 2007), which is highly homologous to ASCT2. Although the overall ASCT2 sequence shares only 30% identity and 50% similarity with GltPh, the critical C-terminal transport domain is much better conserved with 45% identity and 66% similarity. The amino acid residues within 4 Å of the ligand-binding site are even better conserved, with 69% identity and 85% similarity. Therefore, we hypothesize that an ASCT2 homology model based on the GltPh structure will provide a reasonable starting point for MD simulations, at least for the transmembrane domain, while keeping in mind the caveats associated with MD simulations of low resolution or homology models.

We first studied the transporter with bound amino acid (alanine) and with two  $\text{Na}^+$  ions in the sites that are analogous to the Na1 and Na2 sites of GltPh (see Fig. S2 for an illustration of the simulation setup). After 8 ns of simulation, the alanine molecule, as well as the two  $\text{Na}^+$  ions, remained in their binding sites (Fig. 7, A and B). Reentrant loop 2 (RL2) adapted to the lower molecular volume of the bound alanine compared with aspartate in GltPh, resulting in a closing in of the alanine molecule and exclusion of water in a lid-like fashion. These results suggest that the function of RL2 as the external gate that protects the bound amino acid from the water phase is conserved in ASCT2. The  $\text{Na}^+$  ion in the Na2 site is coordinated by a total of five main chain carbonyl oxygens from the SVGA motif at the tip of RL2 (SIGT in GltPh), as well as from the TVN motif in TM7 (TIN in GltPh), which show 86% similarity between ASCT2 and GltPh (Fig. 7 B). The Na1 site is formed by the two carboxylate side chain oxygens of D473, as well as the main chain and side chain oxygens of N384 and two other main chain carbonyls from TM8 (Fig. 7 A).


Both  $\text{Na}^+$  ions were stable in their predicted binding sites in the 2  $\text{Na}^+$ /alanine-loaded transporter within the 8 ns of simulation, as shown for Na2 in Fig. 7 C (red trace;  $n = 3$ ). The sodium ion in the Na1 site also remained stable in the absence of occupation of the Na2 site, as well as when both amino acid and  $\text{Na}^+$  in the Na2 site were absent (Fig. 7 D). Stability is indicated by a lack of significant variability of the D473(Oδ)-Na1 distance with time. This result indicates that occupation of the Na1 site can occur in the empty transporter, i.e., in the absence of bound amino acid substrate. In contrast, a sodium ion in the Na2 site was only stable when the amino acid was bound (Fig. 7 C). In the absence of alanine, the cation dissociated from the Na2 site within 8 ns in two out of three simulation runs (representative traces shown in Fig. 7 C; black line shows dissociating cation).



This result suggests that the cation in the Na2 site is only stable in the presence of bound amino acid.

In some MD simulations, the sodium ion in the Na1 site became coordinated by an oxygen atom in the carboxylate side chain of D386. This configuration is analogous to an intermediate cation site proposed by Huang and Tajkhorshid (2010) for GltPh. This finding raises the possibility that a third Na<sup>+</sup>-binding site exists, in which the Na<sup>+</sup> is coordinated by the D386 side chain as well as by the T139 side chain oxygen from TM3. Evidence for the existence of such a site has been obtained by previous mutagenesis studies showing that neutralization of D386 resulted in a dramatic reduction of Na<sup>+</sup>

affinity of ASCT2 (Tao et al., 2006), as well as MD simulations of glutamate transporters (Huang and Tajkhorshid, 2010; Heinzelmann et al., 2011; Bastug et al., 2012). Therefore, we placed a third sodium ion in the proposed Na3 site and performed MD simulations. The third Na<sup>+</sup> ion in the site analogous to the GltPh Na3 site was stable within 8 ns of simulation (Fig. 7, E and F). These results indicate the potential for a third Na<sup>+</sup>-binding site on ASCT2. It should be noted that Larsson et al. (2010) have proposed an alternative Na3 site in the related glutamate transporters. Because there have been no mutational studies on this site for ASCT2, we have not analyzed it further.



**Figure 7.** Sodium-binding sites are conserved in ASCT2. (A and B) Na1 (A)- and Na2 (B)-binding sites for Na<sup>+</sup> after 8 ns of MD simulation. (C) Representative traces for the distance between the T382  $\alpha$  carbon and the Na<sup>+</sup> ion in the Na2 site as a function of time for the alanine- and Na1-bound transporter (red), and in the absence of alanine (black and green). (D) Time evolution of the distance between D473(O $\delta$ 2) and the cation in the Na1 site in the absence of occupied Na2 (black), as well as in the absence of both Na2 and alanine (red). (E) Distances calculated from an 8-ns MD simulation between D386(O $\delta$ 2) and Na3. The colors indicate distances calculated from the same simulation in the three different subunits (black, chain A; red, chain B; green, chain C). (F) Predicted Na3-binding site at the end of a 10-ns simulation.

## DISCUSSION

In this paper, we have used a combined experimental and computational approach to determine the mechanism and electrogenicity of Na<sup>+</sup> interaction with and translocation by the neutral amino acid exchanger ASCT2. The two major conclusions from these studies are the following: (1) Despite being an electroneutral exchanger, Na<sup>+</sup>-coupled amino acid translocation by ASCT2 is electrogenic and, thus, sensitive to the transmembrane potential. (2) In contrast to previous suggestions, amino acid exchange is coupled to the translocation of at least two but possibly three Na<sup>+</sup> ions. Cation and amino acid binding to ASCT2 is sequential, with at least one Na<sup>+</sup> binding to the *apo*, empty form of the transporter, followed by amino acid binding, followed by binding of another Na<sup>+</sup> to the substrate-bound form (Fig. 4 E).

Overall, the mechanism of amino acid transport by ASCT2 is surprisingly similar to the mechanism of glutamate transport by the related glutamate transporters from the SLC1 family (Wadiche and Kavanaugh, 1998; Watzke et al., 2001; Bergles et al., 2002). Like in the glutamate transporters, amino acid transport requires the cotransport of two to three sodium ions. The sequence of sodium and amino acid binding also appears to be conserved between the ASCT and excitatory amino acid transporter (EAAT) members of the family, as are the major amino acids that contribute oxygen ligands to the cation-binding sites. Finally, Na<sup>+</sup> binding to the empty transporter, or conformational changes associated with it, is associated with charge movement in ASCT2. Similar findings, based on voltage-jump analysis, were reported for several subtypes of EAATs (Wadiche et al., 1995; Watzke et al., 2001).

Despite these similarities, there are also significant differences between the neutral and acidic amino acid transporters of the SLC1 family. First, Na<sup>+</sup> binds to the empty ASCT2 with very high apparent affinity in the submillimolar range (this work and Grewer and Grabsch, 2004). In contrast, the affinity of the *apo* form of EAATs for Na<sup>+</sup> is low, with an apparent affinity of 100 mM for


excitatory amino acid carrier 1 (EAAC1; Tao et al., 2006). The high affinity is important for ASCT2's physiological function as an amino acid exchanger, because the tightly bound sodium ion most likely never dissociates from the transporter (Bröer et al., 2000). Although the intracellular affinity for  $\text{Na}^+$  is not known, it is most likely high, such that  $\text{Na}^+$  has a modulatory function rather than providing a driving force for substrate transport. Second, ASCT2, in contrast to the EAATs, cannot complete a full transport cycle because intracellular  $\text{K}^+$  is unable to drive relocation of the empty transporter, as also noted in Bröer et al. (2000). Clearly, no steady-state component of the transport current was observed in the presence of intracellular  $\text{K}^+$  (Fig. 3 A), indicating that net forward transport of amino acid is not possible, or is at least slow. Furthermore, internal  $\text{Na}^+$ /amino acid are much more capable of rearranging the binding sites to outward-facing than  $\text{K}^+$  (Fig. 3, A and B). Collectively, these results suggest that  $\text{K}^+$  is not able to interact with ASCT2. The preference of  $\text{Na}^+$  over  $\text{K}^+$  may be caused by the high  $\text{Na}^+$  affinity of ASCT2, preventing intracellular  $\text{K}^+$  from displacing  $\text{Na}^+$ , despite its  $\sim 30$ -fold higher intracellular concentration. It is noteworthy that the potentially negatively charged glutamate residue, E373 in EAAC1, is substituted by a glutamine in ASCT2 (Q392). E373 has been implicated in potassium interaction as well as in protonation of the glutamate transporter (Pines et al., 1995; Grewer et al., 2003). A transporter with the E373Q mutation is defective in potassium-induced relocation and is pH insensitive. ASCT2 is also pH insensitive (Grewer et al., 2003), suggesting that the glutamine side chain can mimic a protonated glutamic acid residue. If this hypothesis is correct, electrostatic charge compensation in position in ASCT2 is unchanged compared with the protonated glutamate transporter.

Our results from the concentration-jump analysis directly show that inward amino acid translocation by ASCT2 is coupled to the inward movement of positive charge. These results agree well with the valence calculations from solving the linearized PB equation, predicting a positive valence for the fully loaded transporter. Because the bound amino acid is zwitterionic, but net neutral, the translocated charge could be generated by positive charges of the bound  $\text{Na}^+$  ions, by positively

charged ion/substrate-binding sites, or by both of these effects. However, according to the PB calculations, the empty, unloaded transporter either carries net negative charge in the binding sites (negative valence) or is neutral, providing evidence that the observed inward charge movement is caused by inward movement of the cotransported  $\text{Na}^+$  ions within the membrane electric field.

In a previous report by Bussolati et al. (1992), L-threonine transport by natively expressed system ASC was found to be weakly voltage dependent, with a bell-shaped dependence of flux on voltage. If influx of amino acid takes place only by exchange, a bell-shaped voltage dependence of uptake is expected, as either the forward or reverse rate becomes rate limiting at extreme membrane potentials (see Appendix for a simplified mathematical treatment). Similar considerations have been put forward for electroneutral but voltage-dependent  $\text{Na}^+$  exchange by the  $\text{Na}/\text{K}$  ATPase (De Weer et al., 1988). The slight favoring of inward transport over outward transport of alanine at 0 mV is expected to result in a slight left shift of the maximum of the flux–voltage relationship to positive potentials (see Fig. A1 in Appendix).

Our new results allow us to extend previous mechanistic models proposed for amino acid exchange by ASCTs and system ASC (Bussolati et al., 1992; Bröer et al., 2000). The extended model (Fig. 8) includes a  $\text{Na}^+$ /amino acid stoichiometry of at least 2:1, with sequential binding of  $\text{Na}^+$  and amino acid. The first  $\text{Na}^+$  ion binds to the *apo* state of the transporter in a voltage-dependent manner, or it elicits a voltage-dependent conformational change. The second possibility is more likely, because the MD simulations predict that  $\text{Na}^+$  accesses its  $\text{Na}1$  site through a water-filled channel, which upon binding closes while excluding water molecules from the binding site. Therefore, the overall cation-binding reaction probably includes an initial electroneutral  $\text{Na}^+$  association process through an aqueous environment, followed by electrogenic occlusion of the cation through water exclusion. The function of this first  $\text{Na}^+$ -binding step is modulatory because the high affinity of the transporter for this sodium presumably prevents dissociation from the transporter. It should be noted that we do not know yet the affinity for  $\text{Na}^+$  of the inward-facing transporter. However, based on the predicted modulatory function of this  $\text{Na}^+$ -binding site, this affinity is expected



**Figure 8.** Extended mechanism for amino acid transport by ASCT2. The rate constants for the exchange equilibrium were estimated from Fig. 3 and are given for alanine as the amino acid substrate. The  $\text{Na}^+$ /substrate dissociation sequence on the intracellular side is speculative and not based on experimental data. Electrogenic reactions are illustrated in red.

to be high. In fact, the  $K_m$  of the transporter for extracellular alanine is not significantly changed when the intracellular  $[Na^+]$  is lowered from 140 to 10 mM ( $K_m = 160 \pm 50 \mu M$ ; unpublished data). Therefore, 10 mM of cytosolic  $[Na^+]$  is still saturating the internal  $Na^+$ -binding site.

Binding of the first  $Na^+$  ion is most likely followed by binding of another  $Na^+$  to the empty transporter, resulting in a doubly  $Na^+$ -bound transporter. Both MD simulations and experimental data support the existence of a second  $Na^+$ -binding site on the amino acid-free transporter, although this evidence is less strong than for the first cation-binding step. The  $Na_2$ -bound transporter subsequently binds the amino acid substrate in an electroneutral fashion, followed by association with a third  $Na^+$  ion. The MD results clearly show that this ion, which is expected to bind to the  $Na_2$  site, is unstable in the absence of bound amino acid, suggesting that amino acid binding is a prerequisite for closure of RL2. An occupied  $Na_2$  site may further stabilize the closed RL2, but  $Na^+$  binding to the  $Na_2$  site by itself is clearly not sufficient to stabilize the closed RL2. Overall, this mechanism is in excellent agreement with models on sequential  $Na^+$ /substrate binding to the EAATs (Watzke et al., 2001; Huang and Tajkhorshid, 2008, 2010), suggesting that these aspects of the transport mechanism are conserved within the SLC1 family.

The pre-steady-state kinetic data suggest that amino acid translocation by ASCT2 is fast, occurring on a millisecond time scale. From these data, a maximum exchange rate at optimum voltage for alanine of  $\sim 150$ – $200 s^{-1}$  is estimated (see Appendix), although the existence of slower electroneutral steps that are not seen in our experiments cannot be fully excluded. These results suggest that ASCT2 can release or take up amino acids rapidly, as required if it is involved in processes such as the glutamate–glutamine cycle (Bröer et al., 1999; Deitmer et al., 2003), which recycles the neurotransmitter glutamate in the brain. In contrast to the exchange reaction, the conformational changes of the empty transporter are probably slow. However, the data shown in Fig. 3 A suggest that the rate of rearrangements of the empty transporter, although slow, is nonzero because some of the amino acid-binding sites are facing the extracellular side even in the absence of ions that can promote outward translocation ( $K^+$  inside). Despite this ability of the empty transporter to rearrange binding sites, this process is physiologically not significant because it is too slow to support measurable transport.

ASCT2, as a transporter of the amino acid glutamine, is expressed in many tissues, including the brain (Bröer and Brookes, 2001). It was proposed to be involved in the glutamate–glutamine cycle (Bröer and Brookes, 2001), which recycles the excitatory neurotransmitter glutamate. Recently, ASCT2 has also been shown to be up-regulated in rapidly growing tumor cells, and its knockout led to a


reduction in cell growth (Fuchs and Bode, 2005; Nicklin et al., 2009). Therefore, our results may assist in the understanding of the function of ASCT2 in its physiological environment as well as under pathophysiological conditions.

## Conclusions

In this paper, we demonstrate that an overall electroneutral amino acid exchanger can mediate voltage-dependent amino acid transport through transmembrane charge movement of the amino acid translocation reaction. The molecular basis of the charge movement in the translocation step is the cotranslocation of two to three positively charged  $Na^+$  ions with the zwitterionic, neutral amino acid. This finding is in contrast to previous reports, predicting coupling of amino acid transport to a single cotransported  $Na^+$  ion. The fully loaded complex bears positive charge, resulting in charge movement associated with the amino acid translocation process. However, electroneutral translocation steps may also exist. Binding of the  $Na^+$  ions is modulatory because of the high affinity of the  $Na^+$ -binding sites.

## APPENDIX

For the following simplified one-step amino acid exchange process,



the unidirectional rate for amino acid influx,  $J_1$ , can be written as:

$$J_1 = N_t \frac{k_1 k_{-1}}{k_1 + k_{-1}}. \quad (2)$$

Here,  $N_t$  is the total number of transporters in the membrane. The voltage dependence of flux can then be derived by using the theoretical expressions for the voltage dependence of the forward and reverse rate constants:

$$J_1(V) = N_t \frac{k_1(0)e^{-zFV/RT} k_{-1}(0)e^{zFV/RT}}{k_1(0)e^{-zFV/RT} + k_{-1}(0)e^{zFV/RT}}. \quad (3)$$

Here,  $V$  is the membrane potential;  $z$  is the valence; and  $F$ ,  $R$ , and  $T$  have their usual meaning. This equation represents a bell-shaped curve when plotted as a function of the voltage. The voltage of the maximum flux can be obtained by finding the voltage at which the first derivative is zero:

$$0 = N_t \frac{-k_1(0)k_{-1}(0) [k_1(0)ze^{-zFV/RT} k_{-1}(0)ze^{zFV/RT}]}{[k_1(0)e^{-zFV/RT} + k_{-1}(0)e^{zFV/RT}]^2}. \quad (4)$$

Solving this equation for  $V$  yields the expression:



$$V_{\max} = \frac{-\log(k_1(0)/k_{-1}(0))}{2z}. \quad (5)$$

As expected, the  $V_{\max}$  is 0 when the forward and backward rate constants are identical. If the valence,  $z$ , is positive, the maximum shifts to more positive potentials with increasing forward rate constant, a situation that would be consistent with our kinetic data and the voltage dependence of threonine flux determined by Bussolati et al. (1992). This effect is illustrated in Fig. A1, assuming a valence of  $\pm 0.5$ .

#### Derivation of the equation for the $[\text{Na}^+]$ dependence of the anion current

The total population of anion-conducting states,  $\overline{\text{TNa}^+}$  and  $\overline{\text{TAlaNa}_2^+}$ , according to the kinetic scheme in Fig. 2, is:

$$P = \frac{[\text{Na}^+](K_{\text{Na}_2}K_s + r[\text{Na}^+][\text{S}])}{K_{\text{Na}_2}K_s(K_{\text{Na}_1} + [\text{Na}^+]) + [\text{Na}^+][\text{S}](K_{\text{Na}_2} + [\text{Na}^+])}. \quad (6)$$

Here,  $K_{\text{Na}_1}$  and  $K_{\text{Na}_2}$  are the intrinsic dissociation constants for  $\text{Na}^+$ , as illustrated in Fig. 2 (bottom scheme),  $K_s$  is the intrinsic dissociation constant for the substrate, and  $r$  accounts for the ratio of conductances of the  $\overline{\text{TAlaNa}_2^+}$  over the  $\overline{\text{TNa}^+}$  state.


At saturating alanine concentration, the total current then becomes:

$$I_{\text{total}} = I_{\max}(\text{saturating alanine}) + I_0(\text{no alanine}) \quad (7)$$

$$= -\frac{[\text{Na}^+]}{(K_{\text{Na}_1} + [\text{Na}^+])} + r\frac{[\text{Na}^+]}{(K_{\text{Na}_2} + [\text{Na}^+])}. \quad (8)$$

The equations for the apparent dissociation constant for alanine,  $K_m$ , as a function of  $\text{Na}^+$  have been derived by us in Watzke et al. (2001):

$$K_m = K_s \cdot \left( \frac{K_{\text{Na}_1} + [\text{Na}^+]}{[\text{Na}^+]} \right) \cdot \left( \frac{K_{\text{Na}_2}}{K_{\text{Na}_2} + [\text{Na}^+]} \right). \quad (9)$$



**Figure A1.** Simulation of the voltage dependence of flux according to Eq. 3.

In case of two  $\text{Na}^+$  ions binding before the amino acid, the apparent  $K_m$  is:

$$K_m = K_s \cdot \left( \frac{K_{\text{Na}_1} + [\text{Na}^+]}{[\text{Na}^+]} \right)^2. \quad (10)$$

We are grateful to Dr. Stefan Bröer for helpful comments regarding the manuscript.

This work was supported by the National Institutes of Health (grant 2R01NS049335-06A1 to C. Grewer) and Binational Science Foundation (grant 2007051 to C. Grewer and Baruch I. Kanner).

Lawrence G. Palmer served as editor.

Submitted: 10 December 2012

Accepted: 22 April 2013

## REFERENCES

- Albers, T., W. Marsiglia, T. Thomas, A. Gameiro, and C. Grewer. 2012. Defining substrate and blocker activity of alanine-serine-cysteine transporter 2 (ASCT2) ligands with novel serine analogs. *Mol. Pharmacol.* 81:356–365. <http://dx.doi.org/10.1124/mol.111.075648>
- Baker, N.A., D. Sept, S. Joseph, M.J. Holst, and J.A. McCammon. 2001. Electrostatics of nanosystems: application to microtubules and the ribosome. *Proc. Natl. Acad. Sci. USA.* 98:10037–10041. <http://dx.doi.org/10.1073/pnas.181342398>
- Bastug, T., G. Heinzelmann, S. Kuyucak, M. Salim, R.J. Vandenberg, and R.M. Ryan. 2012. Position of the third  $\text{Na}^+$  site in the aspartate transporter GltPh and the human glutamate transporter, EAAT1. *PLoS ONE.* 7:e33058. <http://dx.doi.org/10.1371/journal.pone.0033058>
- Bergles, D.E., A.V. Tzingounis, and C.E. Jahr. 2002. Comparison of coupled and uncoupled currents during glutamate uptake by GLT-1 transporters. *J. Neurosci.* 22:10153–10162.
- Bode, B.P. 2001. Recent molecular advances in mammalian glutamine transport. *J. Nutr.* 131:2475S–2485S.
- Boudker, O., R.M. Ryan, D. Yernool, K. Shimamoto, and E. Gouaux. 2007. Coupling substrate and ion binding to extracellular gate of a sodium-dependent aspartate transporter. *Nature.* 445:387–393. <http://dx.doi.org/10.1038/nature05455>
- Bröer, A., N. Brookes, V. Ganapathy, K.S. Dimmer, C.A. Wagner, F. Lang, and S. Bröer. 1999. The astroglial ASCT2 amino acid transporter as a mediator of glutamine efflux. *J. Neurochem.* 73:2184–2194.
- Bröer, A., C. Wagner, F. Lang, and S. Bröer. 2000. Neutral amino acid transporter ASCT2 displays substrate-induced  $\text{Na}^+$  exchange and a substrate-gated anion conductance. *Biochem. J.* 346:705–710. <http://dx.doi.org/10.1042/0264-6021:3460705>
- Bröer, S., and N. Brookes. 2001. Transfer of glutamine between astrocytes and neurons. *J. Neurochem.* 77:705–719. <http://dx.doi.org/10.1046/j.1471-4159.2001.00322.x>
- Bussolati, O., P.C. Laris, B.M. Rotoli, V. Dall’Asta, and G.C. Gazzola. 1992. Transport system ASC for neutral amino acids. An electro-neutral sodium/amino acid cotransport sensitive to the membrane potential. *J. Biol. Chem.* 267:8330–8335.
- Callenberg, K.M., O.P. Choudhary, G.L. de Forest, D.W. Gohara, N.A. Baker, and M. Grabe. 2010. APBSmem: a graphical interface for electrostatic calculations at the membrane. *PLoS ONE.* 5:e12722. <http://dx.doi.org/10.1371/journal.pone.0012722>
- Choudhary, O.P., R. Ujwal, W. Kowallis, R. Coalson, J. Abramson, and M. Grabe. 2010. The electrostatics of VDAC: implications for selectivity and gating. *J. Mol. Biol.* 396:580–592. <http://dx.doi.org/10.1016/j.jmb.2009.12.006>
- Crisman, T.J., S. Qu, B.I. Kanner, and L.R. Forrest. 2009. Inward-facing conformation of glutamate transporters as revealed by their

- inverted-topology structural repeats. *Proc. Natl. Acad. Sci. USA*. 106:20752–20757. <http://dx.doi.org/10.1073/pnas.0908570106>
- Darden, T., D.M. York, and L. Pedersen. 1993. Particle mesh Ewald: an  $N \log(N)$  method for Ewald sums in large systems. *J. Chem. Phys.* 98:10089–10092. <http://dx.doi.org/10.1063/1.464397>
- De Weer, P., D.C. Gadsby, and R.F. Rakowski. 1988. Overview: stoichiometry and voltage dependence of the Na/K pump. *Prog. Clin. Biol. Res.* 268A:421–434.
- Deitmer, J.W., A. Bröer, and S. Bröer. 2003. Glutamine efflux from astrocytes is mediated by multiple pathways. *J. Neurochem.* 87:127–135. <http://dx.doi.org/10.1046/j.1471-4159.2003.01981.x>
- Eswar, N., B. John, N. Mirkovic, A. Fiser, V.A. Ilyin, U. Pieper, A.C. Stuart, M.A. Marti-Renom, M.S. Madhusudhan, B. Yerkovich, and A. Sali. 2003. Tools for comparative protein structure modeling and analysis. *Nucleic Acids Res.* 31:3375–3380. <http://dx.doi.org/10.1093/nar/gkg543>
- Forster, I.C., K. Köhler, J. Biber, and H. Murer. 2002. Forging the link between structure and function of electrogenic cotransporters: the renal type IIa Na<sup>+</sup>/Pi cotransporter as a case study. *Prog. Biophys. Mol. Biol.* 80:69–108. [http://dx.doi.org/10.1016/S0079-6107\(02\)00015-9](http://dx.doi.org/10.1016/S0079-6107(02)00015-9)
- Fuchs, B.C., and B.P. Bode. 2005. Amino acid transporters ASCT2 and LAT1 in cancer: partners in crime? *Semin. Cancer Biol.* 15:254–266. <http://dx.doi.org/10.1016/j.semcancer.2005.04.005>
- Grabe, M., H. Lecar, Y.N. Jan, and L.Y. Jan. 2004. A quantitative assessment of models for voltage-dependent gating of ion channels. *Proc. Natl. Acad. Sci. USA*. 101:17640–17645. <http://dx.doi.org/10.1073/pnas.0408116101>
- Grewer, C., and E. Grabsch. 2004. New inhibitors for the neutral amino acid transporter ASCT2 reveal its Na<sup>+</sup>-dependent anion leak. *J. Physiol.* 557:747–759. <http://dx.doi.org/10.1113/jphysiol.2004.062521>
- Grewer, C., N. Watzke, T. Rauen, and A. Bicho. 2003. Is the glutamate residue Glu-373 the proton acceptor of the excitatory amino acid carrier 1? *J. Biol. Chem.* 278:2585–2592. <http://dx.doi.org/10.1074/jbc.M207956200>
- Grewer, C., Z. Zhang, J. Mwaura, T. Albers, A. Schwartz, and A. Gameiro. 2012. Charge compensation mechanism of a Na<sup>+</sup>-coupled, secondary active glutamate transporter. *J. Biol. Chem.* 287:26921–26931. <http://dx.doi.org/10.1074/jbc.M112.364059>
- Groeneveld, M., and D.J. Slotboom. 2010. Na<sup>(+)</sup>:aspartate coupling stoichiometry in the glutamate transporter homologue Glt(Ph). *Biochemistry*. 49:3511–3513. <http://dx.doi.org/10.1021/bi100430s>
- Heinzelmann, G., T. Baştuğ, and S. Kuyucak. 2011. Free energy simulations of ligand binding to the aspartate transporter Glt(Ph). *Biophys. J.* 101:2380–2388. <http://dx.doi.org/10.1016/j.bpj.2011.10.010>
- Huang, Z., and E. Tajkhorshid. 2008. Dynamics of the extracellular gate and ion-substrate coupling in the glutamate transporter. *Biophys. J.* 95:2292–2300. <http://dx.doi.org/10.1529/biophysj.108.133421>
- Huang, Z., and E. Tajkhorshid. 2010. Identification of the third Na<sup>+</sup> site and the sequence of extracellular binding events in the glutamate transporter. *Biophys. J.* 99:1416–1425. <http://dx.doi.org/10.1016/j.bpj.2010.06.052>
- Humphrey, W., A. Dalke, and K. Schulten. 1996. VMD: visual molecular dynamics. *J. Mol. Graph.* 14:33–38. [http://dx.doi.org/10.1016/0263-7855\(96\)00018-5](http://dx.doi.org/10.1016/0263-7855(96)00018-5)
- Larsson, H.P., X. Wang, B. Lev, I. Bacconguis, D.A. Caplan, N.P. Vyleta, H.P. Koch, A. Diez-Sampedro, and S.Y. Noskov. 2010. Evidence for a third sodium-binding site in glutamate transporters suggests an ion/substrate coupling model. *Proc. Natl. Acad. Sci. USA*. 107:13912–13917. <http://dx.doi.org/10.1073/pnas.1006289107>
- Loo, D.D., A. Hazama, S. Supplisson, E. Turk, and E.M. Wright. 1993. Relaxation kinetics of the Na<sup>+</sup>/glucose cotransporter. *Proc. Natl. Acad. Sci. USA*. 90:5767–5771. <http://dx.doi.org/10.1073/pnas.90.12.5767>
- Lu, C.C., and D.W. Hilgemann. 1999. GAT1 (GABA:Na<sup>+</sup>:Cl<sup>-</sup>) cotransport function. Steady state studies in giant *Xenopus* oocyte membrane patches. *J. Gen. Physiol.* 114:429–444. <http://dx.doi.org/10.1085/jgp.114.3.429>
- Mim, C., Z. Tao, and C. Grewer. 2007. Two conformational changes are associated with glutamate translocation by the glutamate transporter EAAC1. *Biochemistry*. 46:9007–9018. <http://dx.doi.org/10.1021/bi7005465>
- Mwaura, J., Z. Tao, H. James, T. Albers, A. Schwartz, and C. Grewer. 2012. Protonation state of a conserved acidic amino acid involved in Na<sup>(+)</sup> binding to the glutamate transporter EAAC1. *ACS Chem Neurosci*. 3:1073–1083. <http://dx.doi.org/10.1021/cn300163p>
- Nicklin, P., P. Bergman, B. Zhang, E. Triantafellow, H. Wang, B. Nyfeler, H. Yang, M. Hild, C. Kung, C. Wilson, et al. 2009. Bidirectional transport of amino acids regulates mTOR and autophagy. *Cell*. 136:521–534. <http://dx.doi.org/10.1016/j.cell.2008.11.044>
- Phillips, J.C., R. Braun, W. Wang, J. Gumbart, E. Tajkhorshid, E. Villa, C. Chipot, R.D. Skeel, L. Kalé, and K. Schulten. 2005. Scalable molecular dynamics with NAMD. *J. Comput. Chem.* 26:1781–1802. <http://dx.doi.org/10.1002/jcc.20289>
- Pines, G., Y. Zhang, and B.I. Kanner. 1995. Glutamate 404 is involved in the substrate discrimination of GLT-1, a (Na<sup>+</sup> + K<sup>+</sup>)-coupled glutamate transporter from rat brain. *J. Biol. Chem.* 270:17093–17097. <http://dx.doi.org/10.1074/jbc.270.29.17093>
- Reyes, N., C. Ginter, and O. Boudker. 2009. Transport mechanism of a bacterial homologue of glutamate transporters. *Nature*. 462:880–885. <http://dx.doi.org/10.1038/nature08616>
- Rostkowski, M., M.H. Olsson, C.R. Søndergaard, and J.H. Jensen. 2011. Graphical analysis of pH-dependent properties of proteins predicted using PROPKA. *BMC Struct. Biol.* 11:6. <http://dx.doi.org/10.1186/1472-6807-11-6>
- Silva, J.R., H. Pan, D. Wu, A. Nekouzadeh, K.F. Decker, J. Cui, N.A. Baker, D. Sept, and Y. Rudy. 2009. A multiscale model linking ion-channel molecular dynamics and electrostatics to the cardiac action potential. *Proc. Natl. Acad. Sci. USA*. 106:11102–11106. <http://dx.doi.org/10.1073/pnas.0904505106>
- Tao, Z., Z. Zhang, and C. Grewer. 2006. Neutralization of the aspartic acid residue Asp-367, but not Asp-454, inhibits binding of Na<sup>+</sup> to the glutamate-free form and cycling of the glutamate transporter EAAC1. *J. Biol. Chem.* 281:10263–10272. <http://dx.doi.org/10.1074/jbc.M510739200>
- Tao, Z., N. Rosental, B.I. Kanner, A. Gameiro, J. Mwaura, and C. Grewer. 2010. Mechanism of cation binding to the glutamate transporter EAAC1 probed with mutation of the conserved amino acid residue Thr101. *J. Biol. Chem.* 285:17725–17733. <http://dx.doi.org/10.1074/jbc.M110.121798>
- Utsunomiya-Tate, N., H. Endou, and Y. Kanai. 1996. Cloning and functional characterization of a system ASC-like Na<sup>+</sup>-dependent neutral amino acid transporter. *J. Biol. Chem.* 271:14883–14890. <http://dx.doi.org/10.1074/jbc.271.25.14883>
- Valdeolmillos, M., J. García-Sancho, and B. Herreros. 1986. Differential effects of transmembrane potential on two Na<sup>+</sup>-dependent transport systems for neutral amino acids. *Biochim. Biophys. Acta*. 858:181–187. [http://dx.doi.org/10.1016/0005-2736\(86\)90304-4](http://dx.doi.org/10.1016/0005-2736(86)90304-4)
- Wadiche, J.I., and M.P. Kavanaugh. 1998. Macroscopic and microscopic properties of a cloned glutamate transporter/chloride channel. *J. Neurosci.* 18:7650–7661.
- Wadiche, J.I., J.L. Arriza, S.G. Amara, and M.P. Kavanaugh. 1995. Kinetics of a human glutamate transporter. *Neuron*. 14:1019–1027. [http://dx.doi.org/10.1016/0896-6273\(95\)90340-2](http://dx.doi.org/10.1016/0896-6273(95)90340-2)

- Watzke, N., E. Bamberg, and C. Grever. 2001. Early intermediates in the transport cycle of the neuronal excitatory amino acid carrier EAAC1. *J. Gen. Physiol.* 117:547–562. <http://dx.doi.org/10.1085/jgp.117.6.547>
- Yernool, D., O. Boudker, Y. Jin, and E. Gouaux. 2004. Structure of a glutamate transporter homologue from *Pyrococcus horikoshii*. *Nature*. 431:811–818. <http://dx.doi.org/10.1038/nature03018>
- Zerangue, N., and M.P. Kavanaugh. 1996a. ASCT-1 is a neutral amino acid exchanger with chloride channel activity. *J. Biol. Chem.* 271:27991–27994. <http://dx.doi.org/10.1074/jbc.271.45.27991>
- Zerangue, N., and M.P. Kavanaugh. 1996b. Flux coupling in a neuronal glutamate transporter. *Nature*. 383:634–637. <http://dx.doi.org/10.1038/383634a0>
- Zhang, Z., G. Papageorgiou, J.E. Corrie, and C. Grever. 2007. Pre-steady-state currents in neutral amino acid transporters induced by photolysis of a new caged alanine derivative. *Biochemistry*. 46:3872–3880. <http://dx.doi.org/10.1021/bi0620860>