

## Research Article

# No Polymorphism in *Plasmodium falciparum* K13 Propeller Gene in Clinical Isolates from Kolkata, India

Moytrej Chatterjee,<sup>1</sup> Swagata Ganguly,<sup>1,2</sup> Pabitra Saha,<sup>1,3</sup> Biswabandhu Bankura,<sup>4</sup> Nandita Basu,<sup>5</sup> Madhusudan Das,<sup>4</sup> Subhasish K. Guha,<sup>6</sup> and Ardhendu K. Maji<sup>1</sup>

<sup>1</sup>Protozoology Unit, Department of Microbiology, Calcutta School of Tropical Medicine, 108 Chittaranjan Avenue, Kolkata 700 073, India

<sup>2</sup>Department of Microbiology, NRS Medical College, 138 AJC Bose Road, Kolkata 700 014, India

<sup>3</sup>Department of Zoology, APC Roy Government College, Himachal Bihar, Matigara, Siliguri, West Bengal 734 010, India

<sup>4</sup>Department of Zoology, Ballygunge Science College, University of Calcutta, 35 Ballygunge Circular Road, Kolkata 700 019, India

<sup>5</sup>Calcutta School of Tropical Medicine, 108 Chittaranjan Avenue, Kolkata 700 073, India

<sup>6</sup>Department of Tropical Medicine, Calcutta School of Tropical Medicine, 108 Chittaranjan Avenue, Kolkata 700 073, India

Correspondence should be addressed to Ardhendu K. Maji; maji.ardhendu@yahoo.com

Received 19 August 2015; Accepted 11 November 2015

Academic Editor: Mariela A. Segura

Copyright © 2015 Moytrej Chatterjee et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Molecular markers associated with artemisinin resistance in *Plasmodium falciparum* are yet to be well defined. Recent studies showed that polymorphisms in K13 gene are associated with artemisinin resistance. The present study was designed to know the pattern of polymorphisms in propeller region of K13 gene among the clinical isolates collected from urban Kolkata after five years of ACT implementation. We collected 59 clinical isolates from urban Kolkata and sequenced propeller region of K13 gene in 51 isolates successfully. We did not find any mutation in any isolate. All patients responded to the ACT, a combination of artesunate + sulphadoxine-pyrimethamine. The drug regimen is still effective in the study area and there is no sign of emergence of resistance against artemisinin as evidenced by wild genotype of K13 gene in all isolates studied.

## 1. Introduction

Emergence of resistance against different antimalarial drugs (AMDs) particularly in *Plasmodium falciparum* is a serious problem in combating malaria. During the 1990s *P. falciparum* acquired resistance to chloroquine (CQ) and sulphadoxine-pyrimethamine (SP) throughout all malaria endemic countries. World Health Organization (WHO) recommended artemisinin combination therapy (ACT) as first line agent to treat all uncomplicated *falciparum* malaria throughout the world. After introduction of the ACTs and increased use of insecticide treated bed nets (ITBNs), the mortality and morbidity due to *falciparum* malaria reduced globally [1] including India [2, 3]. Unfortunately, within a few years *P. falciparum* developed resistance against artemisinin. Artemisinin resistance is defined by longer parasite clearance

time in in vivo efficacy study and in in vitro ring stage survival assay [4–6]. Artemisinin resistance was first reported from Western Cambodia [7, 8] in 2009. Now it has spread throughout Southeast Asian countries [6, 9–13].

ACT, a combination of artesunate + sulphadoxine-pyrimethamine (AS + SP), was first introduced in India in 2009 to treat all uncomplicated *falciparum* cases. Previously ACT was recommended only in some high risk districts like Purulia and Jalpaiguri of West Bengal. Several efficacy studies showed that this combination is still effective in various parts of the country [14, 15] except in northeastern part [13] where AS + SP has been replaced by artemether + lumefantrine. Antimalarial drug efficacy monitoring every two years can predict the emergence of drug resistance, if any. Though in vivo therapeutic efficacy study remains the gold standard for the purpose, molecular markers have also been validated as

TABLE 1: Primers and PCR conditions of primary and nested PCR of *K13* gene.

Primer name	Primer sequence (5'-3')	Mg <sup>2+</sup> Conc. (mM)	PCR programme						Number of cycles	
			Denaturation Temp. (°C)	Time (min)	Annealing Temp. (°C)	Time (min)	Elongation Temp. (°C)	Time (min)	1st PCR	2nd PCR
K13_PCR_F K13_PCR_R	CGGAGTGACCAAATCTGGGA GGGAATCTGGTGGTAACAGC	3	94	0:30	60	1:30	72	1:30	40	—
K13_N1_F K13_N1_R	GCCAAGCTGCCATTCATTTG GCCTTGTGAAAGAAGCAGA	2.5	94	0:30	60	1:30	72	1:30	—	40

alternative tool. A single mutation, K76T in *pfprt* gene, is a hallmark of CQ resistance [16, 17] and a quintuple mutation in *pfdhfr* and *pfdhps* genes strongly correlated with SP resistance [18]. Recently polymorphisms in the propeller region of *P. falciparum* Kelch protein (*K13* gene) have been found to be associated with artemisinin resistance [19]. Polymorphisms in *K13* gene have been reported from different countries [9, 19–26] but such report from India is scarce. Recently a report showed that mutations in *K13* gene are limited [27] in the strains isolated from northeastern part of the country. The present study deals with the pattern of polymorphisms in propeller region of *K13* gene among the clinical isolates collected from Kolkata, India, after five years of ACT implementation.

## 2. Materials and Methods

**2.1. Study Site and Collection of Blood Samples.** The study was conducted at the Malaria Clinic attached to Protozoology Unit of the Calcutta School of Tropical Medicine, Kolkata, from January to December, 2014. In routine practice diagnosis was done by microscopy following Giemsa staining of thick and thin blood films. Report and medicines were distributed on the same day. All treated *P. falciparum* cases were advised to attend the clinic for reexamination on days 1, 2, 3, 7, 14, 21, 28, 35, and 42. Two to three mL EDTA blood samples were collected from the microscopically *P. falciparum* positive patients during distribution of report and medicine after obtaining informed consent or assent from the patients/guardians of the children below 14 years of age. The study protocol was reviewed and cleared by Institutional Ethics Committee of the Calcutta School of Tropical Medicine, Kolkata.

**2.2. DNA Isolation.** *P. falciparum* genomic DNA was isolated from whole blood samples using QIAamp DNA blood mini kit (Qiagen, Hilden, Germany), following the manufacturer's instructions. Extracted DNA of all the samples was preserved at -20°C and an aliquot was used as the DNA source for further study.

**2.3. Nested PCR and Sequencing of *K13* Gene Propeller Region.** The propeller domain of *K13* gene was amplified by nested PCR method as described by Ariey et al., 2014. All amplification reactions were carried out in a final volume of 20 µL for primary PCR and 50 µL for nested PCR. The oligonucleotide primers and PCR conditions are summarized in Table 1.

The quality and concentration of PCR products were ascertained by 1.5% agarose gel electrophoresis following ethidium bromide stain. The nested PCR products were purified using QIAquick PCR Purification Kit (Qiagen). Sequencing was outsourced from Amnion Biosciences Pvt. Ltd., Bangalore, India.

**2.4. Analysis of Sequence.** The sequences were analyzed using the free software Bioedit Sequence Alignment Editor version 7.0.5.2 and aligned with sequences of PF3D7\_1343700 Kelch protein propeller domain using the online sequence alignment tool ClustalW.

## 3. Results

During the study period a total of 8877 febrile patients attended the clinic for diagnosis and treatment for malaria. Among them 1223 were positive for *P. vivax* and 107 were positive for *P. falciparum*. The presence of mixed infection with both *P. vivax* and *P. falciparum* cannot be ruled out as a confirmatory molecular method was not performed. Out of 107 *P. falciparum* cases blood samples were collected from 59 patients who consented. Though it was not a therapeutic efficacy study all treated patients were followed up both clinically and parasitologically as mentioned earlier. It was found that all 59 patients responded to AS + SP therapy, that is, no case of recrudescence recorded during six-week posttreatment period.

Out of 59 samples, sequencing of propeller region of *K13* gene was done in 51 isolates. After alignment with PF3D7\_1343700, it was found that all 51 sequences were perfectly aligned from nucleotide number 1327 to 1998 (codons 443–666), that is, from blade 1 to blade 5 of the propeller region of *K13* gene. In blade 6 sequences were studied partly. All the strains were wild type having no polymorphism in this region of the gene. The nucleotide sequence of *K13* gene of *P. falciparum* isolate from the study site has been submitted in GenBank under Accession number KR779827.1.

## 4. Discussion

Unlike chloroquine and sulphadoxine-pyrimethamine resistance, artemisinin resistance in *P. falciparum* is yet to be well defined as definite phenotypes have not been well established except delayed parasite clearance following chemotherapy

and ring stage survival assay (RSA) [7, 9]. These parameters are indications of emergence of resistance by the parasites against artemisinin derivatives. WHO [28] defined suspected partial artemisinin resistance as follows: persistent parasitemia by microscopy on day 3 after treatment with ACT/artesunate monotherapy in 10%/more patients or presence of resistance-associated mutations in *K13* gene in at least 5% of patients or parasite clearance half-life is more than 5 hours after treatment. Confirmed partial artemisinin resistance is defined as presence of *K13* resistance-associated mutations in at least 5% of patients to have either persistent parasitaemia by microscopy on day 3 or a parasite clearance half-life of  $\geq 5$  hours after treatment with ACT or artesunate monotherapy. To conserve the efficacy of artemisinin derivatives, monotherapy with this drug has been stopped and is recommended with another long acting partner drug. So efficacy study of any ACT does not deal with efficacy of artemisinin alone. As definite phenotype of ACT resistance is very rare, the association of polymorphisms in marker genes is very difficult to correlate with efficacy outcome. Ariey et al., 2014 [19], have recently shown that four polymorphisms, C580Y, Y493H, R539T, and M476L, in propeller region of *K13* gene are strongly associated with artemisinin resistance. Following this discovery several workers reported a number of SNPs in this gene particularly at propeller region from different parts of the world [9, 19–26].

In the present study, we found that parasite was cleared in 53 cases on day 1 and in the remaining 6 cases on day 2. We did not find any mutation in the propeller part of the gene and reports of such mutation from this country are also rare [27]. However mutation at A578S which is adjacent to the C580Y, the major mutation causing delayed parasite clearance, was reported from neighboring country Bangladesh [22] and mutation at F446I from Myanmar [26].

## 5. Conclusion

ACT has been introduced in Kolkata just five years back. As we have not taken into account the parasite clearance half-life or ring stage survival assay in the study and the sample size is also small so it is difficult to comment on emergence of resistance against it. If the association of identified SNPs with artemisinin resistance is true, then this base-line information indicates that in urban Kolkata the prevailing *falciparum* population is still sensitive to artemisinin. Similar studies from different parts of the country and their association with delayed parasite clearance and RSA will be helpful to ascertain the emergence of artemisinin resistance, if any. Both quadruple and quintuple mutations in *psdhfr* and *psdhps* genes have been reported from different parts of the country [29]. As SP is used as partner drug of ACT, the in vivo efficacy study of this combination should be monitored at a regular interval of two years.

## Conflict of Interests

The authors have no conflict of interests concerning the work reported in this paper.

## Authors' Contribution

Ardhendu K. Maji and Swagata Ganguly designed the study; Moytrej Chatterjee, Swagata Ganguly, Pabitra Saha, Subhasish K. Guha, and Nandita Basu clinically examined, recruited, and followed up the patients and collected the blood samples; Swagata Ganguly provided the treatment; Moytrej Chatterjee, Swagata Ganguly, and Pabitra Saha assessed laboratory diagnostics and molecular studies. Moytrej Chatterjee, Pabitra Saha, Biswabandhu Bankura, and Madhusudan Das performed the data analysis and interpretation. Ardhendu K. Maji, Swagata Ganguly, Moytrej Chatterjee, and Nandita Basu prepared the paper.

## Acknowledgments

The authors are thankful to all the patients who participated in this study and to the Department of Science and Technology, Government of West Bengal, for financially supporting the study.

## References

- [1] A. Bhattarai, A. S. Ali, S. P. Kachur et al., "Impact of artemisinin-based combination therapy and insecticide-treated nets on malaria burden in Zanzibar," *PLoS Medicine*, vol. 4, no. 11, pp. 1784–1790, 2007.
- [2] N. K. Shah, P. Tyagi, and S. K. Sharma, "The impact of artemisinin combination therapy and long-lasting insecticidal nets on forest malaria incidence in tribal villages of India, 2006–2011," *PLoS ONE*, vol. 8, no. 2, Article ID e56740, 2013.
- [3] P. Saha, S. Ganguly, S. Dutta et al., "Impact of artemisinin-based combination therapy on *Falciparum* malaria in urban Kolkata: a clinic-based report," *Japanese Journal of Infectious Diseases*, vol. 68, no. 4, pp. 321–323, 2015.
- [4] B. Witkowski, N. Khim, P. Chim et al., "Reduced artemisinin susceptibility of *Plasmodium falciparum* ring stages in western Cambodia," *Antimicrobial Agents and Chemotherapy*, vol. 57, no. 2, pp. 914–923, 2013.
- [5] B. Witkowski, C. Amaratunga, N. Khim et al., "Novel phenotypic assays for the detection of artemisinin-resistant *Plasmodium falciparum* malaria in Cambodia: in-vitro and ex-vivo drug-response studies," *The Lancet Infectious Diseases*, vol. 13, no. 12, pp. 1043–1049, 2013.
- [6] A. P. Phyto, S. Nkhoma, K. Stepniewska et al., "Emergence of artemisinin-resistant malaria on the western border of Thailand: a longitudinal study," *The Lancet*, vol. 379, no. 9830, pp. 1960–1966, 2012.
- [7] A. M. Dondorp, F. Nosten, P. Yi et al., "Artemisinin resistance in *Plasmodium falciparum* malaria," *The New England Journal of Medicine*, vol. 361, no. 5, pp. 455–467, 2009.
- [8] H. Noedl, Y. Se, K. Schaefer, B. L. Smith, D. Socheat, and M. M. Fukuda, "Evidence of artemisinin-resistant malaria in Western Cambodia," *The New England Journal of Medicine*, vol. 359, no. 24, pp. 2619–2620, 2008.
- [9] E. A. Ashley, M. Dhorda, R. M. Fairhurst et al., "Spread of artemisinin resistance in *Plasmodium falciparum* malaria," *The New England Journal of Medicine*, vol. 371, pp. 411–423, 2014.
- [10] C. Amaratunga, S. Sreng, S. Suon et al., "Artemisinin-resistant *Plasmodium falciparum* in Pursat province, western Cambodia:

- a parasite clearance rate study," *The Lancet Infectious Diseases*, vol. 12, no. 11, pp. 851–858, 2012.
- [11] M. P. Kyaw, M. H. Nyunt, K. Chit et al., "Reduced susceptibility of *Plasmodium falciparum* to artesunate in Southern Myanmar," *PLoS ONE*, vol. 8, no. 3, Article ID e57689, 2013.
- [12] T. T. Hien, N. T. Thuy-Nhien, N. H. Phu et al., "In vivo susceptibility of *Plasmodium falciparum* to artesunate in Binh Phuoc Province, Vietnam," *Malaria Journal*, vol. 11, article 355, 2012.
- [13] N. Mishra, K. Kaitholia, B. Srivastava et al., "Declining efficacy of artesunate plus sulphadoxine-pyrimethamine in northeastern India," *Malaria Journal*, vol. 13, article 284, 2014.
- [14] P. Saha, A. Naskar, S. Ganguly et al., "Therapeutic efficacy of artemisinin combination therapies and prevalence of S769N mutation in PfATPase6 gene of *Plasmodium falciparum* in Kolkata, India," *Asian Pacific Journal of Tropical Medicine*, vol. 6, no. 6, pp. 443–448, 2013.
- [15] N. Mishra, J. P. N. Singh, B. Srivastava et al., "Monitoring antimalarial drug resistance in India via sentinel sites: outcomes and risk factors for treatment failure, 2009–2010," *Bulletin of the World Health Organization*, vol. 90, no. 12, pp. 895–904, 2012.
- [16] R. Durand, S. Jafari, J. Vauzelle, J.-F. Delabre, Z. Jesic, and J. Le Bras, "Analysis of pfprt point mutations and chloroquine susceptibility in isolates of *Plasmodium falciparum*," *Molecular and Biochemical Parasitology*, vol. 114, no. 1, pp. 95–102, 2001.
- [17] A. Djimdé, O. K. Doumbo, J. F. Cortese et al., "A molecular marker for chloroquine-resistant falciparum malaria," *The New England Journal of Medicine*, vol. 344, no. 4, pp. 257–263, 2001.
- [18] J. G. Kublin, F. K. Dzinjalama, D. D. Kamwendo et al., "Molecular markers for failure of sulfadoxine-pyrimethamine and chlorproguanil-dapsone treatment of *Plasmodium falciparum* malaria," *Journal of Infectious Diseases*, vol. 185, no. 3, pp. 380–388, 2002.
- [19] F. Arie, B. Witkowski, C. Amaratunga et al., "A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria," *Nature*, vol. 505, no. 7481, pp. 50–55, 2014.
- [20] E. Kamau, S. Campino, L. Amenga-Etego et al., "K13-propeller polymorphisms in *Plasmodium falciparum* parasites from sub-Saharan Africa," *The Journal of Infectious Diseases*, vol. 211, no. 8, pp. 1352–1355, 2015.
- [21] M. D. Conrad, V. Bigira, J. Kapisi et al., "Polymorphisms in K13 and falcipain-2 associated with artemisinin resistance are not prevalent in *Plasmodium falciparum* isolated from Ugandan children," *PLoS ONE*, vol. 9, no. 8, Article ID e105690, 2014.
- [22] A. N. Mohon, M. S. Alam, A. G. Bayih et al., "Mutations in *Plasmodium falciparum* K13 propeller gene from Bangladesh (2009–2013)," *Malaria Journal*, vol. 13, article 431, 2014.
- [23] P. Bosman, J. Stassijns, F. Nackers et al., "*Plasmodium* prevalence and artemisinin-resistant falciparum malaria in Preah Vihear Province, Cambodia: a cross-sectional population-based study," *Malaria Journal*, vol. 13, article 394, 2014.
- [24] S. Takala-Harrison, C. G. Jacob, C. Arze et al., "Independent emergence of artemisinin resistance mutations among *Plasmodium falciparum* in Southeast Asia," *The Journal of Infectious Diseases*, vol. 211, no. 5, pp. 670–679, 2015.
- [25] S. M. Taylor, C. M. Parobek, D. K. DeConti et al., "Absence of putative artemisinin resistance mutations among *Plasmodium falciparum* in Sub-Saharan Africa: a molecular epidemiologic study," *The Journal of Infectious Diseases*, vol. 211, no. 5, pp. 680–688, 2015.
- [26] K. M. Tun, M. Imwong, K. M. Lwin et al., "Spread of artemisinin-resistant *Plasmodium falciparum* in myanmar: a cross-sectional survey of the K13 molecular marker," *The Lancet Infectious Diseases*, vol. 15, no. 4, pp. 415–421, 2015.
- [27] N. Mishra, S. K. Prajapati, K. Kaitholia et al., "Surveillance of artemisinin resistance in *Plasmodium falciparum* in India using the kelch13 molecular marker," *Antimicrobial Agents and Chemotherapy*, vol. 59, no. 5, pp. 2548–2553, 2015.
- [28] WHO-2014—Status report on artemisinin resistance, September, <http://www.who.int/malaria/publications/atoz/status-rep-artemisinin-resistance-sep2014.pdf>.
- [29] A. Ahmed, D. Bararia, S. Vinayak et al., "*Plasmodium falciparum* isolates in india exhibit a progressive increase in mutations associated with sulfadoxine-pyrimethamine resistance," *Antimicrobial Agents and Chemotherapy*, vol. 48, no. 3, pp. 879–889, 2004.