



ORAL PRESENTATION

Open Access

Expanded memory CD4+ T Cells in the fetal and the infant Gut; a mucosal route for mother-to-child transmission of HIV-1

MJ Bunders*, C van der Loos, PL Klarenbeek, J van Hamme, J Wilde, N de Vries, RA van Lier, N Kootstra, ST Pals, TW Kuijpers

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

Cord blood-derived CD4+ T cells have a naïve phenotype and do not express CCR5, the mandatory co-receptor for transmitted HIV-1 R5 strains in infants. This leaves the question unanswered: what are the target cells for MTCT of HIV-1 and where do they reside? We hypothesized that in infant mucosal tissues, CD4+CCR5+ T cells may be present to facilitate mucosal transmission of HIV-1.

Methods

Using multicolor immuno-histochemistry, flowcytometry and next-generation sequencing of the T cell receptor, we analyzed various human fetal and infant tissues to identify memory CD4+ T cells as targets for HIV-1.

Results

Here, we demonstrate the previously unrecognized abundance of memory CD4+CCR5+ T cells in the human fetal and infant gut mucosa. CD4+ T cells from mesenteric lymph node were mostly naïve, similar to blood. T helper differentiation profiles as determined by transcription factors differed by tissue, with T-bet and ROR γ t predominantly expressed by memory T cells in the gut mucosa. Next-generation sequencing for high-resolution screening of the T-cell receptor β -chain repertoire of clonal T cells as a hallmark of memory cells, identified expanded T cell clones in the gut mucosa (30%) and not in lymph node or cord blood. The gut mucosal fetal and infant CD4+ T cells were extremely susceptible to HIV-1 without any prestimulation; pol proviral DNA levels were similar to infected PHA stimulated adult PBMCs.

Conclusion

In conclusion, we show that extensive adaptive immunity, with a tissue-dependent distribution is present before birth, resulting in the gut mucosa as the preferential site for memory CD4+ T cells. These memory CD4+CCR5+T cells provide a large pool of susceptible cells for ingested HIV-1 at birth and during breastfeeding, indicating a mucosal route of MTCT of HIV-1, which can be targeted in future prevention strategies.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-O21

Cite this article as: Bunders *et al.*: Expanded memory CD4+ T Cells in the fetal and the infant Gut; a mucosal route for mother-to-child transmission of HIV-1. *Retrovirology* 2012 **9**(Suppl 2):O21.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Academic Medical Centre, Amsterdam, the Netherlands



© 2012 Bunders *et al.*; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.