



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Janus kinase inhibitor baricitinib is not an ideal option for management of COVID-19

D. Praveen<sup>a</sup>, Ranadheer Chowdary Puvvada<sup>a</sup>, Vijey Aanandhi M<sup>b,\*</sup>

<sup>a</sup> Research Scholar, School of Pharmaceutical Sciences, Vels Institute of Science Technology and Advanced Studies, 600117, Chennai, India

<sup>b</sup> Department of Pharmaceutical Chemistry and Analysis, School of Pharmaceutical Sciences, Vels Institute of Science Technology and Advanced Studies, Chennai, India

### ARTICLE INFO

Editor: Jean-Marc Rolain

**Keywords:**

SARS-CoV-2

COVID 19

Baricitinib

Novel Corona Virus

### ABSTRACT

The Wuhan outbreak of novel Corona virus infection has been the global focus since December 2019. This infection has become a global pandemic. It is highly important to understand the virology of the pathogen and to explore the therapeutic options for management of this pandemic. Drug repurposing strategies are being considered for management of COVID 19. Among the identified drugs, Baricitinib has become a keen interest for researchers because of its ability to inhibit the viral assembly by the prevention of Clarithrin associated endocytosis. We tried to explore the reasons on why Baricitinib is not an ideal option for COVID 19.

© 2020 Elsevier B.V. and International Society of Chemotherapy. All rights reserved.



Drug repurposing strategies are being considered for the management of novel coronavirus disease 2019 (COVID-19). The janus kinase (JAK) inhibitor, baricitinib has gained considerable research interest because it can inhibit viral assembly by preventing AP-2 associated protein kinase 1 (AAK1)-mediated endocytosis.

The most important human receptor for the severe acute respiratory syndrome (SARS) S glycoprotein is the angiotensin-converting enzyme 2 (ACE2) [1]. Novel coronavirus has a similar glycoprotein that may also target this enzyme. ACE2 is predominantly found in the lower respiratory tract, particularly in the lung AT2 alveolar epithelial cells [2], which are prone to viral infections like SARS coronavirus [3]. These AT2 alveolar epithelial cells might contribute to viral reproduction and transmission through endocytosis [4]. AAK1 is a potential promoter of this endocytosis, thus enhancing viral assembly in the intracellular matrix [5]. Cyclin G-associated kinase is another regulator of this endocytosis [6].

Baricitinib is a potential option for the management of disease caused by novel coronaviruses. Baricitinib inhibits both AAK1 and cyclin G-associated kinase, thereby preventing endocytosis and reducing viral assembly. Baricitinib inhibits JAK1 and JAK2 and there-

fore may help manage inflammation [7]. Several studies suggested the use of baricitinib for treating COVID-19 [3,8].

Study results indicate that baricitinib cannot be initiated in patients with an absolute neutrophil count less than  $1 \times 10^9$  cells/L, or in patients with an absolute lymphocyte count less than  $0.5 \times 10^9$  cells/L [9]. Epidemiological studies show the values of the selected patients are closer to the threshold levels at baseline (Table 1) [10–14]. In one epidemiological study, absolute lymphocyte count in non-survivors was reported to be  $0 \cdot 6 \times 10^9$  cells/L (interquartile range:  $0 \cdot 5\text{--}0 \cdot 8 \times 10^9$  cells/L) (Table 2) [14]. Similarly, another study carried out by Huang et al. reported that absolute lymphocyte count in infected patients in ICU was  $0 \cdot 4 \times 10^9$  cells/L (interquartile range:  $0 \cdot 2\text{--}0 \cdot 8 \times 10^9$  cells/L) [12]. The risk of lymphocytopenia may affect the progression of COVID-19. A 26% incidence of anaemia has been reported in non-survivors of COVID-19 infection [14]. Baricitinib therapy is predicted to be associated with incidence of anaemia [15]; therefore, initiation of baricitinib therapy in patients with COVID-19 may further increase anaemia incidence [9].

Elevated creatine kinase has been observed in patients receiving baricitinib therapy [15]. Although the median value of creatine kinase is reported to be in the normal range ( $<175$  U/L) in patients with COVID-19, it is greatly increased in critically ill patients and non-survivors of this disease [10–14]. A total of 46% of patients with COVID-19 in ICU have been reported to have elevated creatine kinase levels [12]. In one critically ill COVID-19 patient, creatine kinase levels were as high as 493 U/L [12]. Elevated creatine kinase levels pose a risk for the initiation of baricitinib therapy.

\* Corresponding author: Dr. M. Vijey Aanandhi, M.Pharm., Ph.D, Professor and Head, Department of Pharmaceutical Chemistry and Analysis, School of Pharmaceutical Sciences, Vels Institute of Science Technology and Advanced Studies (VISTAS), Chennai.

E-mail addresses: [praveennandan.1993@gmail.com](mailto:praveennandan.1993@gmail.com) (D. Praveen), [ranadheer.pharmd@gmail.com](mailto:ranadheer.pharmd@gmail.com) (R.C. Puvvada), [hodpchemistry@velsuniv.ac.in](mailto:hodpchemistry@velsuniv.ac.in) (V. Aanandhi M).

**Table 1**

Published biochemical data of COVID-19 patients.

Parameter	Reference range	Wang et al <sup>10</sup>	Chen et al <sup>11</sup>	Huang et al <sup>12</sup>	Ng et al <sup>13</sup>	Zhou et al <sup>14</sup>
No. of patients	-	99	138	41	21	191
Absolute neutrophil count ( $\times 10^9$ cells/L)	2.0-7.4	3.0 (2.0-4.9)	5.0 (3.3-8.1)	5.0 (3.0-8.9)	3.33 (3-3.91)	NA
Absolute lymphocyte count ( $\times 10^9$ cells/L)	1.1-3.6	0.8 (0.6-1.1)	0.9 (0.5)	0.8 (0.6-1.1)	1.29 (0.7-1.65)	1 · 0 (0 · 6-1 · 3)
Creatine kinase (U/L)	<170	102 (62- 252)	85.0 (51-184)	132.0 (62-219)	78.0 (69-137)	21 · 5 (13 · 0-72 · 4)

Values are median (interquartile range). NA - Not Available

**Table 2**

Published biochemical data of COVID-19 patients in ICU and non-survivors.

Parameter	Reference range	Wang et al <sup>10</sup> (n=138)	Huang et al <sup>12</sup> (n=41)	Zhou et al <sup>14</sup> (n=191)
ICU/non-survivors	-	ICU	ICU	Non-survivors
No. of patients	-	36	13	54
Absolute lymphocyte count ( $\times 10^9$ cells/L)	1.1-3.6	0.8 (0.5-0.9)	0 · 4 (0 · 2-0 · 8)	0 · 6 (0 · 5-0 · 8)
No. of patients (%) with lymphocytopenia	-	NA	11 (85%)	41 (76%)
Creatine kinase (U/L)	<170	102 (62-252)	132 · 0 (82 · 0-493 · 0)	39 · 0 (19 · 5-151 · 0)
No. of patients (%) with elevated creatine kinase	-	NA	6/13 (46%)	11/52 (21%) <sup>#</sup>

Values are median (interquartile range). NA - Not Available. <sup>#</sup>Data not available for 2 patients

Limited data are available on the potential effects of baricitinib in the elderly population aged at least 75 years [15]. Zhou et al. reported that mortality is higher in elderly infected patients [14]. Furthermore, studies have shown increased incidence of respiratory tract infections (16.3%) and incidence of infective diseases (29-42%). Co-infection is one of the most common threats in the management of COVID-19 [10]. There is also a risk of re-activation of latent infections. Infected patients will also be at risk of tuberculosis and hepatitis B [16]. Researchers have concluded that baricitinib therapy can reactivate varicella-zoster, herpes simplex and Epstein-Barr virus strains [17]. Zhou et al. reported that 50% of patients with COVID-19 experienced secondary infections [14].

Baricitinib may not be an ideal drug of choice for management of COVID-19. Available therapeutic options must be explored to prevent mortality in patients with this disease.

## Declarations

**Funding:** No funding.

**Competing Interests** None.

**Ethical Approval** Not required.

## References

- [1] Gruber CC, Steinkellner G. Wuhan coronavirus 2019-nCoV—what we can find out on a structural bioinformatics level. Jan 23, 2020. <https://innophore.com/2019-ncov/> (accessed Feb 10, 2020).
- [2] Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. Lancet 2020;395(10224):565-74.
- [3] Richardson P, Griffin I, Tucker C, Smith D, Oechsle O, Phelan A, et al. Baricitinib as potential treatment for 2019-nCoV acute respiratory disease. Lancet 2020;395(10223):e30-1.
- [4] Pu SY, Xiao F, Schor S, Bekerman E, Zanini F, Barouch-Bentov R, et al. Feasibility and biological rationale of repurposing sunitinib and erlotinib for dengue treatment. Antiviral Res 2018;155:67-75.
- [5] Sorrell FJ, Szklarz M, Abdul Areez KR, Elkins JM, Knapp S. Family-wide structural analysis of human numb-associated protein kinases. Structure 2016;24:401-11.
- [6] Mesa RA. Ruxolitinib, a selective JAK1 and JAK2 inhibitor for the treatment of myeloproliferative neoplasms and psoriasis. IDrugs 2010;13(6):394-403.
- [7] Channappanavar R, Fett C, Mack M, Ten Eyck PP, Meyerholz DK, Perlman S. Sex-based differences in susceptibility to severe acute respiratory syndrome coronavirus infection. J Immunol 2017;198(10):4046-53.
- [8] Stebbing J, Phelan A, Griffin I, Tucker C, Oechsle O, Smith D, et al. COVID-19: combining antiviral and anti-inflammatory treatments. Lancet Infect Dis 2020;20(4):400-2.
- [9] Chaplin S. Baricitinib: a new oral treatment for rheumatoid arthritis. Prescriber 2017;28(6):44-6.
- [10] Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. JAMA 2020 Feb 7:1585. doi:10.1001/jama.2020.
- [11] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. Lancet 2020;395:507-13. doi:10.1016/S0140-6736(20)30211-7.
- [12] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet 2020;395:497-506. doi:10.1016/S0140-6736(20)30183-5.
- [13] Ng M, Lee E, Yang J, Yang F, Li X, Wang H, et al. Imaging profile of the COVID-19 infection: radiologic findings and literature review. Radiology: Cardiothoracic Imaging 2020;2(1):e200034.
- [14] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet 2020;395(10229):1054-62.
- [15] Ema.europa.eu. [cited 20 March 2020]. Available from: [https://www.ema.europa.eu/en/documents/product-information/olumiant-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/olumiant-epar-product-information_en.pdf).
- [16] Mehta SK, Crucian B, Pierson DL, Sams C, Stowe RP. Monitoring immune system function and reactivation of latent viruses in the Artificial Gravity Pilot Study. J Gravit Physiol 2007;14(1):21-5.
- [17] Harigai M, Winthrop K, Takeuchi T, Hsieh T, Chen Y, Smolen J, et al. Evaluation of hepatitis B virus in clinical trials of baricitinib in rheumatoid arthritis. RMD Open 2020;6(1):e001095.