



OPEN

The superior beneficial effects of exercise training versus hormone replacement therapy on skeletal muscle of ovariectomized rats

Sara Barros Silva^{1,2}, Kinulpe Honorato-Sampaio^{2,3,4,5}, Sabrina Paula Costa², Talita Emanuela Domingues^{2,3}, Timilly Mayra Martins da Cruz², Cíntia Maria Rodrigues^{2,3}, Karine Beatriz Costa^{2,3}, Jousielle Márcia dos Santos^{2,3}, Vanessa Kelly da Silva Lage^{2,3}, Thais Peixoto Gaiad^{1,6}, Ana Paula Santos^{1,6}, Marco Fabrício Dias-Peixoto^{2,3,4,6}, Cândido Celso Coimbra⁷, Adelina Martha dos Reis⁷, Raphael Escorsim Szawka⁷, Pedro Henrique Scheidt Figueiredo^{1,2,6}, Henrique Silveira Costa^{1,2,6}, Murilo Xavier Oliveira^{1,2,6}, Vanessa Amaral Mendonça^{1,2,3,4,6} & Ana Cristina Rodrigues Lacerda^{1,2,3,4,6}✉

Previous studies have highlighted the positive effects of Estradiol (E2) replacement therapy and physical exercise on skeletal muscle during menopause. However, the comparison effects of exercise training (ET) and estradiol replacement therapy during menopause on skeletal muscle have not been investigated to date. This study aimed to compare the effects of endurance exercise training *versus* E2 replacement therapy on mitochondrial density, redox status, and inflammatory biomarkers in the skeletal muscle of ovariectomized rats. Thirty female *Wistar* rats (12-week-old) were randomly assigned into three groups: Untrained ovariectomized rats (UN-OVX, n = 10); untrained ovariectomized rats treated with estradiol replacement therapy (E2-OVX); and, trained ovariectomized rats (TR-OVX). After ovariectomy, the E2-OVX rats were treated subcutaneously with E2 (implanted Silastic® capsule containing 360 µg of 17β-estradiol/mL) while the TR-OVX group performed an exercise training protocol (50–70% of maximal running speed on a treadmill, 60 min/day, 5 days/week for 8 weeks). After euthanasia, the soleus muscle was processed for histological and biochemical evaluations. Only exercise prevented the reduction of maximal oxygen consumption and increased mechanical efficiency (ME). While mitochondrial muscle density, total antioxidant capacity (FRAP), catalase (CAT) activity, and interleukin 10 levels were higher in TR-OVX, only OVX-E2 presented higher CAT activity and lower interleukin 6 levels. Endurance exercise training compared with E2 replacement therapy maintains the aerobic capacity improving the ME of OVX rats. In addition, only endurance exercise training raises the skeletal muscle mitochondrial content and tends to balance the redox and inflammatory status in the skeletal muscle of OVX rats.

The postmenopausal period is associated with several metabolic and musculoskeletal chronic disorders, such as osteoporosis, tendinopathies, and arthritis^{1,2}. Thus, maintaining skeletal muscle integrity during menopause plays a central role in preventing musculoskeletal disorders¹. The mechanisms through which ovarian hormone

¹Programa de Pós-Graduação em Reabilitação e Desempenho Funcional (PPGREab), Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Campus JK-Highway MGT-367-Km 583, N°. 5000-Alto da Jacuba, Diamantina 39100-000, Brazil. ²Centro Integrado de Pós-Graduação e Pesquisa em Saúde (CIPq-Saúde), Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, Brazil. ³Programa Multicêntrico de Pós-graduação Multicêntrico em Ciências Fisiológicas (PPGMCF), Sociedade Brasileira de Fisiologia (SBFis), Diamantina, Brazil. ⁴Programa de Pós-Graduação em Ciências da Saúde (PPGCS), Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, Brazil. ⁵Faculdade de Medicina, Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, Minas Gerais, Brazil. ⁶Faculdade de Ciências Biológicas e da Saúde, Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, Minas Gerais, Brazil. ⁷Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Minas Gerais, Brazil. ✉email: lacerda.acr@ufvjm.edu.br

deficiency negatively affects the muscle integrity during the postmenopausal period are multifactorial, involving a pro-inflammatory profile and mitochondrial dysfunction^{1,2}. Studies using ovariectomized (OVX) rats, the most common experimental menopause model³, have reported that the skeletal muscle from OVX rats presents a reduction in mitochondrial biogenesis and an increase in oxidative stress^{4,5}. In addition, ovariectomy alters the intramuscular levels of several cytokines. Ovariectomy enhances tumor necrosis factor- α (TNF- α) expression and interleukin 6 (IL-6) levels, pro-inflammatory cytokines associated with muscle dysfunction^{6–9}, and reduces interleukin 10 (IL-10) levels, an anti-inflammatory^{8,9}. Additionally, chronic stimulation of pro-inflammatory cytokines, e.g., TNF- α , impairs mitochondrial function and biogenesis, promoting a vicious pro-inflammatory cycle^{10,11}.

Many studies have highlighted the positive effects of estrogen replacement therapy during menopause^{4,5}. Moreover, a growing body of evidence has also proven several beneficial effects of physical exercise therapy in counteracting the deleterious effects of menopause in the skeletal and cardiac muscles. In previous studies, we and others found that exercise training improved the cardiovascular and cardiac function of OVX rats^{5–7}. However, comparison effects of endurance exercise training versus estradiol (E2) replacement therapy after menopause on skeletal muscle mitochondrial profile and redox/inflammatory status have not been investigated to date.

Given that endurance exercise training stimulates mitochondrial biogenesis and improves redox and inflammatory status in the skeletal muscle under physiological and pathological conditions^{12–14}, we hypothesized that endurance exercise training would have similar or higher benefits compared to E2 replacement therapy to the skeletal muscle mitochondrial content and the redox and inflammatory status in OVX rats. Thus, we aimed to investigate the effects of endurance exercise training versus E2 replacement therapy on muscle mitochondrial density and redox/inflammatory status in OVX rats.

Methods

Animals. 12-week-old female *Wistar* rats ($n=30$, body mass = 216 ± 2.54 g) were provided by the animal facility of the Universidade Federal de Minas Gerais, Brazil. The rats' age in this study was determined according to previous studies^{4,15,16} and based on the premise that at 12 weeks of age, *Wistar* female rats are sexually mature¹⁶.

The animals were maintained in a temperature-controlled room (22 °C), in a 12 h dark: light cycle, and had free access to standard chow (Nuvilab Nutrients LTDA, Colombo, PR, Brazil) and water. All groups received the same diet (free from phytoestrogen) throughout the experimental period. All rats were treated similarly in terms of daily manipulation. All surgical procedures and protocols used were approved by the Animal Use Ethics Committee of the Universidade Federal dos Vales do Jequitinhonha e Mucuri (protocol n°015/2019) and conducted in accordance with the National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals.

Experimental design and sampling. The rats were randomly assigned into three groups: (1) Untrained ovariectomized rats (UN-OVX, $n=10$), (2) Ovariectomized rats treated with Estradiol replacement (E2-OVX, $n=10$), and (3) Trained ovariectomized rats (TR-OVX, $n=10$). Of note, all animals were ovariectomized and received a subcutaneously implanted Silastic® capsule containing vehicle (UN-OVX and TR-OVX groups) or 17 β -estradiol (E2-OVX group). The UN-OVX and E2-OVX groups were similarly handled and placed near the treadmill during the training sessions of the TR-OVX group to match the same environmental exposure conditions. Two weeks after ovariectomy, TR-OVX rats were submitted to an endurance exercise training protocol for eight weeks. All animals were familiarized with the maximal effort treadmill test and exercise protocol.

The rats were euthanized by decapitation. Both right and left soleus muscles were harvested, washed in ice-cold PBS (0.15 M, pH 7.34), frozen in liquid nitrogen, and stored at – 80 °C. The left soleus muscles were processed for oxidative stress evaluation and the right ones for inflammatory analyses. In addition, posterior mid-belly fragments of the right soleus from three animals per group were dissected and chemically fixed for mitochondrial density assessment by transmission electron microscopy.

Ovariectomy. The animals were anesthetized (Ketamine 80 mg/kg + Xylazine 12 mg/kg), both lateral abdominal walls were trichotomized and an incision was made. The ovaries were located, the oviduct was sectioned to remove the ovaries and the incisions were stitched^{3,4}. The animals received one dose of antibiotics (Pentabiotic, 24,000 UI/kg) immediately after surgery, and two doses of analgesic (Flunixin meglumine, 2.5 mg/kg), immediately and 24 h after surgery. Animals had two weeks to recover from surgery before the maximal aerobic capacity test. All rats recovered successfully.

Estradiol (E2) replacement therapy. Immediately after ovariectomy, all animals received a subcutaneously implanted Silastic® capsule containing 360 μ g of 17 β -estradiol/mL in corn oil or vehicle (corn oil). The Silastic® capsules were made of 20-mm segments of Silastic® tubing (inner/outer diameter: 1.02/2.16 mm). An incision was made in the rat dorsal region (10 mm) to implant the Silastic® capsule using forceps¹⁵. The incision was subsequently stitched. Silastic® capsules were re-implanted after five weeks in order to maintain concentrations within the physiological range^{15,17}. The efficacy of this hormone replacement protocol was confirmed by previous studies^{4,15,18,19}.

Maximal aerobic exercise test. All animals were familiarized with running on the treadmill (0.3 km/h, 10 min/day, 5 days) (Panlab, Havard Apparatus, Spain)^{20,21}. All rats were familiarized successfully.

The maximal effort exercise test consisted of 3 m/min increments every 3 min until the rat could no longer keep pace^{20,21}. The purpose was to evaluate the maximal aerobic capacity and determine exercise training intensity. Maximal oxygen consumption was accessed during the maximal exercise test (VO_{2max}) by indirect

calorimetry (Panlab, Harvard Apparatus, Spain) coupled to the treadmill (airflow = 1.0 L/min). VO_{2max} was measured continuously by a computerized system (Metabolism, Panlab, Harvard Apparatus, Spain)²⁰.

Mechanical efficiency (ME) was calculated by the formula: $ME = (\text{workload}/\text{energetic cost}) \times 100$ ²². Workload (W; kgm) was calculated as $W = \text{body weight (kg)} \times \text{TTF} \times \text{treadmill speed (m min}^{-1}) \times \text{sine (treadmill inclination)}$, where TTF is time to fatigue (min).

Endurance exercise training. The TR-OVX group performed the exercise on a motor treadmill (Insight®, SP, Ribeirão Preto, Brazil) at low-moderate intensity (~ 50–70% maximal running speed) 60 min/day, 5 days/week for eight weeks (total of 40 sessions), with a gradual increase in speed from 0.7 to 1.2 km/h weekly^{20,22}.

Transmission electron microscopy. Fragments of the soleus muscle were fixed in Karnovsky's solution (2.5% glutaraldehyde and 2% paraformaldehyde) in 0.1 M cacodylate buffer pH 7.4 overnight at 4 °C. Then, samples were post-fixed in a mixture of 2% (w/v) osmium tetroxide and 1.5% (w/v) potassium ferrocyanide for a minimum of 2 h to enhance the contrast of organelles. Thereafter, samples were washed in distilled water and kept in 2% uranyl acetate (en bloc staining) overnight, serially dehydrated in graded ethanol baths, and embedded in Epon 812. Finally, 50 nm ultrathin sections were stained with Reynolds lead citrate. Transmission electron microscopy (TEM) was performed using a FEI Tecnai G2-12 Spirit at 80 kV. The images were acquired in a SIS-MegaView 3 CCD camera with 1376 × 1070 pixels. Twenty-four electron micrographs per animal were taken at a × 11,000 magnification. Images were randomly selected from central parts of muscle fibers and were analyzed with ImageJ. Volume densities (Vv) of mitochondria were determined with the classic point counting method using a 252-point-grid (500 × 500 nm grid) projected onto each image^{23,24}.

Redox status and antioxidant enzyme activities. The entire soleus muscle samples were defrosted gradually from – 80 to 4 °C. The left soleus muscles were processed for oxidative stress evaluation. Thereafter, they were homogenized in extraction solution (1 mL/muscle 100 g) containing PBS 01x (125 mL), NaCl (2.925 g), BSA (0.625 g), EDTA (46.5 mg), PMSF (2.125 mg), benzethonium chloride (5.6 mg), Tween 20 (62.5 µL), aprotinin (2.5 µ) using an manual macerator. To evaluate the level of lipid peroxidation in the skeletal muscle, 1 mL of the homogenate was centrifugated at 5000×g for 5 min at 4 °C. For the analysis of the activity and protein expression of the antioxidant enzymes in the skeletal muscle, 1 mL of the homogenate was centrifugated at 10,000×g for 5 min at 4 °C. The Bradford method using bovine serum albumin was used as a standard to determine the sample's protein levels²⁵. The thiobarbituric acid reaction with malondialdehyde was used to determine lipid peroxidation by thiobarbituric acid reactive substances (TBARS) levels²⁶. The ferric reducing ability of plasma (FRAP), i.e., the reduction of ferric-tripyridyltriazine [Fe(III)-TPTZ] complex to ferrous-tripyridyltriazine [Fe(II)-TPTZ] was used to determine the total antioxidant capacity²⁷. The quantification of superoxide dismutase activity (SOD) was based on the inhibition of the reaction between $O_2^{\cdot -}$ and pyrogallol²⁸. Catalase activity (CAT) was determined by measuring the decrease in H_2O_2 absorbance at 240 nm²⁹.

Inflammatory biomarkers. The entire soleus muscle samples were defrosted gradually from – 80 to 4 °C. The right soleus muscles were processed for inflammatory analyses. Thereafter, they were homogenized in extraction solution (1 mL/muscle 100 g) containing PBS 01x (125 mL), NaCl (2.925 g), BSA (0.625 g), EDTA (46.5 mg), PMSF (2.125 mg), benzethonium chloride (5.6 mg), Tween 20 (62.5 µL), aprotinin (2.5 µ) using a manual macerator. The homogenate was then centrifugated at 10,000×g for 10 min at 4 °C. The supernatant was separated and used for analyses of IL-6, IL-10, and TNF-α according to the manufacturer's instructions by ELISA kits (DuoSet, R&D Systems, United States).

Statistical analyses. Data are reported as mean ± standard error (S.E.M.). Differences between groups were analyzed using one or two-way ANOVA followed by Dunnett posthoc test. The confidence interval (CI) and effect size (ES) for each significant difference is also shown. The correlation between variables was evaluated using the Pearson coefficient. The significance level for all tests was set at 5%. Statistical analyses were performed with GraphPad Prism 5.0 and G Power 3.1.9.2.

Ethics approval. All surgical procedures and protocols used were approved by Animal Use Ethics Committee of the Universidade Federal dos Vales do Jequitinhonha e Mucuri (protocol nº015/2019) and conducted in accordance with National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals. Of note, we confirm that our work is reported as described by the ARRIVE guidelines.

Consent for publication. The researchers of this study confirm that they have given due consideration to protecting the intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing, we confirm that we have followed the regulations of our institutions concerning intellectual property.

Results

Figure 1 presents the body mass and the uterus/body mass ratio. Final body mass did not differ between groups after 10 weeks of ovariectomy (UN-OVX: 290.50 ± 6.38 g; E2-OVX: 287.0 ± 3.59 g; TR-OVX: 293.00 ± 3.88 g, Fig. 1A). However, the uterus/body mass ratio was higher in the E2-OVX rats compared with both UN-OVX and TR-OVX groups (E2-OVX: 0.00070 ± 0.00003 g; UN-OVX: 0.00030 ± 0.00002 g; and TR-OVX: 0.00039 ± 0.00002 g, Fig. 1B).




Figure 1. (A) Body mass and (B) uterus mass/body mass ratio of untrained ovariectomized rats (UN-OVX), ovariectomized rats treated with Estradiol replacement (E2-OVX), and trained ovariectomized rats (TR-OVX). N = 10 per group. Data are presented as mean \pm S.E.M. 95% confidence interval (CI). Effect size (ES) * $p < 0.05$ (TR-OVX vs E2-OVX). One-way ANOVA followed by Dunnet posthoc test.

Measurement	UN-OVX (n = 10)	E2-OVX (n = 10)	TR-OVX (n = 10)	CI (TR vs E2)	CI (UN vs TR)	ES	p ¹	p ²	p ³
VO_{2max} (mL kg⁻¹ min⁻¹)									
Initial	30.34 \pm 0.36	30.57 \pm 0.45	29.35 \pm 0.91	- 2.53 to 3.00	- 3.76 to 1.78	1.73	< 0.001	< 0.001	0.055
Final	25.48 \pm 0.75	24.84 \pm 0.62	30.59 \pm 0.62*	- 3.41 to 2.12	2.32 to 7.87				
Mechanical efficiency (%)									
Initial	18.98 \pm 0.75	19.55 \pm 0.81	21.91 \pm 1.20	- 0.05 to 0.06	- 0.02 to 0.08	0.95	< 0.001	< 0.001	< 0.001
Final	22.94 \pm 1.94	23.64 \pm 1.94	37.51 \pm 2.94* ^Δ	- 0.04 to 0.06	0.09 to 0.20				
Distance (m)									
Initial	233.00 \pm 15.44	248.40 \pm 22.08	255.20 \pm 13.70	- 90.97 to 121.70	- 84.10 to 128.60	1.56	< 0.001	0.079	< 0.001
Final	162.5 \pm 21.74	156.00 \pm 16.99	530.40 \pm 72.85*	- 112.80 to 99.85	262.60 to 475.30				
Time (s)									
Initial	1200.00 \pm 42.19	1237.00 \pm 58.85	1262.00 \pm 36.38	- 195.90 to 268.70	- 170.50 to 294.10	1.49	< 0.001	0.635	< 0.001
Final	970.70 \pm 71.99	955.20 \pm 60.94	1846.75 \pm 93.69*	- 247.80 to 216.80	643.40 to 1108.00				
Speed (m/s)									
Initial	21.8 \pm 0.70	22.5 \pm 1.00	22.5 \pm 0.50	- 0.20 to 0.27	- 0.20 to 0.27	1.41	< 0.001	0.640	< 0.001
Final	18.0 \pm 1.00	17.7 \pm 1.00	32.3 \pm 1.50*	- 0.25 to 0.22	0.62 to 1.10				

Table 1. Aerobic capacity. Data are presented as mean \pm S.E.M. 95% confidence interval (CI). untrained ovariectomized rats (UN-OVX), ovariectomized rats treated with Estradiol replacement (E2-OVX), and trained ovariectomized rats (TR-OVX). VO_{2max}: maximal oxygen consumption. N = 10 per group. Effect size (ES). p1 interaction, p2 time, p3 treatment. *TR-OVX vs UN-OVX and E2-OVX. ^ΔTR-OVX Final vs TR-OVX Initial. Two-way ANOVA followed by Dunnet posthoc test.

Table 1 shows the results from the maximal aerobic capacity test. Before the training protocol, VO_{2max}, mechanical efficiency, final distance and time, and average speed were similar among the groups. After the training period, mechanical efficiency improved only in TR-OVX rats. Moreover, VO_{2max}, mechanical efficiency, final distance and time, and average speed were higher in the TR-OVX rats compared with both UN-OVX and E2-OVX rats (Table 1).

Figure 2 highlights the results of skeletal muscle mitochondrial density. Of note, the skeletal muscle mitochondrial density was higher (~20%) in the TR-OVX compared with both E2-OVX and UN-OVX groups (UN-OVX: 15.59 \pm 0.47%; E2-OVX: 16.50 \pm 0.59%; TR-OVX: 19.48 \pm 0.57%; Fig. 2A–D).

Figure 3 shows the results of muscle redox status. TBARS levels did not differ among the groups (UN-OVX: 0.37 \pm 0.04; E2-OVX: 0.35 \pm 0.01; TR-OVX: 0.37 \pm 0.02 mmolMDA/mg protein, Fig. 3A). The total antioxidant capacity (FRAP) levels (UN-OVX: 436.10 \pm 25.06; E2-OVX: 433.20 \pm 23.94; TR-OVX: 563.60 \pm 10.80 mmolFeSO₄/L/mg protein, Fig. 3B) and catalase activity (UN-OVX: 0.67 \pm 0.03; E2-OVX: 0.80 \pm 0.026; TR-OVX:




Figure 2. Mitochondrial muscle density. (A). Quantification of total mitochondria muscle density in mid-belly fragments of the right soleus from untrained ovariectomized rats (UN-OVX), ovariectomized rats treated with Estradiol replacement (E2-OVX), and trained ovariectomized rats (TR-OVX). $N = 72$ fields from three animals per group. (B–D) Transmission electron micrographs (TEM) of transverse sections of muscle fibers (ultra-structural view), in which mitochondria is highlighted in green. Data are reported as mean \pm S.E.M. 95% confidence interval (CI). Effect size (ES) * $p < 0.01$. One-way ANOVA followed by Dunnett posthoc test.

0.96 ± 0.04 nmol/mg protein, Fig. 3C) were higher in the TR-OVX compared with both E2-OVX and UN-OVX groups. Otherwise, SOD activity was higher in the E2-OVX compared with both TR-OVX and UN-OVX rats (UN-OVX: 1.28 ± 0.01 ; E2-OVX: 1.40 ± 0.03 ; TR-OVX: 1.33 ± 0.00 USOD/mg protein, Fig. 3D).

Figure 4 presents the results of inflammatory biomarkers. TNF- α levels did not differ among the groups (UN-OVX: 283.4 ± 3.343 ; E2-OVX: 284.4 ± 6.652 ; TR-OVX: 271.5 ± 4.173 pg/mg, Fig. 4A). However, the IL-6 levels were lower (~44%) in the E2-OVX group compared with both TR-OVX and UN-OVX groups (UN-OVX: 378.9 ± 8.875 ; E2-OVX: 262.3 ± 13.13 ; TR-OVX: 398.5 ± 27.53 pg/mg, Fig. 4B), and the IL-10 levels were higher (~61%) in the TR-OVX rats compared with the E2-OVX and UN-OVX rats (UN-OVX: 232.3 ± 5.249 ; E2-OVX: 216.0 ± 14.18 ; TR-OVX: 375.9 ± 15.97 pg/mg, Fig. 4C).

The correlations of FRAP with IL-10, IL-6, and VO₂max with FRAP and IL-10 are shown in Fig. 5. The analyses demonstrated a moderate positive correlation of FRAP with IL-10 (R squared 0.69, $p < 0.001$, Fig. 5A) while IL-6 showed no significant correlation with FRAP (R squared 0.40, $p < 0.001$, Fig. 5B). Additionally, we also observed a moderate positive correlation between FRAP and VO₂max (squared 0.66, $p < 0.001$, Fig. 5C), and a strong positive correlation between IL-10 and VO₂max (squared 0.82, $p < 0.001$, Fig. 5D).

Discussion

The present study was designed to compare the effects of endurance exercise training versus E2 replacement therapy on mitochondrial density, redox status, and inflammatory select biomarkers in the skeletal muscle of OVX rats. The most important finding is that both therapies exert beneficial effects on OVX rat muscles, but endurance exercise therapy was superior to E2 replacement therapy. Endurance exercise training prevented the reduction of aerobic capacity and improved mechanical efficiency. In addition, skeletal muscle mitochondrial density, CAT activity, FRAP, and IL-10 levels were higher in the exercised group. However, in OVX rats, the muscle SOD activity was higher and IL-6 levels lower only in the E2 replacement therapy.

Studies demonstrated that ovariectomy reduces rats' performance during maximal exercise tests^{21,30}. This physical performance deterioration in rats may be a consequence of the decrease in skeletal muscle mitochondrial content and function induced by ovariectomy^{4,5}.

In our study, endurance exercise training effectively prevented VO₂max reduction, improved mechanical efficiency, and promoted higher skeletal muscle mitochondrial density (an important physiological indicator of muscle mitochondrial function improvement)³¹ in OVX rats. On the other hand, E2 replacement therapy did not improve any of these parameters. It is noteworthy that ovariectomy may increase mitochondrial ROS




Figure 3. Redox status and antioxidant enzymes assays in the left soleus muscle of untrained ovariectomized rats (UN-OVX), ovariectomized rats treated with Estradiol replacement (E2-OVX), and trained ovariectomized rats (TR-OVX). (A). Thiobarbituric acid reactive substances assay (TBARS). (B). The ferric reducing ability of plasma (FRAP). (C). Superoxide dismutase activity (SOD). (D). Catalase activity (CAT). N = 10 per group. Data are reported as mean \pm S.E.M. 95% confidence interval (CI). Effect size (ES) * $p < 0.01$. One-way ANOVA.




Figure 4. Inflammatory biomarkers assays in the right soleus muscle of untrained ovariectomized rats (UN-OVX), ovariectomized rats treated with Estradiol replacement (E2-OVX), and trained ovariectomized rats (TR-OVX). (A). Tumour necrosis factor-alpha (TNF- α). (B). Interleukin 6 (IL-6). (C). Interleukin 10 (IL-10). N = 10 per group. Data are reported as mean \pm S.E.M. 95% confidence interval (CI). Effect size (ES) * $p < 0.01$. One-way ANOVA.




Figure 5. Correlations of the ferric reducing ability of plasma (FRAP) with interleukin 10 (IL-10) (A), interleukin 6 (IL-6) (B), and maximal oxygen consumption (VO_{2max}) with FRAP (C) and IL-10 (D). The open circle (white) represents untrained ovariectomized rats (UN-OVX). A closed circle (black) represents ovariectomized rats treated with Estradiol replacement (E2-OVX). A closed triangle (black) represents trained ovariectomized rats (TR-OVX).

production^{3–5}, favoring a pro-oxidant and inflammatory status leading to progressive mitochondrial dysfunction and cell death^{10,11}. To our knowledge, this is the first study highlighting the effects of endurance exercise training versus E2 replacement therapy on the skeletal muscle mitochondrial density from OVX rats using transmission electron microscopy, the "gold standard" to measure mitochondrial content. This assay exhibits more accuracy in measuring mitochondrial content than other measures, such as transcription factors levels, once changing in transcription factors levels does not necessarily induce changes in mitochondrial content.

Our results indicate that both E2 replacement therapy and endurance exercise training may protect OVX rat muscle from oxidative stress through distinct pathways. For example, while the SOD activity was higher in the E2 replacement therapy group, the CAT activity was higher in the endurance exercise training group. However, only endurance exercise training had higher total antioxidant capacity (FRAP) compared with the other OVX groups.

Our data also revealed that the skeletal muscle IL-6 levels were lower only in the E2 replacement therapy group compared to the other OVX groups. The complex physiological role of IL-6 has been a matter of debate. While some studies suggest IL-6 as an immune-modulatory cytokine that induces low-grade inflammation in some chronic diseases and a detrimental intramuscular modulatory factor in specific conditions such as cachexia, other studies suggest IL-6 as a key factor that induces positive effects on muscle metabolism and myogenesis⁹. Despite these controversial effects, we believe that the IL-6 reduction induced by E2 replacement therapy in OVX rat skeletal muscle is beneficial and contributes to an anti-inflammatory profile during menopause. Accordingly, a recent study indicated that E2 replacement therapy improved the IL-6-induced mitochondrial dysfunction through the activation of both GPER and Era receptors³². However, the mechanism behind the protective effect of E2 on mitochondrial dysfunction induced by a proinflammatory profile in OVX rats remains unknown, and a complete characterization of the modulatory role of E2 replacement therapy on intramuscular cytokines deserves future investigations.

Although endurance exercise training did not change IL-6 muscle levels in OVX rats, the TR-OVX group presented higher levels of IL-10 compared with both UN-OVX and E2-OVX groups. IL-10 is an important anti-inflammatory myokine that down-regulates pro-inflammatory signaling and protects the muscle against oxidative damage^{10,11,33}. Moreover, the positive correlation between FRAP, IL-10 and VO_{2max} reinforces the contribution of endurance exercise training to the improvement of total antioxidant capacity and anti-inflammatory

status in the OVX rat muscle. Of note, previous studies showed that IL-10 expression is coupled to the nuclear transcriptional network of mitochondrial biogenesis³⁴ and the genetic deletion for IL-10 increases damaged mitochondria in skeletal muscles, reinforcing the idea of the benefits of endurance exercise training through mitochondria-cytokine crosstalk³⁵.

The probable mechanisms underlying the improvement in skeletal muscle mitochondrial density, redox balance, and anti-inflammatory profile induced by endurance exercise training in OVX rats must be further elucidated. Furthermore, although it is plausible that there would be an additive effect of E2 replacement therapy and endurance exercise training on the evaluated parameters in our study, this experimental design is beyond the scope of the present study and deserves future investigations.

Conclusion

Overall, endurance exercise training compared to E2 replacement therapy was effective to maintain aerobic capacity and improve mechanical efficiency in skeletal muscle of OVX rats. In addition, endurance exercise training compared to E2 replacement therapy raises the skeletal muscle mitochondrial content and tends to balance the redox and inflammatory status in the skeletal muscle of OVX rats. Thus, endurance exercise training may be an alternative therapy to hormonal replacement for the treatment of musculoskeletal disabilities during menopause.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Received: 20 January 2022; Accepted: 6 May 2022

Published online: 24 May 2022

References

- Collins, K. H. *et al.* Obesity, metabolic syndrome, and musculoskeletal disease: Common inflammatory pathways suggest a central role for loss of muscle integrity. *Front. Physiol.* <https://doi.org/10.3389/fphys.2018.00112> (2018).
- Lizcano, F. & Guzmán, G. Estrogen deficiency and the origin of obesity during menopause. *Biomed. Res. Int.* <https://doi.org/10.1155/2014/757461> (2014).
- Rodrigues, C. M. *et al.* Cardioprotective effects of severe calorie restriction from birth in adult ovariectomized rats. *Life Sci.* **275**, 119411. <https://doi.org/10.1016/j.lfs.2021.119411> (2021).
- Sutham, W. *et al.* Ovariectomy and obesity have equal impact in causing mitochondrial dysfunction and impaired skeletal muscle contraction in rats. *Menopause* **25**, 1448–1458. <https://doi.org/10.1097/GME.0000000000001149> (2018).
- Capllonch-Amer, G. *et al.* Estradiol stimulates mitochondrial biogenesis and adiponectin expression in skeletal muscle. *J. Endocrinol.* **221**, 391–403. <https://doi.org/10.1530/JOE-14-0008> (2014).
- Dagdeviren, S. *et al.* Tumor necrosis factor- α antagonist administration recovers skeletal muscle dysfunction in ovariectomized rats. *J. Orthop. Res.* **29**, 275–280. <https://doi.org/10.1002/jor.21226> (2011).
- Costa, S. P. *et al.* Does endurance training prior to ovariectomy protect against myocardial contractility dysfunction in rats?. *Exp. Gerontol.* **155**, 111556. <https://doi.org/10.1016/j.exger.2021.111556> (2021).
- Kim, J. S., Jeon, J., An, J. J. & Yi, H. K. Interval running training improves age-related skeletal muscle wasting and bone loss: Experiments with ovariectomized rats. *Exp. Physiol.* **104**, 691–703. <https://doi.org/10.1113/EP087458> (2019).
- Muñoz-Cánoves, P. *et al.* Interleukin-6 myokine signaling in skeletal muscle: A double-edged sword?. *FEBS J.* **280**, 4131–4148. <https://doi.org/10.1111/febs.12338> (2013).
- Cherry, A. D. & Piantadosi, C. A. Regulation of mitochondrial biogenesis and its intersection with inflammatory responses. *Antioxidants Redox Signal.* **22**, 965–976. <https://doi.org/10.1089/ars.2014.6200> (2015).
- Valerio, A. *et al.* TNF- α downregulates eNOS expression and mitochondrial biogenesis in fat and muscle of obese rodents. *J. Clin. Invest.* **116**, 2791–2798. <https://doi.org/10.1172/JCI28570> (2006).
- Lundby, C. & Jacobs, R. A. Adaptations of skeletal muscle mitochondria to exercise training. *Exp. Physiol.* **101**, 17–22. <https://doi.org/10.1113/EP085319> (2016).
- Pedersen, B. K. Muscles and their myokines. *J. Exp. Biol.* **214**, 337–346. <https://doi.org/10.1242/jeb.048074> (2011).
- Egan, B. & Zierath, J. R. Exercise metabolism and the molecular regulation of skeletal muscle adaptation. *Cell Metab.* **17**, 162–184. <https://doi.org/10.1016/j.cmet.2012.12.012> (2013).
- Barbosa, M. R. *et al.* Resistance training and ovariectomy: Antagonistic effects in mitochondrial biogenesis markers in rat skeletal muscle. *Int. J. Sports Med.* **37**, 841–848. <https://doi.org/10.1055/s-0042-107247> (2016).
- Quinteiro, H. *et al.* Aerobic exercise training promotes additional cardiac benefits better than resistance exercise training in postmenopausal rats with diabetes. *Menopause* **22**, 534–541. <https://doi.org/10.1097/GME.0000000000000344> (2015).
- Ström, J. O., Theodorsson, E. & Theodorsson, A. Order of magnitude differences between methods for maintaining physiological 17 β -oestradiol concentrations in ovariectomized rats. *Scand. J. Clin. Lab. Investig.* **68**, 814–822. <https://doi.org/10.1080/00365510802409703> (2008).
- Yousefzadeh, N. *et al.* Review article : Ovariectomised rat model of osteoporosis. *EXCLI J.* **19**, 89–107 (2020).
- Pignon, A. *et al.* Exercise training in ovariectomized rats stimulates estrogenic-like effects on expression of genes involved in lipid accumulation and subclinical inflammation in liver. *Metabolism* **60**, 629–639. <https://doi.org/10.1016/j.metabol.2010.06.012> (2011).
- Leandro, C. G. *et al.* A program of moderate physical training for Wistar rats based on maximal oxygen consumption. *J. Strength Cond. Res.* **21**, 751–756. <https://doi.org/10.1519/00124278-200708000-00016> (2007).
- Mostarda, C. *et al.* Impact of aging on cardiac function in a female rat model of menopause: Role of autonomic control, inflammation, and oxidative stress. *Clin. Interv. Aging.* <https://doi.org/10.2147/cia.s88441> (2016).
- Brooks, G. A., Donovan, C. M. & White, T. P. Estimation of anaerobic energy production and efficiency in rats during exercise. *J. Appl. Physiol.* **56**, 520–525. <https://doi.org/10.1109/T-ED.1984.21594> (1984).
- Honorato-Sampaio, K. *et al.* Physical activity prevents alterations in mitochondrial ultrastructure and glucometabolic parameters in a high-sugar diet model. *PLoS ONE* **12**, e0172103. <https://doi.org/10.1371/journal.pone.0172103> (2017).
- Broskoy, N. T. *et al.* Skeletal muscle mitochondrial and lipid droplet content assessed with standardized grid sizes for stereology. *J. Appl. Physiol.* **115**, 765–770. <https://doi.org/10.1152/japplphysiol.00063.2013> (2013).
- Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254. <https://doi.org/10.1016/j.cj.2017.04.003> (1976).

26. Ohkawa, H., Ohishi, N. & Yagi, K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* **95**, 351–358. [https://doi.org/10.1016/0003-2697\(79\)90738-3](https://doi.org/10.1016/0003-2697(79)90738-3) (1979).
27. Benzie, I. F. & Strain, J. J. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: The FRAP assay. *Anal. Biochem.* **239**, 70–76. <https://doi.org/10.1006/abio.1996.0292> (1996).
28. Marklund, S. L. Superoxide dismutase isoenzymes in tissues and plasma from New Zealand black mice, nude mice and normal BALB/c mice. *Mutat. Res. Fundam. Mol. Mech. Mutagen.* **148**, 129–134. [https://doi.org/10.1016/0027-5107\(85\)90216-7](https://doi.org/10.1016/0027-5107(85)90216-7) (1985).
29. Aebi, H. Catalase in vitro. *Methods Enzymol.* **105**, 121–126. [https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3) (1984).
30. Conti, F. F. *et al.* Influence of sexual hormones on oxygen consumption in rats. *Rev. Bras. Med. Do Esporte.* **20**, 421–423. <https://doi.org/10.1590/1517-86922014200601665> (2014).
31. Broskey, N. T. *et al.* Exercise efficiency relates with mitochondrial content and function in older adults. *Physiol. Rep.* **3**, 1–9. <https://doi.org/10.14814/phy2.12418> (2015).
32. Bauzá-Thorbrügge, M. *et al.* Antioxidant treatment induces reductive stress associated with mitochondrial dysfunction in adipocytes. *J. Biol. Chem.* **7**, 2340–2352. <https://doi.org/10.1074/jbc.RA118.004253> (2019).
33. Dagdeviren, S. *et al.* IL-10 prevents aging-associated inflammation and insulin resistance in skeletal muscle. *FASEB J.* **31**, 701–710. <https://doi.org/10.1096/fj.201600832R> (2017).
34. Piantadosi, C. A. *et al.* Heme oxygenase-1 couples activation of mitochondrial biogenesis to anti-inflammatory cytokine expression. *J. Biol. Chem.* **18**, 16374–16385. <https://doi.org/10.1074/jbc.M110.207738> (2011) (**Epub 2011 Mar 18**).
35. Qualls, A. L., Southern, W. M. & Call, J. A. Mitochondria-cytokine crosstalk following skeletal muscle injury and disuse: A mini-review. *Am. J. Physiol. Cell Physiol.* <https://doi.org/10.1152/ajpcell.00462> (2021).

Acknowledgements

Part of this research derives from the Master’s Thesis of Sara Barros Silva. We would like to thank the Federal University of the Jequitinhonha and Mucuri Valleys (Universidade Federal dos Vales do Jequitinhonha e Mucuri) for institutional support. We also thank the National Council for Scientific and Technological Development (CNPq), the Research Support Foundation for the state of Minas Gerais (FAPEMIG), and the Coordination for the Improvement of Higher Education Personnel (CAPES—finance code 001). Besides, the assistance of the Brazilian Society of Physiology is gratefully acknowledged.

Author contributions

S.B.S.: Formal analysis, data curation, methodology, writing—review & editing—original draft. K.H.S.: Conceptualization, data curation, writing—review & editing—original draft. S.P.C.: Writing—review & editing—original draft. T.E.D.: Writing—review & editing—original draft. T.M.M.C.: Writing—review & editing—original draft. C.M.R.: Writing—review & editing—Original Draft. K.B.C.: Methodology, writing—review & editing—original draft. J.M.S.: Methodology, writing—review & editing—original draft. V.K.S.L.: Writing—review & editing—original draft. T.P.G.: Writing—review & editing—original draft. A.P.S.: Writing—review & editing—original draft. M.F.D.P.: Writing—review & editing—original draft. C.C.C.: Writing—review & editing—original draft. A.M.R.: Writing—review & editing—original draft. R.E.S.: Writing—review & editing—original draft. P.H.S.F.: Writing—review & editing—original draft. H.S.C.: Writing—review & editing—original draft. M.X.O.: Writing—review & editing—original draft. V.A.M.: Data curation, writing—review & editing—original draft. A.C.R.S.: Conceptualization, data curation, writing—review & editing—original draft.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to A.C.R.L.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022