

RESEARCH

Open Access



High smoking and low cessation rates among patients in treatment for opioid and other substance use disorders

Endre Dahlen Bjørnstad^{1,2*}, John-Kåre Vederhus¹ and Thomas Clausen^{2,3}

Abstract

Background: Smoking is a well-documented cause of health problems among individuals with substance use disorders. For patients in opioid maintenance treatment (OMT), the risk for somatic health problems, including preventable diseases associated with tobacco smoking, increases with age. Our aim was to describe smoking among patients entering substance use disorder (SUD) treatment, investigate changes in smoking from the start of treatment to 1-year follow-up, and explore factors related to smoking cessation.

Methods: We employed data from the Norwegian Cohort of Patient in Opioid Maintenance Treatment and Other Drug Treatment Study (NorComt). Participants were 335 patients entering SUD treatment at 21 participating facilities across Norway. They were interviewed at the start of treatment and at 1-year follow-up. The main outcomes were smoking and smoking cessation by treatment modality. A logistic regression identified factors associated with smoking cessation.

Results: High levels of smoking were reported at the start of treatment in both OMT (94%) and other SUD inpatient treatment patients (93%). At 1-year follow-up most patients in OMT were still smoking (87%), and the majority of the inpatients were still smoking (69%). Treatment as an inpatient was positively associated and higher age was negatively associated with smoking cessation. Most patients who quit smoking transitioned to smokeless tobacco or kept their existing smokeless habit.

Conclusion: As illustrated by the high smoking prevalence and relatively low cessation levels in our sample, an increased focus on smoking cessation for patients currently in OMT and other SUD treatment is warranted. Harm-reduction oriented smoking interventions may be relevant.

Keywords: Opioid maintenance treatment, SUD Treatment, Tobacco, Smoking cessation

*Correspondence:

Endre Dahlen Bjørnstad
endre.dahlen.bjornestad@sshf.no

¹Addiction Unit, Sørlandet Hospital HF, Po. box 416, N-4604 Kristiansand, Norway

²Norwegian Centre for Addiction Research (SERAF), University of Oslo, Kirkeveien 166, N-0407 Oslo, Norway

³Addiction Unit, Sørlandet Hospital HF, Po. box 416, N-4604 Kristiansand, Norway



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

The burden of disease associated with opioid use disorder (OUD) is substantial [1, 2]. Mortality among patients in opioid maintenance treatment (OMT) is related not only to overdose deaths [3] but also to somatic and infectious diseases that are treatable if detected and handled appropriately [4, 5]. Smoking is a well-documented cause of health problems for individuals with substance use disorders (SUDs) in general [6–8]. Individuals with opioid use disorders appear especially at risk, given established associations between opioid and nicotine use [9–11]. Research has indicated that smoking has generally been higher among individuals with substance use disorder compared with the general population [12]. Whilst prevalence of cigarette smoking in the general population has declined over the past decades [13], smoking rates appear to have remained relatively high among individuals with SUDs including OUD, both inside and outside treatment [8, 14]. Smoking cessation intervention is generally recommended for individuals with SUD [15], but also specifically for patients with OUD [7, 16]. Research has suggested that patients with OUD in treatment are motivated to cease smoking when asked [7]. In clinical practice several treatment options are, in principle, available [17]. At the same time, individuals with OUD and in OMT who are attempting to quit smoking face extra challenges, such as nicotine opioid interactions [9], complex needs [18], and a range of other individual-level challenges that may demand special attention from treatment providers [7].

Some countries have ageing OMT populations [19, 20], and this can be considered an indicator of the success of OMT [21, 22]. However, the burden of somatic health problems among individuals with OUD, including those receiving OMT, increases with age [23–25]. Older patients in OMT have reported a higher prevalence of somatic comorbidities [26, 27] and sub-clinical, self-reported impairments [28, 29]. Lifestyle-related factors such as smoking [3] that contribute to somatic diseases among ageing OMT patients [25] are therefore of concern.

As OMT populations age, the need for more descriptive information on smoking is important in the design of appropriate preventive programs.

The aim of the present study was to explore smoking rates and smoking cessation following SUD-treatment entry among patients in OMT and other SUD treatment. Specifically, we aimed to.

1. Describe smoking among patients entering SUD treatment (baseline).
2. Investigate change in smoking between start of treatment (T0) and 1-year follow-up (T1) by treatment modality, contrasting patients that entered OMT with patients that entered inpatient treatment.

3. Explore factors associated with smoking cessation in OMT and other SUD treatment.

Methods

Study design

Data were drawn from the Norwegian Cohort of Patients in Opioid Maintenance Treatment and Other Drug Treatment (NorComt) study [30]. NorComt is a longitudinal, naturalistic, multi-site study that was designed to increase understanding of factors influencing treatment adherence and outcomes for a diverse patient population across standard care treatment modalities (OMT and other inpatient SUD treatments). For this study, our primary patient group of interest comprised individuals who entered OMT. Inpatients served as a comparison treatment group, to put the results from OMT into context.

Setting

There were 21 participating facilities across Norway: 14 were OMT outpatient centers, and 7 were inpatient centers (predominately non-OMT). OMT in Norway is generally provided on an outpatient basis by publicly funded health services, following a national treatment guideline [31]. The 2010–2022 OMT guideline was in use when the data for the present study were collected. Although the specialist healthcare service serves as the overall responsible provider, the treatment is provided in collaboration with primary healthcare and social services. Apart from an established opioid use disorder diagnosis, there are no further criteria for entering this type of treatment, although substitution-free treatment is generally recommended as a first option. Inpatient SUD treatment represented in the present project typically has a duration of 6–9 months, primarily with non-OMT treatment in therapeutic community-like settings. Some patients transfer to an outpatient treatment after completing inpatient treatment. There are no specific regulations on tobacco use in Norwegian SUD treatment, other than compliance with Norwegian legislation restricting smoking inside public buildings, including health institutions. This means that, when receiving either type of treatment, patients would need to smoke outdoors, for example in a designated area in close proximity to the treatment site. The study setting has been described in more detail in previous publications [32, 33].

Participants

To participate, the only formal inclusion criterion was admittance to a SUD treatment facility, and there were no formal exclusion criteria. Participants were consecutively enrolled in the study when beginning treatment (N=548) and consented at baseline to be contacted for additional data collection one year later for a follow-up interview [30]. Clinicians at each respective treatment

center conducted the baseline interviews of consecutively enrolled patients within an average of 3 weeks from treatment initiation. The interview questions were framed to reflect the period prior to the start of treatment. Thus, “T0” refers to the start of treatment for each specific patient. There were 341 participants included at follow-up (62% of initial participants). Of these, 335 (61%) had measurements of smoking at both T0 and T1 and were included in the further analysis.

Measures

A structured interview included questions on sociodemographic variables, substance use, and a variety of measurements ranging from mental health to quality of life [32, 33]. The main outcomes were smoking and whether patients reported smoking cessation. Participants were asked both at T0 and T1 whether they had smoked cigarettes the past 6 months or not. At both assessments, they were also asked to estimate the number of cigarettes smoked per day. It was therefore possible to explore whether smoking status had changed from baseline to follow-up. Patients were also asked whether they used smokeless tobacco or not, as well as how many days one box lasted: a low number of days per box implies higher use intensity. Excerpts from the EuropASI, a validated version of the Addiction Severity Index adapted for European use [34], were used to collect data on most used substances or addictive medications. Only data on the four most used substance types were collected. The Severity of Dependence Scale (SDS) [35] was employed as a measure of psychological dependence, and as a severity measure. The SDS is a validated five-item scale that was designed to measure dependence on specific substances (e.g., “Did you think your use of heroin was out of control?”), but here the items have been rephrased to reflect general dependence on substances (e.g., “Did you think your use of substances was out of control?”). Responses were given in a 4-point format ranging from 0 to 3, with 0 corresponding to “Never” and 3 corresponding to “Always”. The summed scale ranged from 0 to 15, with higher scores representing higher severity. The Hopkins Symptom Checklist 25-item version (HSCL-25) [36] was included as a measure of mental distress [37]. The version employed in our study used a 5-point Likert-type response format [38, 39]. Each participant’s mean score (range 0–4) was used in the analysis. A score of 1.0 indicates mental distress of clinical concern [39, 40]. Somatic health-related variables were selected for analysis primarily based on their relevance to smoking. Self-reported physical health was measured by an item from the QOL10 [41], and patients were asked how they considered their physical health prior to treatment in a 5-point Likert-type format ranging from 0 to 4, with 0 corresponding to “very poor” and 4 to “very good”. Somatic

health complaints were self-reported using a structured questionnaire that included multiple complaints. Patients were asked to indicate the degree which they “had been bothered by respiratory ailments in the past 2 weeks” on a 5-point scale from 0 to 4, with 0 corresponding to “not at all”, 1 “a little”, 2 “moderately”, 3 “a lot”, and 4 “very much”.

Analysis strategy/Statistical analysis

Participant sociodemographic data, substance-use variables and health-related variables were summarized using descriptive statistics. Continuous variables are reported as means (M) and standard deviations (SD), or medians (Mdn) and interquartile ranges (IQR). Categorical variables are reported as frequencies and percentages. As we were interested in tobacco smoking by treatment modality, we tested differences in background variables at baseline between patients in OMT and inpatient treatment with t-tests or chi-square tests, depending on variable type (continuous or categorical). Categorical outcomes within the full sample and within each treatment modality were tested using McNemar’s test. Changes in continuous outcomes were tested using the matched-pairs Wilcoxon signed-rank test, as the data deviated from normal distribution. Associations between relevant independent factors and smoking cessation were investigated with bivariate logistic regression. We calculated the differences between follow-up and baseline and recoded the resulting data into a dichotomous variable where 0 was “no change” or “started smoking”, and 1 represented a change in smoking, that is, a healthy or positive change. From the bivariate analyses, variables with a p-value < 0.2 were retained for further analysis [42]. A final logistic regression was conducted to determine the strength of the associations between these variables and change in smoking. Results are presented as odds ratio (ORs) with 95% confidence interval (CI). P-values < 0.05 were considered statistically significant. Analyses were performed with IBM SPSS Statistics, version 27.

Results

Among 335 respondents included in the analysis, 175 were from the outpatient OMT group, and 160 were from the inpatient group (Table 1). At T0, patients in the OMT group reported opioids as the most used substance group (78%). Among the inpatient group, the substances most used were stimulants (38%) and cannabis (28%). Patients in both groups reported polysubstance use 4 weeks prior to treatment, with a mean above 2 drugs for both groups. Both groups had a mean score on the lower side of the self-reported physical health scale. Patients did however report only modest levels of respiratory ailments. Asthma was reported by roughly 1 in 5 in the OMT group, and by about 1 in 8 among in the inpatient

Table 1 Baseline demographics and other relevant variables of the patients that were reached at the 1-year follow-up grouped by OMT and other SUD inpatient treatment

		OMT (N = 175)	Inpa- tient (N = 160)	p-value ^a ;	All patients (N = 335)
Sociodemographics					
Age	M (SD)	39 (10)	29 (7)	< 0.001 ^b	34 (10)
Male	n (%)	127 (73)	110 (69)	0.443	237 (71)
2° education or higher	n (%)	83 (47)	65 (41)	0.230	148 (44)
Employed/education	n (%)	25 (14)	21 (13)	0.728	46 (14)
Stable living conditions	n (%)	147 (84)	111 (69)	0.002	258 (77)
Somatic health					
Asthma	n (%)	31 (18)	20 (13)	0.249	51 (15)
Respiratory ailments ^c	M (SD)	0.76 (1.16)	0.83 (1.16)	0.550	0.79 (1.16)
Physical health ^d	M (SD)	1.58 (1.05)	1.65 (1.09)	0.527	1.61 (1.07)
Mental health and other variables					
Mental distress ^e	M (SD)	1.26 (0.87)	1.22 (0.74)	0.604	1.24 (0.81)
Severity of dependence ^f	M (SD)	10.36 (3.17)	10.08 (3.06)	0.536	10.23 (3.12)

^a p-values are based on t-tests and chi-square tests
^b Equal variance not assumed
^c Item from somatic complaints questionnaire
^d Item from QOL10
^e HSCl-25
^f SDS

group. Sociodemographic variables were generally similar between the two groups. Around half had completed secondary education or higher. Only a minority reported being employed or enrolled in education. There was no significant difference between the groups' HSCl-25 scores, but both groups had means above 1.0, suggesting levels of mental distress at T0 of clinical concern. The mean SDS scores were > 10 for both groups, indicating a high level of substance use severity. The major differences between the groups were that patients in OMT were

Table 2 Smoking and tobacco use among OMT patients (N = 175) and other SUD treatment inpatients (N = 160) from treatment start to 1-year follow-up

Tobacco	OMT			Inpatient			
		T0	T1	p-value ^a ;	T0	T1	p-value ^a ;
Smoking	n (%)	165 (94)	153 (87)	< 0.001	149 (93)	110 (69)	< 0.001
Cigarettes	Mdn (IQR)	15 (10)	10 (10)	< 0.001	13 (13)	6 (13)	< 0.001
Smokeless tobacco	n (%)	47 (27)	59 (34)	0.074	66 (41)	83 (52)	0.023
Days/box	Mdn (IQR)	4 (5)	4 (5)	0.123	3 (4.75)	3 (3)	0.347

Notes: Cigarettes, daily cigarettes smoked; Days/box, days required to finish one box of smokeless tobacco

^a p-values based on McNemar's test and Wilcoxon signed-rank test

Table 3 Logistic regression of relevant covariates for smoking cessation.^a; (N = 335)

Variables	Bivariate analysis OR (95% CI)	p-value ^a ;	Multivariate analysis OR (95%CI)	p-value ^b ;
Age	0.90 (0.87–0.94)	< 0.001	0.93 (0.89–0.97)	0.002
Sex, Female	1.52 (0.84–2.77)	0.169	1.64 (0.85–3.17)	0.141
Education level	0.71 (0.40–1.26)	0.235		
Employed or in school	0.84 (0.35–1.98)	0.684		
Treatment, inpatient	5.32 (2.69–10.49)	< 0.001	2.71 (1.25–5.86)	0.011
Cigarettes per day	1.0 (0.97–1.02)	0.928		
Respiratory ailments ^d	1.11 (0.93–1.41)	0.378		
Physical health ^e	1.21 (0.93–1.59)	0.156	1.31 (0.98–1.76)	0.065
Mental distress ^f	0.82 (0.57–1.19)	0.300		
Severity of dependence ^g	0.99 (0.91–1.09)	0.866		

^a The dependent variable was dichotomized smoking change
^b p-value obtained from bivariate logistic regression
^c p-value obtained from multivariable logistic regression; multivariable analysis included variables with p-values < 0.20 in bivariate analyses
^d Item from somatic complaints questionnaire
^e Item from QOL10
^f HSCl-25
^g SDS

roughly 10 years older than the inpatients and reported more stable living conditions 4 weeks prior to start of treatment (Table 1).

In the full sample of 335 patients there was a significant ($p < 0.001$) decrease in smoking from T0 (94%) to T1 (79%). There was also a significant ($p < 0.001$) decrease in cigarette use, from a median of 15 cigarettes per day at T0 to a median of 10 cigarettes per day at T1. For smokeless tobacco, we found a significant ($p = 0.003$) increase in use from T0 (34%) to T1 (42.5%).

The within-group changes from T0 to T1 are shown in Table 2. We found a small albeit statistically significant

($p < 0.001$) decrease among the 175 OMT patients, as 12 (7%) patients reported smoking cessation, 10 (6%) continued as non-smokers, and 153 (87%) still reported smoking. Among 160 inpatients a significant ($p < 0.001$) decrease was also found, with 45 (28%) reporting smoking cessation, 5 (3%) remaining non-smokers, 6 (4%) starting to smoke, and 104 (65%) reporting that they were still smoking. Daily cigarettes smoked decreased significantly ($p < 0.001$) in both groups. A median reduction of 5 daily cigarettes was observed in patients in OMT, and among the inpatients there was a median reduction of 7 daily cigarettes.

Of the 57 subjects in the full sample who reported smoking cessation, 22 (39%) started using smokeless tobacco, 21 (37%) continued their smokeless tobacco habit, 8 (14%) quit both cigarettes and smokeless tobacco, and 6 (10%) did not use or commence using smokeless tobacco. Thus, a total of 14 (4%) former tobacco smokers reported total tobacco abstinence at follow-up. When investigating within-group smoking cessation and smokeless tobacco use, it was found that of the 12 OMT patients who reported smoking cessation at T1, 11 (92%) commenced or kept their existing smokeless tobacco habit, and 1 (0.6%) reported total tobacco abstinence. Among the 45 inpatients who reported smoking cessation, 32 (71%) either transitioned to or kept their existing smokeless tobacco habit at T1, and 13 (9%) reported total tobacco abstinence.

A logistic regression was performed to assess factors associated with smoking cessation at follow-up. Treatment modality and age were the only two variables that were significantly associated with smoking cessation (Table 3). The odds of reporting smoking cessation were nearly 3-fold higher for inpatients than for OMT-patients in the adjusted model. Higher age was negatively associated with smoking cessation.

Discussion

There was a very high prevalence of smoking among patients entering SUD treatment, and a low cessation rate 1 year after admission. Higher age and being in the OMT group (versus the inpatient group) were associated negatively with quitting smoking.

The high prevalence of smoking among patients entering treatment was observed during a period in which smoking had been on a decline for years in the Norwegian general population, to around 13% daily smokers in 2015 [13]. This is consistent with previous research that suggested smoking prevalence is still extremely high among patients with SUDs [43]. We interpret these findings to highlight the struggles with inherent challenges faced by individuals with OUD and other SUDs.

When we investigated change in smoking by treatment modality, we found that most patients in OMT were

smokers both at the start of treatment and after about 1 year of treatment. This is consistent with findings from a relatively recent systematic review on smoking prevalence in addiction treatment [8]. Among inpatients, the numbers were similar to those among OMT subjects at the start of treatment, but inpatient smoking cessation levels were higher at the 1-year follow-up. Despite the relatively low quit rate, the reported number of daily cigarettes was reduced significantly in both groups, similar to findings from previous research [44]. Although the number of patients reporting smoking cessation in OMT was relatively low, our findings suggest that it is possible for individuals in OMT to quit smoking.

When we investigated factors associated with smoking cessation, treatment modality and age were the two variables contributing significantly in the regression model. The treatment modality association may be partly explained by a difference in time of exposure to active psychosocial treatment between the out-patient OMT and inpatient SUD treatment of several months' duration. OMT is often considered life-long [31] and may be referred to as a "long-term low-intensity treatment". In the Norwegian OMT model, somatic health follow-up including smoking cessation counseling has often been organized as the responsibility of the patient's general practitioner, with overall responsibility for OMT residing with the specialist healthcare service [31]. Inpatient treatment or residential treatment, in comparison, can be characterized as "shorter-term and higher-intensity treatment". Inpatients may spend 6–9 months in an environment where goals revolve around lifestyle change as well as abstinence from substances. It may involve several social activities that increase the possibility of patients influencing each other in the recovery process. As such, the social environment in a residential treatment facility will perhaps have a stronger impact on tobacco habits than will the environment in OMT treatment, where patients relate less to other patients. We note that differences between the patients in the two treatment modalities—namely, differences in age, stability of living conditions, and primary underlying SUDs—may also have contributed to the observed differences in smoking cessation between modalities.

The age association, although modest, indicated that smoking cessation was less likely to occur as patients aged. This is concordant with findings in general population samples that have indicated that as individuals age, their intention to quit smoking decreases and their nicotine dependence increases [45]. Along similar lines, positive associations between age and being a hardcore smoker (i.e., less inclined to quit) have also been reported [46, 47]. In other words, having a long history of tobacco use and being a hardcore smoker may be confounders for age. The literature on the relationship between age and

smoking in SUD populations is sparse [48]. Nevertheless, with the increasing age of the OMT population it appears that treatment providers need to be mindful of age when considering smoking counseling and treatment, independent of where in the treatment trajectory an individual resides.

An interesting observation made relates to smokeless tobacco use. We found that most patients that quit smoking kept their existing smokeless tobacco habit or started using smokeless tobacco during the follow-up period. Smokeless tobacco has previously been described as a means of quitting for individuals already smoking in the general population [49, 50]. Tobacco abstinence would usually be the recommended outcome. However, given the difference in risk for developing health problems between smoking and smokeless tobacco, the transition from smoking to smokeless tobacco may be considered an improvement from a harm reduction perspective [51, 52].

It has previously been noted that there appears to be an unmet need for smoking cessation interventions in the addiction services [53, 54]. Our findings support this notion in the time frame of our data collection (2012–2016), especially within OMT. In 2018, after the start of the present study, standardized clinical pathways for mental health and SUD were published by the Norwegian directorate of health [55]. Improved somatic health follow-up was noted as a specific aim [55]. The aim of these standardized clinical pathways is to ensure that patients receive the same quality level of treatment and care, independent of, for example, geographical location. Even though a large part of the clinical pathways relates to quality outcomes, patient logistics, and structure, they also include recommendations and criteria of direct relevance to clinical practice and treatment outcomes. Included within the structure of the standardized clinical pathway are specific recommendations for somatic health check-ups [56], as well as recommendations on how to address smoking and smoking cessation. At the present time, not much is known about to the degree to which these clinical pathway recommendations have affected clinical practice with regard to somatic health, or if they have affected smoking cessation among patients currently enrolled in treatment. It will be of interest to researchers, decision makers, and clinicians to monitor smoking rates and cessation rates in SUD populations in the years to come, concurrent with the further implementation of the national guidelines and standardized clinical pathways.

Strengths and limitations

A strength of this study is its prospective design employing data from a relatively large clinical cohort. The data were collected through interviews, with a low prevalence

of missing data. The first interviews were conducted in a treatment setting, and the follow-up interviews were conducted by independent researchers external to the treatment process. The use of external researchers would lessen the risk of patient response bias, but likely not eliminate it completely. Some limitations should be considered when interpreting the results. With self-reported measurements there is always a chance of social desirability and recall biases. Results should thus be interpreted with caution, as self-reported smoking status was not independently verified. Furthermore, we do not have data on whether formal smoking cessation interventions and smoking cessation aids were provided as part of treatment, nor whether smoking cessation benefits were communicated. Future studies should include questions concerning smoking cessation motivation among patients in order to increase our understanding of the mechanisms underlying the cessation outcomes. We note also that, due to the inherent differences between the OMT and inpatient groups, the findings should be used to put the results from OMT into context and not as a direct comparison.

Conclusion

As illustrated by the high smoking prevalence and relatively low cessation levels in our sample, an increased focus on smoking cessation among patients currently in OMT is warranted. Harm reduction-oriented smoking interventions (e.g., smokeless tobacco) may also be of relevance and interest for future research.

List of abbreviations

EuropASI	Addiction Severity Index adapted for European use.
HSCl-25	Hopkins Symptom Checklist 25.
IQR	Interquartile range.
Mdn	Median.
NorComt	Norwegian Cohort of Patients in Opioid Maintenance Treatment and Other Drug Treatment.
OMT	Opioid maintenance treatment.
ODU	Opioid use disorder.
QOL10	Ten-item general quality of life instrument.
SD	Standard deviation.
SDS	Severity of Dependence Scale.
SUD	Substance use disorder.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12888-022-04283-6>.

Supplementary Material 1. NorComt structured interview (excerpts)

Acknowledgements

The authors wish to thank NorComt participants for their willingness to contribute to this study, and facility staff for collecting data.

Authors' contributions

EDB and JKV participated in the study design, analyzed the dataset, drafted the manuscript, and interpreted the results. TC was the NorComt's principal investigator, designed the overall study, and participated in the design of the

present study, interpretation of the results, and drafting the manuscript. All authors read and approved the final manuscript.

Funding

This study received funding from the South-East Regional Health Authority in Norway.

Data Availability

The dataset used during the current study is available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The study was approved by The Regional Committee for Research Ethics in Norway (REK 2012/1131). All methods were used in accordance with relevant guidelines and regulations. Written informed consent was obtained from all study participants.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no conflicts of interest.

Clinical trial registration

Clinicaltrials.gov no. NCT05182918. Registered 10/01/2022, i.e., the study was retrospectively registered

Received: 8 February 2022 / Accepted: 30 September 2022

Published online: 19 October 2022

References

- Strang J, Volkow ND, Degenhardt L, Hickman M, Johnson K, Koob GF, Marshall BDL, Tyndall M, Walsh SL. Opioid use disorder. *Nat Rev Dis Primers*. 2020;6(1):3.
- Degenhardt L, Charlson F, Ferrari A, Santomauro D, Erskine H, Mantilla-Herrera A, Whiteford H, Leung J, Naghavi M, Griswold M, et al. The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Psychiatry*. 2018;5(12):987–1012.
- Bech AB, Clausen T, Waal H, Šaltytė Benth J, Skeie I. Mortality and causes of death among patients with opioid use disorder receiving opioid agonist treatment: a national register study. *BMC Health Serv Res*. 2019;19(1):440.
- Larney S, Tran LT, Leung J, Santo T Jr, Santomauro D, Hickman M, Peacock A, Stockings E, Degenhardt L: All-Cause and Cause Specific Mortality Among People Using Extramedical Opioids: A Systematic Review and Meta-analysis. *JAMA Psychiatry* 2020, **77**(5):493–502.
- Troberg K, Håkansson A, Dahlman D. Self-Rated Physical Health and Unmet Healthcare Needs among Swedish Patients in Opioid Substitution Treatment. *J Addict*. 2019;2019:7942145.
- Hurt RD. Mortality Following Inpatient Addictions Treatment. *JAMA*. 1996;275(14):1097–103.
- Vlad C, Arnsten JH, Nahvi S. Achieving Smoking Cessation Among Persons with Opioid Use Disorder. *CNS Drugs*. 2020;34(4):367–87.
- Guydish J, Passalacqua E, Pagano A, Martinez C, Le T, Chun J, Tajima B, Docto L, Garina D, Delucchi K. An international systematic review of smoking prevalence in addiction treatment. *Addiction*. 2016;111(2):220–30.
- Rajabi A, Dehghani M, Shojaei A, Farjam M, Motevalian SA: Association between tobacco smoking and opioid use: A meta-analysis. *Addict Behav* 2019, **92**:225–235.
- Lichenstein SD, Zakiniaez Y, Yip SW, Garrison KA. Mechanisms and Clinical Features of Co-occurring Opioid and Nicotine Use. *Curr Addict Rep*. 2019;6(2):114–25.
- Karila L, Petit A, Zarmndini R, Coscas S, Lowenstein W, Reynaud M. [Tobacco use and illicit substance use disorders: what should we have to do?]. *Presse Med*. 2013;42(5):795–805.
- Kelly JF, Greene MC, Bergman B, Hoepfner B. Smoking cessation in the context of recovery from drug and alcohol problems: Prevalence, predictors, and cohort effects in a national U.S. sample. *Drug Alcohol Depend*. 2019;195:6–12.
- Gartner CE, Lund KE, Barendregt JJ, Mohamed Nor N, Hassan H, Vedøy TF, Kvaavik E. Projecting the future smoking prevalence in Norway. *Eur J Public Health*. 2017;27(1):139–44.
- Haas AL, Sorensen JL, Hall SM, Lin C, Delucchi K, Sporer K, Chen T. Cigarette Smoking in Opioid-Using Patients Presenting for Hospital-Based Medical Services. *Am J Addict*. 2008;17(1):65–9.
- Lien L, Bolstad I, Bramness JG. Smoking among inpatients in treatment for substance use disorders: prevalence and effect on mental health and quality of life. *BMC Psychiatry*. 2021;21(1):244.
- Morris CD, Garver-Apgar CE. Nicotine and Opioids: a Call for Co-treatment as the Standard of Care. *J Behav Health Serv Res*. 2020;47(4):601–13.
- Thurgood SL, McNeill A, Clark-Carter D, Brose LS. A Systematic Review of Smoking Cessation Interventions for Adults in Substance Abuse Treatment or Recovery. *Nicotine Tob Res*. 2016;18(5):993–1001.
- Ainscough TS, Brose LS, Strang J, McNeill A. Contingency management for tobacco smoking during opioid addiction treatment: Implementation challenges. *Drug Alcohol Rev*. 2021;40(4):658–61.
- Carew AM, Comiskey C. Treatment for opioid use and outcomes in older adults: a systematic literature review. *Drug Alcohol Depend*. 2018;182:48–57.
- Waal H, Bussestund K, Clausen T, Skeie I, Lilleveld P. Status rapport 2017: LAR 20 år - Status, vurdering og perspektiver [Status report 2017: OMT 20 years - Status, review and perspectives]. Oslo: Senter for rus- og avhengighetsforskning, Nasjonal kompetansetjeneste for tverrfaglig spesialisert rusbehandling; 2018.
- Nyhagen HT, Waal H. Den aldrende LAR pasienten [The ageing OMT patient]. *Nord Stud Alcohol Drugs*. 2017;34(1):72–9.
- Degenhardt L, Grebely J, Stone J, Hickman M, Vickerman P, Marshall BDL, Bruneau J, Altice FL, Henderson G, Rahimi-Movaghar A, et al. Global patterns of opioid use and dependence: harms to populations, interventions, and future action. *The Lancet*. 2019;394(10208):1560–79.
- Gaulen Z, Alpers SE, Carlsen SL, Nesvag S. Health and social issues among older patients in opioid maintenance treatment in Norway. *Nord Stud Alcohol Drugs*. 2017;34(1):80–90.
- Kielland KB, Skaug K, Amundsen EJ, Dalgard O. All-cause and liver-related mortality in hepatitis C infected drug users followed for 33 years: a controlled study. *J Hepatol*. 2013;58(1):31–7.
- Rosen D, Hunsaker A, Albert SM, Cornelius JR, Reynolds CF. 3rd: Characteristics and consequences of heroin use among older adults in the United States: a review of the literature, treatment implications, and recommendations for further research. *Addict Behav*. 2011;36(4):279–85.
- Firoz S, Carlson G. Characteristics and treatment outcome of older methadone-maintenance patients. *Am J Geriatr Psychiatry*. 2004;12(5):539–41.
- Lofwall MR, Brooner RK, Bigelow GE, Kindbom K, Strain EC. Characteristics of older opioid maintenance patients. *J Subst Abuse Treat*. 2005;28(3):265–72.
- Medved D, Clausen T, Bukten A, Bjørnstad R, Muller AE. Large and non-specific somatic disease burdens among ageing, long-term opioid maintenance treatment patients. *Subst Abuse Treat Prev Policy*. 2020;15(1):87.
- Han B, Polydorou S, Ferris R, Blaum CS, Ross S, McNeely J. Demographic Trends of Adults in New York City Opioid Treatment Programs—An Aging Population. *Subst Use Misuse*. 2015;50(13):1660–7.
- Muller AE, Skurtveit S, Clausen T. Building abstinence networks is an important resource in improving quality of life. *Drug Alcohol Depend*. 2017;180:431–8.
- The Norwegian Directorate of Health (Helsedirektoratet). Nasjonal retningslinje for legemiddelassistert rehabilitering ved opioidavhengighet (National guideline for medication assisted treatment for opioid dependence). Oslo: The Norwegian Directorate of Health (Helsedirektoratet); 2010.
- Muller AE, Skurtveit S, Clausen T. Many correlates of poor quality of life among substance users entering treatment are not addiction-specific. *Health Qual Life Outcomes*. 2016;14:39.
- Skjærvo I, Skurtveit S, Clausen T, Bukten A. Substance use pattern, self-control and social network are associated with crime in a substance-using population. *Drug Alcohol Rev*. 2017;36(2):245–52.
- Rivndal E, Lauritzen G. A prospective study of drug abusers in Norway: Client characteristics and treatment course. *Nord Stud Alcohol Drugs*. 2004;21(6):381–97.
- Gossop M, Darke S, Griffiths P, Hando J, Powis B, Hall W, Strang J. The Severity of Dependence Scale (SDS): psychometric properties of the SDS in English and Australian samples of heroin, cocaine and amphetamine users. *Addiction*. 1995;90(5):607–14.

36. Derogatis LR, Lipman RS, Rickels K, Uhlenhuth EH, Covi L. The Hopkins Symptom Checklist (HSCL): a self-report symptom inventory. *Behav Sci.* 1974;19(1):1–15.
37. Abel KF, Skjærø I, Ravndal E, Clausen T, Bramness JG. Perceived Self-Control is Related to Mental Distress in Patients Entering Substance Use Disorder Treatment. *Subst Use Misuse.* 2018;53(9):1454–62.
38. Ravndal E, Lauritzen G. Opiate users in methadone-assisted rehabilitation one year and two years after admission. *Tidsskr Nor Lægeforen.* 2004;3(124):329–31.
39. Andreas JB, Lauritzen G, Nordfjaern T. Co-occurrence between mental distress and poly-drug use: a ten year prospective study of patients from substance abuse treatment. *Addict Behav.* 2015;48:71–8.
40. Lauritzen G, Ravndal E, Larsson J. Gjennom 10 år - En oppfølgingsstudie av narkotikabrukere i behandling. Oslo: Statens institutt for rusmiddelforskning; 2012.
41. Ventegodt S, Andersen NJ, Merrick J. QOL10 for clinical quality-assurance and research in treatment-efficacy: Ten key questions for measuring the global quality of life, self-rated physical and mental health, and self-rated social, sexual- and working ability. *J Altern Complement Med.* 2009;1(2):113–22.
42. Altman DG. *Practical statistics for medical research.* London: Chapman and Hall/CRC; 1991.
43. Weinberger AH, Gbedemah M, Wall MM, Hasin DS, Zvolensky MJ, Goodwin RD. Cigarette use is increasing among people with illicit substance use disorders in the United States, 2002–14: emerging disparities in vulnerable populations. *Addiction.* 2018;113(4):719–28.
44. Yee A, Hoong MC, Joyce YC, Loh HS. Smoking Cessation Among Methadone-Maintained Patients: A Meta-Analysis. *Subst Use Misuse.* 2018;53(2):276–85.
45. Arancini L, Borland R, Le Grande M, Mohebbi M, Dodd S, Dean OM, Berk M, McNeill A, Fong GT, Cummings KM. Age as a predictor of quit attempts and quit success in smoking cessation: findings from the International Tobacco Control Four-Country survey (2002–14). *Addiction.* 2021;116(9):2509–20.
46. Jarvis MJ, Wardle J, Waller J, Owen L. Prevalence of hardcore smoking in England, and associated attitudes and beliefs: cross sectional study. *BMJ.* 2003;326(7398):1061.
47. Lund M, Lund KE, Kvaavik E. Hardcore smokers in Norway 1996–2009. *Nicotine Tob Res.* 2011;13(11):1132–9.
48. Weinberger AH, Funk AP, Goodwin RD. A review of epidemiologic research on smoking behavior among persons with alcohol and illicit substance use disorders. *Prev Med.* 2016;92:148–59.
49. Lund KE, McNeill A, Scheffels J. The use of snus for quitting smoking compared with medicinal products. *Nicotine Tob Res.* 2010;12(8):817–22.
50. Lund KE, Vedøy TF, Bauld L. Do never smokers make up an increasing share of snus users as cigarette smoking declines? Changes in smoking status among male snus users in Norway 2003–15. *Addiction.* 2017;112(2):340–8.
51. Maki J. The incentives created by a harm reduction approach to smoking cessation: Snus and smoking in Sweden and Finland. *Int J Drug Policy.* 2015;26(6):569–74.
52. Lund I, Lund KE. How Has the Availability of Snus Influenced Cigarette Smoking in Norway? *Int J Environ Res Public Health.* 2014;11(11):11705–17.
53. Cookson C, Strang J, Ratschen E, Sutherland G, Finch E, McNeill A. Smoking and its treatment in addiction services: clients' and staff behaviour and attitudes. *BMC Health Serv Res.* 2014;14(1):304.
54. Bowman J, Wiggers J, Colyvas K, Wye P, Walsh RA, Bartlem K. Smoking cessation among Australian methadone clients: Prevalence, characteristics and a need for action. *Drug Alcohol Rev.* 2012;31(4):507–13.
55. The Norwegian Directorate of Health (Helsedirektoratet). Nasjonal plan for implementering av pakkeforløp for psykisk helse og rus 2018–2020 (National Action Plan for the Treatment and Rehabilitation of Substances Use Disorders. Oslo: The Norwegian Directorate of Health (Helsedirektoratet); 2018.
56. The Norwegian Directorate of Health (Helsedirektoratet). Ivaretagelse av somatisk helse og levevaner ved psykiske lidelser og/eller rusmiddelproblemer. The Norwegian Directorate of Health (Helsedirektoratet); 2018.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.